



PHD

Density distribution of peripheral blood lymphocytes in health and rheumatoid disease.

Carter, S. D.

Award date:
1979

Awarding institution:
University of Bath

[Link to publication](#)

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

Copyright of this thesis rests with the author. Access is subject to the above licence, if given. If no licence is specified above, original content in this thesis is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC-ND 4.0) Licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). Any third-party copyright material present remains the property of its respective owner(s) and is licensed under its existing terms.

Take down policy

If you consider content within Bath's Research Portal to be in breach of UK law, please contact: openaccess@bath.ac.uk with the details. Your claim will be investigated and, where appropriate, the item will be removed from public view as soon as possible.

DENSITY DISTRIBUTION OF PERIPHERAL BLOOD
LYMPHOCYTES IN HEALTH AND RHEUMATOID DISEASE

Submitted by,

S. D. CARTER. B.Sc.

for the Degree of Doctor of Philosophy

at the University of Bath

1979

COPYRIGHT

"Attention is drawn to the fact that copyright of this thesis rests with its author. This copy of the thesis has been supplied on condition that anyone who consults it is understood to recognize that its copyright rests with its author and that no quotation from the thesis and no information derived from it may be published without the prior written consent of the author".

This thesis may be photocopied or lent to other libraries for the purposes of consultation.

S. D. Carter


ProQuest Number: U437523

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest U437523

Published by ProQuest LLC(2015). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code.
Microform Edition © ProQuest LLC.

ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

ACKNOWLEDGEMENTS

I would like to thank Professor R. J. Ancill for allowing me to work in his department. The constant support and wise council of my supervisors, Dr. P. A. Bacon and Dr. A. J. Collins (succeeded by Dr. N. D. Hall) has enabled this project to come to fruition.

I give thanks to the hospital administrator, Mr. C. Quinnell, for permitting me the use of the laboratory at the Royal National Hospital for Rheumatic Diseases, Bath.

The patience shown by Miss E. F. Lupton in the typing of this thesis was admirable; however, I feel sure it has put her off for life! Miss V. Winrow was most enthusiastic in collaborating in some of the experiments. My parents, Jennifer Jones and Wanda Lewcun all helped to encourage me when I most needed it and I am indebted to them. Without the courage given to me by Vic and Aillean Chester this thesis would not have been finished.

My major thanks go to the staff and patients at the "Min" who gave so freely of their blood. It wouldn't have been possible without them.

SUMMARY

The incidence, nature and significance of activated lymphocytes has been investigated in R.A. and healthy subjects.

The incidence of these cells using discontinuous Ficoll gradient analysis was studied in R.A. patients with different disease activity and in those undergoing gold therapy. The density gradient also provided a method for separating lymphocyte subpopulations for the further study of surface markers. These included the possession of surface Ig and receptors for sheep erythrocytes and IgG (Fc).

R.A. patients, particularly those with clinically active disease, had more circulatory activated lymphocytes (immunoblasts) and low density lymphocytes than healthy subjects. This increase in low density cells was not seen in patients receiving gold therapy.

Surface marker analysis showed that the population of low density (activated) lymphocytes in R.A. were enriched in surface Ig bearing cells and depleted of cells with sheep erythrocyte receptors, in comparison with the high density (small) lymphocytes. However, by comparison, high avidity Fc receptors were found predominantly on the dense lymphocytes whereas low avidity Fc receptors were more common on the low density cells.

The role of the activated cells in R.A. is considered with particular reference to the increased B cell activation in peripheral blood. The propagation of R.A. by activated lymphocytes is discussed in relation to their abnormal migratory patterns to inflammatory sites.

ACKNOWLEDGEMENTS	ii
SUMMARY	iv
<u>CHAPTER ONE</u> <u>INTRODUCTION</u>	1
SECTION 1 LEUCOCYTES	3
a) Introduction	3
b) Polymorphonuclear cells	3
i) Life History	4
ii) Motility	5
iii) Phagocytosis	5
c) Monocytes	6
SECTION 2 LYMPHOCYTES AND THE	
LYMPHATIC SYSTEM	7
a) Differences between lymphocytes and other WBC	8
i) Morphology	9
ii) Adherence	9
iii) Life span	9
iv) Mitotic potential	9
v) Traffic pathways	9
b) Lymphocyte sub-populations	10
c) Lymphocyte antigen interaction	11
d) Immunoblasts	13
i) Morphology	13
ii) Life Span	17
iii) Function	20
iv) Metabolism, markers and motility	23

SECTION 3	RHEUMATOID ARTHRITIS	30
a)	Definition	30
b)	Pathology of joint inflammation	30
i)	Synovial fluid	31
ii)	Synovial membrane	32
c)	Immunopathogenesis	34
i)	Immune reactions in joints	35
ii)	Lymphocyte sub-populations	35
iii)	Abnormal lymphocyte responses in vitro	40
iv)	Lymphocyte activation in joints	45
v)	Peripheral blood immunoblasts	46
<u>CHAPTER TWO</u>	<u>MATERIALS AND METHODS</u>	49
SECTION 1	MATERIALS	50
SECTION 2	METHODS	53
a)	Blood collection	53
b)	Lymphocyte preparation	53
i)	Methyl cellulose + carbonyl iron	53
ii)	Ficoll/Triosil and Ficoll-Paque	54
c)	Cell counts	55
d)	Cell smears	55
e)	E- rosettes and E- rosette depletion	56
f)	EA- rosettes	58
g)	EAC- rosette depletion	59
h)	Surface Ig	60
i)	Mitogen stimulation of lymphocytes	61

<u>CHAPTER THREE</u>	<u>FICOLL DENSITY GRADIENT SEPARATION</u>	
	<u>OF LYMPHOCYTES</u>	63
SECTION 1	AIMS	64
SECTION 2	INTRODUCTION	64
SECTION 3	DISCONTINUOUS FICOLL GRADIENT METHOD	79
a)	Lymphocyte preparation	79
b)	Discontinuous Ficoll gradient generation	80
c)	Monocyte depletion	82
d)	Density gradient centrifugation	82
SECTION 4	INVESTIGATION OF PARAMETERS	84
a)	Density gradient material	84
b)	Duration and force of centrifugation	88
c)	Cell load	89
d)	Rebanding of a gradient fraction	89
e)	Reproducibility	95
f)	Yield and viability	98
g)	Cytological examination of fractions	100
h)	³ H-thymidine incorporation	104
i)	Density distribution of mitogen stimulated lymphocytes	106
j)	Monocytes	110
<u>CHAPTER FOUR</u>	<u>ENUMERATION OF CIRCULATING ACTIVATED</u>	
	<u>LYMPHOCYTES IN RHEUMATOID ARTHRITIS</u>	118
SECTION 1	INTRODUCTION	119
SECTION 2	MATERIALS AND METHODS	121
a)	Patients and controls	121
b)	Lymphocyte preparation	121

c) Lymphocyte volume analysis	124
d) Density distribution analysis	125
SECTION 3 RESULTS	126
a) Immunoblast numbers	126
b) Lymphocyte diameters	129
c) Electronic sizing	129
d) Lymphocyte density distribution	130
SECTION 4 DISCUSSION	151

CHAPTER FIVE THE NATURE OF THE LYMPHOCYTE

SUB-POPULATIONS SEPARABLE BY DENSITY

IN HEALTH AND R.A. 153

SECTION 1. INTRODUCTION 154

SECTION 2. MATERIALS AND METHODS 156

a) Density distribution of enriched sub-populations 156

b) Surface markers on lymphocytes in Ficoll gradient fractions 156

i) E- rosettes 156

ii) EA- rosettes 156

iii) Surface immunoglobulin 157

SECTION 3. RESULTS 160

a) Rosette depletion 160

i) E- rosette depletion 160

ii) EAC- rosette depletion 163

b) Direct surface markers 163

i) E- rosettes 163

ii) EA- rosettes 166

iii) Surface immunoglobulin 181

SECTION 4. DISCUSSION	181
<u>CHAPTER SIX</u> <u>GENERAL DISCUSSION</u>	194
SECTION 1. LYMPHOCYTE ACTIVATION IN R.A.	195
SECTION 2. SIGNIFICANCE OF LYMPHOCYTE ACTIVATION IN R.A.	204
REFERENCES	215

CHAPTER ONE

INTRODUCTION

INTRODUCTION

This introduction will discuss the functions of leucocytes and their role in the pathogenesis of rheumatoid arthritis.

Immunoblasts and lymphocyte activation in rheumatoid disease will be highlighted.

.....

SECTION I LEUCOCYTES

a) Introduction

William Hewson, more than 200 years ago, gave the first description of the white blood cells (leucocytes). Under the light microscope, the leucocytes were identifiable as colourless refractile bodies, and because of the affinities of the nuclear and cytoplasmic materials for certain dyes of the Romanovsky type, a morphological distinction between two major cell populations was possible. Cells possessing distinct cytoplasmic granules and a round to deeply lobulated nucleus were termed granulocytes or polymorphonuclear cells; cells with sparse visible granules and a larger rounded nucleus were classified as agranulocytes or mononuclear cells.

On a histological and functional basis, the polymorphonuclear cells can be further sub-divided into three sub-populations: neutrophils, comprising 50-70% of the total leucocyte population, and eosinophils and basophils, approximately 1.4% and 0.1% respectively. Lymphocytes (20-40%) and monocytes (2-8%) constitute the remaining mononuclear cell population of the leucocytes.

b) POLYMORPHONUCLEAR CELLS

There are three types of polymorphonuclear leucocytes. The predominant type is the neutrophil, so called because of the neutral nature of their cytoplasm. Less common are the basophils, which stain with basic dyes and the eosinophils, which are acidophilic.

Granulocytes are produced from a common haematopoietic, pluripotential, stem cell in the bone marrow. The process of maturation and division requires 4-7 cell divisions (Craddock, 1972).

and a 9-10 day transit time before entering the peripheral blood.

During acute inflammation there is a loss of circulating PMNs to a defence role at the sight of local demand. This is rapidly adjusted by a compensatory discharge of reserve cells from the marrow with a stimulated proliferation of their precursors (Cartwright, Athens and Wintrobe, 1964). At the present time the factors controlling maturation and granulocyte production are poorly understood.

I. Life History

A dynamic perspective of the life history of granulocytes emerges from kinetic studies using radio-isotope labelling to monitor the rate of formation of new cells in the marrow and the duration of their brief transit (6-10 hours) in the blood before they emigrate into the extravascular tissues and body cavities. The normal history of granulocytes is a progressive journey through three compartments. i.e, from the bone marrow into the circulating blood and thence out into the tissues with no recirculation. In the steady state efflux of circulating granulocytes is matched by influx of new cells from marrow reserves. The marrow reserve of granulocytes is considerable, amounting to between 10 and 20 times the total population in blood and tissues. Granulocytes survive 4-5 days in tissues producing a total life span in the order of 10-14 days. Boggs (1967), using radiolabelling of granulocytes, calculated their blood half life at 6.7 hours, implying a renewal of the total blood population of these cells about two and a half times each day.

A minority of PMNs complete their life span in the circulation

and are removed from it by hepatic and splenic macrophages. The majority of PMNs pass into the marginating pool in venules and thence enter the extravascular tissues where, as free wandering cells, they are able to engage in the phagocytic ingestion and neutralisation of particulate antigens, such as bacteria. The PMNs disintegrate soon after this and the remains are cleared by local macrophages. Not all PMNs die in the tissues, a proportion must emigrate across epithelia for they are a normal constituent of the secretions of the lung and oral cavity.

II. Motility

Granulocytes are actively motile and the movement has been compared to the amoeboid movement of free living protozoans (Brewer, 1972). This has been tentatively explained as a result of gel-sol transformation, aided by extremely fine contractile microfilaments. It is aided by the firm adherence of granulocytes to surfaces, as readily demonstrated on glass, and these cells have been calculated to travel at 40-50 $\mu\text{m}/\text{minute}$ (Dittrich, 1962).

III. Phagocytosis

The adherence of granulocytes has been regarded by some workers as an extreme attempt to phagocytose (ingest) this surface, the cells which most readily adhere are those which are the most actively phagocytic. Phagocytosis by living granulocytes may be studied under phase contrast when bacteria are mixed with the blood sample, granulocytes extend as they move chemotactically, that is purposeful directional movement towards strains of common bacteria. After pursuit individual bacteria may be phagocytosed. In this glass slide system

granulocytes show no chemotactic interest in virulent strains of bacteria and even after chance contact fail to ingest them; adding specific antibody causes avid phagocytosis. Killing of these bacteria occurs by degranulation of the granulocytes and the release of antibacterial proteins (lyozyme and lactoferrin), alkaline phosphatase and a range of acid hydrolase enzymes. Most of these enzymes are released intracellularly, but an explosive degranulation or frequent exposure to certain antigens may cause extracellular enzyme release. Not only antigens are chemotactic, a number of serum proteins under certain conditions are also effective, notably components of the complement system.

Neither eosinophils or basophils are strongly motile or phagocytic and neither are important in the phagocytosis of bacteria. However, there is good evidence that eosinophils ingest and destroy antigen-antibody complexes (Litt, 1964), possibly preferring IgE containing complexes. Basophils are only believed to survive a few hours in tissues (Parwaresch et al, 1973).

c) MONOCYTES

Monocytes circulate for about 32 hours in the blood before emigrating through the venules to populate the peritoneum, lungs and liver where they can persist for 40-60 days (Van Furth, 1970), although they invariably develop into fixed tissue macrophages (Vernon-Roberts, 1969; Spector, 1969). Monocytes adhere most easily to glass, are actively phagocytic and have lysosomal granules which store latent hydrolytic enzymes; thus, monocytes can act in a similar manner to granulocytes in phagocytosis and degranulation. Monocytes are also chemotactically motile, but appear to have separate and distinct chemotactic factors

to polymorphonuclear cells (Spector, Walters and Willoughby, 1967). Lymphocytes stimulated by specific antigen may release a soluble factor (lymphokine) which inhibits the migration of normal macrophages. This migration inhibitory factor (MIF) is only produced in response to the specific sensitizing antigen (David et al, 1964). Lymphocytes also produce chemotactic factors which influence the behaviour of macrophages, working in harmony with MIF.

SECTION 2 LYMPHOCYTES AND THE LYMPHATIC SYSTEM

..... Lymphocytes are named from their predominance in lymph, the clear protein-rich fluid in the lymphatic system. Morphologically the latter arises in the capillary beds as blind- ending terminal lymphatics which converge and unite to form a tree-like system, ending in the thoracic and right lymphatic ducts which open into the vena cava near the heart. Functionally the lymphatic vessels provide a unidirectional, low pressure drainage pathway through which excess tissue fluid and leaked plasma protein molecules are restored to the blood. But the walls of the terminal lymphatics are more permeable than those of nearby small blood vessels, so that many wandering cells, sub-cutaneously injected drugs and, during infections, bacteria and soluble antigens also leave the connective tissues by this route. However, the contents of the peripheral lymph are monitored as it seeps through the regional lymph nodes strategically sited at the convergence of lymphatics e.g., in the mesentery and near the root of the lung, or interposed in the axis of lymphatic pathways e.g., the popliteal, anguinal and iliac nodes in the leg.

Each lymph node may be regarded as a filter screen of macrophages

through which a changing population of small lymphocytes is continually directed. In normal states these macrophages remove various non-lymphoid cells e.g., senescent granulocytes, monocytes and stray red blood cells and sometimes also particulate debris e.g., carbon dust particles from pulmonary lymph. In disease bacteria and soluble antigens leaving inflamed sites are also carried into the nodes, where macrophages and sensitised lymphocytes react with them. In the ensuing response large numbers of new lymphocytes and sometimes also of antibody-synthesising cells, are generated in the enlarged node and are dispersed via the efferent lymph.

Delivery of lymphocytes from the thoracic duct occurs on a massive scale, sufficient in the rat to renew the blood lymphocyte population between five and ten times a day. Gowans and Knight (1964) showed that, within minutes of being infused into the blood labelled lymphocytes localised in and around the walls of certain post-capillary venules (PCVs) present in the deep cortex of lymph nodes and in the Peyer patches of the intestine. Ultra-structurally, the walls of these PCVs contain numerous small lymphocytes (Marchesi and Gowans, 1964). From this grew the hypothesis that lymphocytes leaving the blood migrated through the tissues to enter the lymphatic system and thence were redirected into the blood - the concept of recirculation of lymphocytes.

a) DIFFERENCES BETWEEN LYMPHOCYTES AND OTHER WBC

In human blood, lymphocytes comprise 20-25% of the differential WBC count. Most of these are small lymphocytes about 6-9 μ m in diameter. A few are medium or large lymphocytes. The recirculating small lymphocyte differs significantly from granulocytes and monocytes in several respects.

I MORPHOLOGY

The small lymphocyte is the smallest WBC, since its dense round nucleus is enclosed by only a sparse amount of cytoplasm. This cytoplasm stains blue with Romanovsky dyes due to its slightly basophilic nature caused by the ribosomes present. Ultrastructurally, lysosomes are rare, the meagre development of the few organelles present e.g., scattered mitochondria, sparse rough endoplasmic reticulum and modest Golgi, conveys the impression of a cell which is functionally quiescent, but with proliferative potential.

II ADHERENCE AND PHAGOCYTOSIS

Small lymphocytes rarely adhere to glass and do not form pseudopodia. Movement is slow and there is no evidence of phagocytosis. Mouse lymphocytes have had directional movement measured at an average of 19 μ m/minute (Matthes, Ax and Fischer, 1971).

III LIFE SPAN

Some lymphocytes last only a few days, but others can survive for months or years in repeated recirculation.

IV MITOTIC POTENTIAL

Unlike the granulocytes the recirculating small lymphocyte is not a fully differentiated post-mitotic end cell. After recognition of antigen it divides and differentiates to mount an immune response against the antigen.

V TRAFFIC PATHWAYS

Granulocytes and monocytes have a unidirectional pathway from bone marrow through the blood to tissues. Small lymphocytes co-

exist with other WBC in these pools, but retain the ability to recirculate through their fourth compartment, the lymphatic system, and return to the blood.

b) LYMPHOCYTE SUB-POPULATIONS

Although the recirculating small lymphocytes appear morphologically identical, they are separable by difference in origin in reaction to antigen, in subsequent differentiation potential and in functional roles in the immune response.

It is generally accepted that there are two major different small lymphocyte sub-populations (Raff, 1970):-

a) T-lymphocytes, processed by, or in some way dependent on the thymus, and responsible for cell-mediated immunity.

b) B-lymphocytes, bursa-dependent and concerned in the synthesis of circulating antibody (Roitt, 1974). The bursa, discovered in chickens, has no equivalent mammalian organ, although gut associated lymphoid tissues and the bone marrow itself have been nominated as possible candidates.

T lymphocytes may be identified in vitro by anti-T antisera (Aiuti and Wigzell, 1973; Williams et al, 1973), by the binding of sheep erythrocytes in the cold (Wybran, Carr and Fudenberg, 1972; Jondal, Holm and Wigzell, 1972) and their lack of B cell markers.

The presence of integral surface immunoglobulin is often used to identify B cells (Van Furth et al, 1965; Lawton et al, 1972; Vossen and Hijams, 1975). Other markers used are surface receptors for the C3 component of complement (Ross et al, 1973). Receptors for the Fc portion of immunoglobulin (Dickler, 1976)

are possessed by B cells but some other cell types also possess them.

Recently cells with neither T or B cell markers have been described. Perlmann and Holm (1969) described K cells and Horwitz and Lobo (1975) defined L cells.

C) LYMPHOCYTE-ANTIGEN INTERACTION

Immune responses depend on the interaction of antigens with cells of both the mononuclear phagocytic system and the lymphoid system (Schrader and Feldmann, 1973; Askonas and Roelants, 1974). This often happens in the regional lymph node, the antigen arriving via the afferent lymph. There is a sudden fall in the number of lymphocytes that leave the node in the efferent lymph (Hall and Morris, 1965). Ford and Gowans (1969) suggested that this 'lymphocyte trapping' was part of the role of lymphocyte recirculation to offer to the lymph nodes a large diverse population of lymphocytes from which those with appropriate immunological potentials could be selected and retained in order to react with antigens that had been fixed by the phagocytic cells. From this general concept it is a short step to view this trapping as a specific immunological selection 'recruitment'. (Strober and Dilley 1973).

Immediately after this shut-down period, which lasts about 24 hours, lymphocytes reappear in the lymph in increased numbers so that the extent of lymphocyte recirculation may be double or treble the basal level (Hall et al, 1967). The characteristic feature of the lymph cells during this time is the presence of

large numbers of immunoblasts (Hall and Morris, 1963). These are a mixture of both T and B cells, like other lymphoid populations, depending upon the stimulus. In many experiments B cell responses predominate so it is easy to demonstrate that these lymph-borne blasts contain and secrete antibody and that many become plasma cells (Hall and Morris, 1963; Hall et al, 1967; Birbeck and Hall, 1967; Hall, 1970; Hall, Parry and Smith, 1971; Murphy et al, 1972), which are the major antibody-producing cells.

Unlike small lymphocytes the immunoblasts incorporate DNA precursors rapidly and some may be specifically cytotoxic in vitro (Denham et al, 1969). Immunoblasts produced by the caudal lymph nodes of rats can be obtained by collecting the thoracic duct lymph 3-6 days after antigenic stimulation of the hindquarter regions (Delorme et al, 1969). Because immunoblasts are released in large numbers before antibodies appear in the blood it has been suggested that their most important function is to extravasate into the tissues and so provide, by local synthesis, the antibody that the blood is unable to supply (Hall, 1969). This is supported by evidence that only a small percentage of immunoblasts (less than 5%) re-enter the lymph in rats (Hall, 1974). Some colonize lymphoid tissue in the spleen and nodes remote from the site of the localisation of antigen, and many go to the gut.

Gowans and Knight (1964) first showed that the small number of immunoblasts present in normal thoracic duct lymph 'homed' preferentially to the lamina propria of the small gut after I.V. injection into syngeneic recipients. This 'homing' is not influenced

by gut contents; it is found in the small gut of Caesarian-delivered neonates where the recipient is agammaglobulinaemic and the gut is sterile (Halstead and Hall, 1972). Parrott and Ferguson (1974) showed that a foetal gut graft under the kidney capsule, away from the influence of normally situated gut attracted a normal share of immunoblasts.

There can be little doubt that most immune responses bring about an investment of activated cells in the gut; the B blasts go to the lamina propria and become IgA-secreting plasma cells (Hall and Smith, 1970), the T blasts go to the Peyer patches (Ford, 1974). Recent evidence (Hopkins and Hall, 1976) suggests that this applies more so to blasts from gut associated lymphoid tissue (GALT).

Evidence for the localization of activated lymphoid cells in GALT has been reviewed by Hall (1974); unfortunately, attempts to demonstrate selective 'homing' of sensitized lymphocytes into delayed-type hypersensitivity (DTH) lesions, allografts or tumours have not met with conspicuous success (McCluskey, Benacerraf and McCluskey, 1963; Hall, 1967; Moore and Hall, 1973).

In humans the increased levels of immunoblasts from lymph nodes caused by antigen activation is reflected in an increase in the numbers found in peripheral blood (Crowther, Hamilton-Fairley and Sewell, 1969a).

d) IMMUNOBLASTS

I MORPHOLOGY

The first description of an atypical lymphocyte was given by Turk

in 1907; greater detail was provided by Downey and McKinlay (1923). The term immunoblast was first suggested by Dameshak (1963) and the morphological characteristics of these cells from various sources reviewed by Wood and Frenkel (1967) and Chalmers et al (1966). Stimulation of lymphocytes in vitro with both antigens and mitogens has been used to study the various stages of activation. It is apparent that both the morphology and time course of activation by antigens, mitogens and mixed lymphocytes culture are very similar in vitro (Inman and Cooper, 1965), the main differences being in the T or B cell origin of the activated cell (Asherson, Allwood and Mayhew, 1973).

The small lymphocyte measures 6-9 μ m in diameter and is characterised by its dense nucleus which occupies the large part of the cell, leaving a thin rim of clear cytoplasm. Intermediate lymphocytes measure 9-12 μ m in diameter, also have a dense nucleus sometimes with a slight nuclear cleft, and a greater amount of cytoplasm, in relation to the nucleus, than seen in small lymphocytes. Large lymphocytes (immunoblasts) range from 12-25 μ m in diameter and have a strongly basophilic cytoplasm with an eccentrically placed nucleus. The chromatin of these cells is in interphase and appears to have a combination of a stippled and a fine reticular pattern. The nucleus contains two or more nucleoli and some cells may have azurophilic granules in the cytoplasm. These characteristics are all easily seen on dry smears stained with a Romanovsky dye.

Electron microscopical examination emphasises the difference between resting and transformed cells.

The cytoplasm of the small cells contains a granular matrix which is thought to be mainly structural protein. There are a few regular-shaped mitochondria and a relatively undeveloped Golgi zone. Lysosome-like bodies are occasionally seen. The cytoplasm of the transformed cells has a similar granular matrix, but also contains large numbers of free ribosomes; high magnification has shown that polyribosomes are present in their cytoplasm. The Golgi zone is highly elaborate and irregularly-shaped mitochondria and lysosomes lie within this region. Other features seen in a number of cells include a high degree of irregularity of the outer nuclear membrane with marked extensions of this membrane into the cytoplasm and a prominent nuclear Hof .

Smetana and Potmesil (1970) have shown that resting, small lymphocytes may be characterised by the presence of ring-shaped nucleoli whilst transformed lymphocytes have larger dense nucleoli with a greater content and uniform distribution of ribonucleic basophil structures. This has been confirmed in infectious diseases (Skramkova and Fortova, 1972) and in experimental allergic encephalomyelitis in the guinea pig (Pekarek et al, 1972). Regions of the cytoplasm of some transformed cells contains fibrillar material and stain deeply with pyronin.

These characteristics are typical of the immunoblast, the largest lymphoid cell type. Other authors have disagreed as to the morphological criteria for the various atypical types and a large number of names have been used, presumably for cells in different parts of the cell cycle. These include lymphoblast, plasmablast and lymphoplasmacyte (Avrameas and Leduc, 1970).

However, to a haematologist a lymphoblast is a bone marrow-derived small lymphocyte precursor. The terms plasmablast and lymphoplasmacyte assume that the blasts are precursors of antibody forming cells.

Evidence has suggested that immunoblasts with a B cell origin have a rough endoplasmic reticulum (ER) whilst the T cell has a smooth E.R., The B cell E.R having attached clusters of polyribosomes. During the development of B cells to plasma cells, there is a more extensive E.R. with electron-dense material in the distended sacs which are arranged concentrically around the nucleus. (Birbeck and Hall, 1967).

Polliack et al (1973) and Lin et al (1973) reported that, by scanning electron microscope investigation, B cells had a villous (hairy) surface and T cells were smooth. However, it now appears that the development of a villous surface is either part of both T and B cell differentiation (Baur et al, 1975) or a reflection of lymphocyte activation. Criswell et al (1975) stimulated mouse lymphocytes with various T and B cell mitogens and found increased numbers of highly villous cells. Examination of the surface morphology of human T and B cells showed no difference between these two subpopulations and was dramatically affected by the experimental parameters and preparative techniques for SEM (Alexander and Wetzel, 1975). To date, the surface morphology of resting and activated human lymphocytes has not been investigated. However, the formation of spontaneous rosettes between human T cells and sheep erythrocytes stimulates formation of villi (Polliack et al, 1974; Lin et al, 1973; Wetzel et al, 1974). Van Ewijk et al

(1975) suggest that the presence of microvilli is more related to activation at the lymphocyte-endothelial contact point in post-capillary venules (PCVs) during lymphocyte recirculation. Microvilli also depend upon the method of cell fixation (Scott-Linthicum and Sell, 1974). Reducing the temperature of cell culture media for the bovine lymphocyte line, NBC-6 affected the size and number of microvilli. At 25°C there were a large number and at 4°C there were only a few of increased diameter (Lin, Wallach and Tsai, 1973).

17

II LIFE SPAN

Immunoblasts do not last for many hours before further development. Cell division follows and results in smaller lymphocytes (and possibly plasma cells, if of B cell origin). After antigenic stimulation of small lymphocytes in a lymph node the peak number of immunoblasts in efferent lymph is reached within 80-120 hours (Hall, Parry and Smith, 1971). At this stage 15-30% of the lymph cells are immunoblasts. Most of these appear to extravasate into the tissues (Hall, 1969; Hall and Smith, 1970). Certainly, very few, if any, recirculate from blood to lymph. This has been related to putative complementary receptors on the immunoblasts and vascular endothelia which allow their extravasation into the tissues. On the other hand the small lymphocytes may possess a receptor for the endothelia of PCVs, which allows their passage from blood to lymph (Gesner and Ginsburg, 1964).

Apart from extravasation, these immunoblasts may be removed from the circulation and destroyed by the monocyte/macrophage system (Wood and Frenkel, 1967).

A resting lymphocyte (at phase G_0 of the cell cycle) takes up to 12 hours after antigenic stimulation to enter DNA synthesis in vitro (G phase). The next stage is of increased DNA synthesis, during which the cell enlarges and is easily detected by increased uptake of DNA precursors, such as 3H -thymidine (S-phase). The increase in cell size is more related to the presence of cells in the S and $G_2 + M$ phases in the cell cycle, rather than to blast transformation per se (Steinman et al, 1978). Gunthe et al (1974) obtained information that stimulated cells in G_1 are irreversibly committed to entering S phase many hours before they actually synthesize DNA.

Reports on the duration of S-phase have been wildly different, probably due to the source of lymphocytes and the antigen used. Yoshida and Osmond (1971) reported that the S-phase lasted 13-14 hours whereas Steinman et al (1978) recorded a period of 18-24 hours using flow microfluorometry. However, maximal incorporation of tritiated thymidine takes place after 60-70 hours of mitogen stimulation (Ling, 1968), almost certainly reflecting in vitro recruitment.

Correlation of these numerous in vitro findings with in vivo events is difficult. The role of cell-cell co-operation is very important. The role of macrophages and regulator T cells in vivo has not been accurately determined in man. Polyclonal stimulation by mitogens is not always comparable to antigen stimulation, for example, maximum 3H -thymidine uptake with mitogens is after 60-70 hours of culture whereas it is 110-120 hours with antigens.

The failure of lymphocytes to proliferate continuously in vitro has often been assumed to be a reflection of poor tissue culture conditions. However, it may be that rapid maturation of the blast to an end stage cell limits proliferation. For example, Schooley's data (1961) indicate that at least 80% of mature plasma cells in situ may live less than 12 hours.

Feldbush and Stewart (1978) have also shown that B blasts do not recirculate in the rat. Large memory cells were shown to be formed in antigen draining lymph nodes and able to circulate, but not recirculate in the normal way by passing from blood to lymph.

Steinman et al (1978) suggest that not all blast cells divide to produce viable, haploid cells. Some blasts die, presumably effector cells e.g., plasma cells and cytotoxic cells, even those which divide do not produce all viable progeny.

Gowans et al (1962) and Birbeck and Hall (1967) have attempted to maintain in vivo generated immunoblasts in tissue culture and been uniformly unsuccessful, concluding that the requirements of these cells for survival and development are fairly critical and best provided by their normal physiological environment, inside a lymph node. This may be true, to some extent, but allied to the rapid maturation and the spontaneous death of blasts, the reasons for the rapid loss of these cells from the circulation are complex. Their life span in tissues has not been measured.

The life span of small lymphocytes is relatively well documented (Gowans, 1966; Parrott and de Sousa, 1971).

There are two populations of lymphocytes called short lived and long lived; the short lived consist of a population of cells, the majority of which will incorporate ^3H -thymidine within 4-5 days

whereas the long lived population consists of cells, many of which remain unlabelled over weeks or months. Rieke and Schwarz (1967) related the long lived population to T cells and the short lived population to B cells. This postulate has enjoyed general support, with some reservations (Parrott and de Sousa, 1971). Asherson, Allwood and Mayhew (1973) have shown that T blasts, produced by contact sensitivity to oxazolone, in the mouse, arise in the draining lymph nodes with the same time course as B blasts produced by antigenic challenge at lymph nodes. This confirms the work of Pritchard and Micklem (1972) and Asherson and Barnes (1973), who analysed the time course of DNA synthesis.

III FUNCTION

Immunoblasts are believed to be the effector cells of the immune response and to be the precursors of immunological memory. B blasts invariably become plasma cells and memory cells. T blasts become T effector cells and are involved in cell mediated immunity and the immunoregulation of T and B cells. Memory cells are also produced by T blasts.

Antigenically stimulated lymph nodes release immunoblasts in the efferent lymph which have been shown to contain (Hall and Morris, 1963) and to secrete (Cunningham, Smith and Mercer, 1966; English, Morris and Adams, 1976) specific antibody even though they display little organised endoplasmic reticulum (Hall et al, 1967). Immunoblasts normally extravasate into the tissues (Hall, 1969) and, certainly, some such process is necessary if effective titres of antibody are to be established in the extravascular tissue spaces (Hall et al, 1967).

These antibodies may sensitize macrophages and mast cells in the tissues rather than simply produce conventional circulating immunoglobulins. B blasts which enter the gut secrete IgA as the major immunoglobulin (Hopkins and Hall, 1976) and surface IgA before release from mesenteric lymph nodes (McWilliams et al, 1977). The immunoblasts which develop into plasma cells in the lymph or blood are believed to produce the circulating antibodies. Plasma cells, themselves do not extravasate into tissues. The nature of the lymph may have a suppressive action on the production of antibody from B blasts, but the nature of this inhibitor is so far unknown (English, Morris and Adams, 1977). Immunoblasts from a primary response undergo a fundamental physiological change to produce recirculating long lived cells involved in the secondary antibody response (Strober, 1972). Memory cells from draining lymph nodes are blast cells soon after antigenic challenge, but after many days the memory cells are all small lymphocytes. This supports the views that immunoblasts do not recirculate, that small lymphocytes do recirculate and that some B blasts divide in the circulation to provide B memory cells (Feldbush and Stewart, 1978).

The generation of T blasts heralds a more diverse range of potential effects and function being performed by a particular lymphocyte sub-set, conforming with the doctrine of clonal selection. T-immunoblasts have been demonstrated to be cytotoxic in vitro to alloantigens (Cerottini et al, 1970; Roberts and Hayry, 1977). T blasts also give rise to memory cells to alloantigens (Wilson and Nowell, 1971; Hollander et al, 1974). T blasts can also modulate the effectiveness of antibody production

by B blasts and plasma cells. (Strelkauskas et al, 1977) This modulation is demonstrated in vitro by the generation of T cells with either helper or suppressor functions in cultures, containing low or high levels of protein antigen, respectively. (Kontainen and Feldman, 1975, 1976). The first helper and suppressor cells to be detected are T blasts whereas later both help and suppression is mediated by medium and small sized lymphocytes. The blasts are proliferating to increase the number of small effector cells, but are still themselves active, particularly as suppressors (Andersson et al, 1977). T blasts may produce small lymphocytes which carry specific memory and have the ability to recirculate. (Sprent and Miller, 1972 a, b, c.).

Specific antigen stimulation of T cells causes the release of soluble, non-antibody mediators - lymphokines (Dumonde et al, 1969). To date, a large number have been reported with widely differing effects on various cells and tissues. Some even activate other lymphocytes in vitro. Immunoblasts can produce lymphokines before cell division occurs (Rocklin, 1973). Thus, antigen-activation of T cells can trigger off a whole sequence of events.

After immunization with the contact sensitizing agent Oxazolone the cells that move to sites of inflammation are predominantly T blasts and have the properties of non-specific cytotoxicity and the capacity to passively transfer cellular immunity (Asherson et al, 1973).

IV METABOLISM, MARKERS AND MOTILITY

As proliferating cells, immunoblasts have increased rates of metabolism measurable by several parameters. The more easily detectable change are increases in newly formed DNA and RNA, both easily shown with Romanovsky stains or methyl green pyronin (Elves et al, 1963).

Increases in DNA content have been demonstrated by increased incorporation of radiolabelled thymidine in lymphocytes stimulated in vitro with the mitogens phytohaemagglutinin (PHA) (Waithe et al, 1971), Concanavalin A (Con A) (Stobo et al, 1972), Pokeweed (PWM) (Froebel et al, 1975), with numerous antigens including bacterial products (Blomgren, 1975) and viral antigens (Mochizuki et al, 1977). Lymphocytes activated in vivo also incorporate ^3H -thymidine in vitro (Carter, 1965; Yoshida and Osmond, 1971). The proliferative response has been demonstrated with ^3H -thymidine and autoradiography (Schrek, 1963) or ^{14}C -thymidine and scintillation (Dutton and Eady, 1964). Flow microfluometric analysis of DNA content has been used to measure blastogenic responses (Cassidy et al, 1976).

The increased synthesis of RNA is detectable using radiolabelled uridine (Epstein and Brecher, 1965) and flow cytofluorometry (Braunstein et al, 1976).

Biochemical studies of membrane receptor sites for various activators and kinetic and inhibition studies of the early events indicate that the initial events of activation occur at the level of the lymphocyte membrane (Fischer and Mueller, 1969; Coulson, 1969). Increased turnover of membrane phospholipids (Fischer and Mueller, 1969; Lucas et al, 1971), induction of pinocytosis

and labilization of lysosomes (Hirschhorn and Hirschhorn, 1965; Hirschhorn et al, 1968), activation of an adenyl cyclase 'message' (Smith et al, 1971; Watson, 1975), increases in carbohydrate metabolism (Roos and Loos, 1973; Wang, Marquardt and Foker, 1976) and other biochemical changes, all occurring within the first hour or two of activation, are followed by events associated with derepression of eukaryote genomes (Hirschhorn et al, 1969). Ling (1968) has reviewed lymphocyte stimulation in greater detail than given here.

Apart from the internal metabolic changes there are a number of changes which are detectable on the lymphocyte surface, thus allowing analysis of individual cells and the use of definitive markers for lymphocyte activation in vivo, often before the appearance of immunoblasts.

A number of workers have analysed the T and B cell markers on blast transformed cells and found some striking differences by comparison with resting lymphocytes. Blast cells produced by PHA, Con A, pokeweed and allogeneic cell culture all had either E-rosette markers or surface Ig. B blasts lost their C_3 and Fc receptors (Jondal, 1974 a, b). However, Svedmyr, Wigzell and Jondal (1974) found that T blasts produced by sensitization to allogeneic or autologous lymphoblastoid cell lines were 'null' cells i.e., had neither surface Ig, SRBC receptors, C_3 receptors or Fc receptors, suggesting that the nature of the antigen can affect the marker expressed on the T blast. Whiteside and Rabin (1976) found that the predominantly T cell mitogens (PHA and Con A) produced E-rosetting blasts which had surface IgM. IgG and IgA and light chain determinants were only detected in PWM-induced

blasts. Hellstrom et al (1971) found that mitogenic stimulation of human T cells lead to the presence of easily detectable concentrations of a light chain. Vitetta, Forman and Kettman (1976) showed that B lymphocytes with different surface Ig markers gave rise to different responses to bacterial lipopolysaccharide (LPS), those with IgM elaborate an IgM response, those with IgD only undergo proliferation to produce numbers of memory cells.

Immature bone marrow B cells have a reduction in the number of cells with receptors for C_3 or Fc and a large number of bone marrow cells ($60\% \pm 13\%$) have been classified as null cells (Abdou, Alavi and Abdou, 1976).

Cytochemical methods have been used by Catovsky et al (1974) to differentiate T and B blasts in lymphoproliferative disorders. B leukaemias were PAS positive and T leukaemias were PAS negative whilst T leukaemias were acid phosphatase positive and the B leukaemias were negative. The PAS (Periodic acid Schiff) positivity is diastase - sensitive and therefore represents cytoplasmic glycogen content. Chronic lymphocytic leukaemias have been divided into a B or T cell origin by the presence of surface Ig on the neoplastic cell with reservations as to its use for functional studies (Knapp et al, 1974; Sagone and Murphy, 1975).

A certain proportion of T cells form E-rosettes easily - called 'active' or 'early' rosettes (Wybran and Fudenberg, 1973). Immature tonsillar T cells and MLC blasts form mainly, if not exclusively, early rosettes. Mitogens stimulate 'late' rosetting T cells to become early rosetting T cells. (Gergely, Vanky and Klein, 1976). Felsburg and Edelman (1977) showed that sensitized

lymphocytes incubated with the testing antigen increasingly formed early rosettes, thus showing another in vitro correlate for DTH as well as confirming this system as a marker for activated T cells.

Peripheral blood lymphocytes form low numbers of rosettes with autologous erythrocytes - auto-rosettes. This rosetting is a property of a high percentage of human thymocytes, and of lymphocytes treated with neuraminidase or stimulated to blast transformation by mitogens and is probably related to the low content of cell surface sialic acid (Sandilands et al, 1975).

The expression of Fc receptors (FcR) has been related to the activation of T lymphocytes, on which FcR are not usually found (Yoshida and Andersson, 1972). Cell maturation may also effect FcR expression. Miyama et al (1978) showed that precursor B cells do not express FcR. However, these are probably blasts which are plasma cell precursors. Recently, attention has centred on the function of FcR + and FcR- T blasts. Stout and Herzenberg (1975 a, b) found that helper T cells were FcR-. Van Boxel and Rosenstreich, (1974) demonstrated that FcR + cells were found as a result of antigen activation. T Blasts that become cytotoxic T cells are also FcR+ (Stout et al, 1976), but the presence of FcR on cytotoxic effector cells is equivocal. Suppressor T blasts are FcR+ (Moretta et al, 1977) and modulate the immune response by synergistic interaction with FcR - helper cells (Miyama et al, 1978). Helper T cells do not develop FcR during any stage of cooperation (Yodoi, Miyama and Masuda, , 1978).

Electrophoresis has been used to estimate lymphocyte surface charge differences. Andersson et al (1973) demonstrated that T and

B cells could be electrophoretically separated, T cells having a greater negative charge than B cells. Neuraminidase treatment of lymphocytes resulted in a disappearance of the normal biphasic distribution, suggesting that sialic acid content exerts a strong influence on the surface charge (Nordling, Andersson and Hayry, 1972). Also, the presence of membrane Ig reduces the negative charge of lymphocytes, explaining the lower electrophoretic mobility (EPM) of B cells. This surface Ig is lost by B cells triggered to antibody production (Takahasi et al, 1971) and thus these have an increased EPM. Von Boehmer et al (1974) confirmed this finding and showed that null cells also have an EPM intermediate between that of T and B cells, and these cells are FcR+. T blasts have a slower EPM than small T lymphocytes (Wiig, 1973). Some of the cytotoxic activity in an allograft is mediated by T cells that are electrokinetically blasts (Roberts and Hayry, 1977). The comparison of EPMs of activated T and B blasts has not yet been made, and although B cell activation causes an increased EPM, T cell activation has only been studied with mitogens, which decreased the EPM (Sato and Kojima, 1976).

Sensitized B cells have been shown to have specific endogenously synthesized surface immunoglobulin as antigen receptors (Julius and Herzenberg, 1974). However, the identification of specific antigen receptors on the various T cell subsets has not been successful (Elliott and Haskill, 1975; Elliott, Haskill and Axelrad, 1975), except on T blasts in mixed leucocyte cultures (Nagy et al, 1976; Krammer, 1978) suggesting that either blasts have more receptors or that in the development to memory cells the antigen receptors are masked. The theta (Θ) antigen of mouse thymocytes (Thy 1) is also believed to vary in

distribution during differentiation and maturation, the populations being separable by cell density (Shortman and Jackson, 1974). Further work to characterize the cell surface changes of T blasts has shown differences in ^{125}I -labelling patterns from those of a starting T cell population (Dunlap, Bach and Bach, 1978).

Antibody-forming immunoblasts degenerate quickly in culture, but Hall (1970) has developed a way to assess specific antibody production using bacterial adherence. Human immunoblasts produced in culture with mitogen have been used to provide evidence for membrane antigens specific for human B blasts in the 'S' phase of the cellcycle (Thomas and Phillips, 1973).

Subsets of mouse T cells have been defined by the presence or absence of surface antigens Ly 1, 2 and 3. T cells with $\text{Lyl}^+ \text{Ly23}^-$ are programmed for helper function and those with $\text{Lyl}^- \text{Ly23}^+$ for killer/suppressor function (Huber et al, 1976). Antigen stimulation of some Ly 123^+ cells gives rise to Lyl-Ly23^+ cells (Cantor, Shen and Boyse, 1976).

Thus, there are a number of ways of investigating the derivation of immunoblasts, firstly to identify a T or B cell origin, and, if T cell, to assess which subset has been stimulated.

Lymphocytes are relatively non-motile until they are activated (Wilkinson et al, 1976, 1977). The invasiveness of cancer tissue has been related to an increase of intracellular contractile proteins (Gabbiani, Trenchev and Holborow, 1975). Immunoblasts have orientated motility - chemotaxis, which is shown more by B blasts than T blasts. However, B blasts have a lower unstimulated motility than T blasts (Russell et al, 1975). This chemotaxis has been shown

to be increased towards the specific priming antigen (Wilkinson et al, 1977). The lymphocytes which can be shown most clearly to migrate into inflammatory lesions in vivo are blast-transformed cells (Moore and Hall, 1973; Asherson, Allwood and Mayhew, 1973).

SECTION RHEUMATOID ARTHRITIS

a) DEFINITION

Rheumatoid arthritis (R.A.) is a systemic disease in which an inflammatory polyarthritis in the diarthrodial joints is the predominant feature. Characteristically it involves peripheral joints (often symmetrically) and runs a sub-acute to chronic course of exacerbation and remission, depending on the degree of synovial involvement.

That joint involvement is the predominant feature is exemplified by the American Rheumatism Associations Diagnostic Criteria, in which nine of the 11 criteria refer to joint involvement (Rodnan, 1973).

The aetiology of R.A. is unknown. Glynn (1969), Bland and Phillips (1972), Barland (1973) and Denman (1975) have all put forward the hypothesis that R.A. is a disease of viral origin which manifests itself only in genetically pre-disposed people. The pathologically damaged joint which is characteristic of R.A. is due to an on-going immune response, either to the persisting virus or to antigens evoked during the initial reaction to the infective agent.

b) THE PATHOLOGY OF JOINT INFLAMMATION IN R.A.

The pathology of joint inflammation in the chronic lesion has been extensively reviewed (Gardner, 1972; Zvaifler, 1973). The synovitis of R.A. is a predominantly polymorphonuclear (PMN) exudation of the synovial fluid, and a mononuclear cell infiltration of the deep synovium. Weissman (1971) characterised the lesions in R.A. joints as follows:

- a) Margination of leucocytes and their appearance in synovial fluid.
- b) Hypertrophy and hyperplasia of synovial lining cells.
- c) Infiltration of the synovium by many lymphocytes.
- d) The transformation of synovium to granulation tissue which, as pannus, overgrows and invades cartilage.
- e) Erosion of cartilage; initially protoglycon, followed by collagen breakdown, accompanied by chondrocyte death and attempts at regeneration.

I SYNOVIAL FLUID

In chronic R.A. the volume of synovial fluid increases from 0.1 - 0.4ml to as much as 200 - 400ml. When seen at arthroscopy the fluid appears turbid, containing fibrinous clots or strands of fibrin which are most conspicuous on the synovial lining cells. The protein content increases to levels obtained for serum. Many agents capable of influencing the coagulation mechanisms are found, e.g., the Hageman Factor (Kellermeyer and Brekennage, 1966) and fibrin. Complement levels are characteristically low when compared with other joint diseases. Patients seropositive for Rheumatoid Factor (R.F) usually display high synovial fluid levels of R.F. The fluid also contains immunoglobulin (Ig) of other specificities. Many enzymes are found including proteases, lipases and collagenases. The predominant nucleated cells are PMNs, accounting for 75 - 90% of the population; lymphocytes, monocytes and occasional macrophages constituting the remainder. Cell counts vary from

2,500 to 25,000 per cubic millilitre in an exudation which shows remarkable population stability over a long time period even though the PMNs half life in synovial fluid is only four hours (Zvaifler, 1972). Hence, the joint cell turnover is very high and Hollingworth, Siegel and Creasey (1967) estimated that the daily joint cell turnover may exceed 1,000 million cells.

II SYNOVIAL MEMBRANE

Rheumatoid synovitis is characterized by a large number of histologic changes which are characteristic but not pathognomonic of this disease (Hamerman et al, 1969).

The synovial lining layer proliferates and the synovium appears oedematous and inflamed and protrudes into the joint space as slender villous projections. The lining cells are multilayered, reaching to a depth of 6-10 cells, as compared to the normal 1-3 layers. Multinucleated giant cells are common and located in areas of lining cell hyperplasia or in the sub-intimal layer. These are not unique to R.A. and have been observed in other forms of chronic synovitis (Bhan and Roy, 1971).

Oedema of the connective tissue stroma beneath the lining cells is a conspicuous feature of the early rheumatoid lesion. This contains fibrin or fibrinoid masses, occasionally interrupted by cytoplasmic processes from the synovial lining cells (Hamerman et al, 1969). The small blood vessels immediately beneath the lining cells appear to be increased in number and their lumens may be occluded by hypertrophied endothelial cells (Kulka, 1966). Schumacher and Kitridou (1972) found that the prominent vascular

changes mainly affected small venules which showed congestion and obliteration of the lumen by inflammatory cells, fibrin, platelets and enlarged endothelial cells.

Synovial cells speed up their metabolism with the onset of inflammation and Lindner (1971) suggests that this may contribute to the self-perpetuating lesion of R.A. This characteristic of involved synovial cells is one of a number of abnormalities which may persist through several generations as shown in cell culture (Gardner, 1972).

Synovial and subsynovial tissues are found to be infiltrated by many mononuclear cells. Generally, the lymphocytic infiltrate is diffuse and varies from sparse to prolific. They are thought to originate from blood and to leave through the endothelial cells of the post-capillary venules (Ziff, 1974). Within the hyperaemic, swollen synovial tissue the diffuse lymphocyte infiltrate is sometimes accompanied by a follicular arrangement and occasionally secondary germinal centres are seen. Investigations of the early lesions of the active synovitis have failed to establish the sequential order of the cellular exudation. A mixture of cell types is found in the extravascular region: small lymphocytes, undifferentiated blast cells, plasma blasts, plasma cells and macrophages, all in significant numbers. Close association of these cell types is suggestive of cell-cell interaction having a role in the prominent feature of chronic synovitis. Macrophages are found in both superficial and deep layers of the synovium (Muirden, 1966).

Granulation tissue formation at the margin of joints affected

by progressive rheumatoid arthritis is invariable and is generally regarded as a key to the peripheral, progressive destruction of articular cartilage that accounts for so many of the principal pathological features of the disease. This tissue, pannus, consists of proliferating fibroblasts, numerous small blood vessels, various numbers of inflammatory cells and occasional collagen fibres. It extends over, and takes the place of, peripheral cartilage; although invariably associated with pannus cartilage destruction can occur where there is no overlying pannus (Hamerman et al, 1967).

34

The prevailing view is that the pannus is responsible for the observed articular cartilage injury, although there is no general agreement how this comes about. Many workers support the theory that pannus is responsible for enzymatic digestion of constituents of cartilage (Hamerman, et al, 1967; Felland Thomas, 1960; Fell and Dingle, 1963). However, other authors (Hamerman et al, 1969; Tateishi et al, 1971) believe the release of enzymes from PMNs in the synovial fluid is of greater importance, suggesting that pannus is the result rather than the cause of cartilage injury, pannus formation following. Enzyme production has been shown from both chondrocytes and synovial fluid PMNs, including acid phosphatase, B- glucuronidase and collagenases (Page-Thomas, 1967, 1969; Chayen and Bitensky, 1971). These enzymes are believed to attack the cartilage proteoglycans which are leached out, making the remaining collagen fibres susceptible to mechanical stress (Holt, 1971).

c) IMMUNOPATHOGENESIS

Despite the lack of progress in identifying the cause of rheumatoid arthritis, there is considerable evidence implicating immune mechanisms.

I IMMUNE REACTIONS IN JOINTS

In the early 1960's the discovery, by Hedberg (1903) and Pekin and Zvaifler (1964), that synovial fluid haemolytic complement activity was reduced in comparison with simultaneously determined levels in the serum suggested that immune mechanisms were involved in rheumatoid synovitis. Vaughan Barnett and Sobel (1968) showed that PMNs in synovial fluid had ingested immunoglobulins which were associated with the C_3 component of complement. The association of low synovial fluid complement and seropositivity for 19S antiglobulin (R.F.) has implicated these rheumatoid factors in the immunopathogenesis of rheumatoid synovitis (Vaughan Jacox and Noell, 1968). Patients with the highest titres of serum R.F. are more efficient in utilizing complement and are the patients most likely to have vasculitis and depressed serum haemolytic complement titres (Franco and Schur, 1971). IgG antiglobulins have been noted and the resulting complexes also reduce complement levels (Britton and Schur, 1971).

The inter action of antigen and antibody initiates the complement sequence in the synovial fluid, generating a number of biologically active products. Some of those cause increased vascular permeability, allowing an influx of serum proteins and cellular blood elements into the site where the complexes reside. PMNs are attracted by complement-derived chemotactic factors and the complexes are attached to their cell surfaces by receptors for IgG and C_3 . Engulfment follows, with a concomitant release of large quantities of hydrolytic enzymes.

It is these lysosomal enzymes that are responsible for much of the inflammation and some of the tissue damage. Immune complexes alone cannot explain all the articular changes in R.A.. Of late there has been a greater appreciation of the role of cellular hypersensitivity and granulation tissues in the proliferative and destructive aspects of rheumatoid joint disease. Any theoretical explanation of rheumatoid joint disease must consider these findings.

Cryoprecipitable complement fixing immunoreactants have also been found in rheumatoid synovial fluid by Marcus and Townes (1971). Antibodies to IgG, IgM, fibrinogen, cartilage and DNA have been identified in these cryoprecipitates.

The earliest and most severe cartilage damage, however, does not occur over the broad, exposed articular surface, as one might expect, from the potentially damaging events of lysosomal enzyme release caused by immune complex ingestion occurring in the free synovial fluid, but in those areas where synovial membranes and cartilage are in close apposition. The synovial membrane contains large numbers of lymphocytes, frequently collected into aggregates or follicles, particularly around the small blood vessels, but true germinal centres are rarely seen (Gardner, 1972). The predominant cell in these nodules is the small lymphocyte, but about the periphery are typical plasma cells that, by immunofluorescent studies are found to contain immunoglobulin deposits (Fish et al, 1966).

What kind of lymphocytes infiltrate rheumatoid synovial membrane? Electron micrographs of areas rich in lymphocytes reveal predominantly small lymphocytes with less than five per cent

transformed blast-like cells and variable numbers of plasma cells (Kobayashi and Ziff, 1973). Fluoresceinated rabbit anti-human T cell antiserum applied to frozen sections of human rheumatoid synovium stains the majority of the cells within the lymphoid infiltrates, suggesting that the predominant cell is a T lymphocyte (Williams, 1975). The technique of cytoadherence, applied to the study of rheumatoid synovial membranes, has yielded conflicting data. When suspensions of sheep erythrocytes (E) or erythrocytes coated with antibody and the C_3 component of complement (EAC) are layered over frozen serial sections of rheumatoid synovium, the respective T and B markers adhere to the relevant cell in the lymphoid infiltration. Van Boxel and Paget (1975) claim that the major lymphocyte is a T cell, whereas Sheldon and Holborow (1975) concluded that B lymphocytes predominate. There is general agreement, however, that the majority of viable lymphocytes extracted from rheumatoid synovial membrane are T cells and the proportion of B cells (identified by membrane bound immunoglobulin or C_3 receptors) is usually below that of levels normally found in the blood (Van Boxel and Paget, 1975; Abrahamsen et al, 1975; Wangel and Klockars, 1977).

There is ample evidence that the rheumatoid synovium contains large amounts of immunoglobulin, much of it synthesized locally. (Smiley, Sachs and Ziff, 1968; Sliwinski and Zvaifler, 1970). Immunoglobulin G, generally found in abundance is in areas that also stain for complement components (C_3 and C_4); whereas the IgM is in lesser amounts and not identified with complement.

IgG is the predominant immunoglobulin in the synovial membrane plasma cells of both seropositive and seronegative patients and possesses in many cells, antiglobulin (IgGRF) activity (Munthe and Natvig, 1972). The authors interpret their findings to indicate that many of the plasma cells in the rheumatoid synovium make an IgG R.F. which combines with similar IgG molecules (Self associating IgG') within the cell. Although these complexes do not appear to activate or bind complement within the plasma cell cytoplasm, they might have important complement fixing activities after secretion from the cells.

II. LYMPHOCYTE SUB-POPULATIONS

A number of investigations have enumerated the T and B cells in the blood of patients with R.A., attempting to provide insights into the pathogenesis of the disease, and to find correlates of disease activity. For a variety of reasons, these studies have yielded conflicting data.

Several reports have cited normal percentages of T cells in the blood of patients with R.A. (Micheli and Bron, 1974; Tannenbaum and Schur, 1974; Zeiders et al, 1974; Brenner, Scheinberg and Cathcart, 1975; Clements et al, 1974). Williams et al (1973) noted a tendency for the proportion of T cells to decrease below normal with the development of severe disease or vasculitis. Utsinger and Bluestein (1974) demonstrated an overall decrease in the absolute number of lymphocytes, as well as a preferential decrease in T cells, associated with increases in disease activity. Froland et al (1975) found an increased percent of E rosette-forming cells in R.A. patients. Hoyeraal, Froland and Wisloff (1975) found both normal proportions and normal absolute

numbers of T lymphocytes in juvenile R.A. in a comparison with age and sex matched controls.

Most investigators have used anti-immunoglobulin antisera to assess the percent of B cells in peripheral blood. R.A. patients are generally equivalent to normal controls (Williams, et al, 1973; Zeiders et al, 1973; Froland, Natvig and Wisloff, 1975; Micheli and Bron, 1974; Tennenbaum and Schur, 1974; Utsinger and Bluestein, 1971) but an increased (Papamichael, Brown and Holborow, 1971) or decreased (Mellbye et al, 1972)

percent of B cells in peripheral blood has been reported.

In addition, several studies have noted increased numbers of null cells in patients with R.A. (Williams et al, 1973; Zeiders et al, 1973; Tannenbaum and Schur, 1974; Brenner, Scheinberg and Cathcart, 1975).

Since data from peripheral blood analysis have been of only limited usefulness, some investigators have attempted to classify lymphocytes present in synovial effusions. Again the results are conflicting. The percent of T cells was increased in some studies (Froland et al, 1975; Vernon-Roberts, Currey and Perrin, 1974). Utsinger (1975) and Brenner et al (1975) have found normal levels of T cells, a result confirming the work of Winchester et al (1973) who also found that the joints with a marked diminution of complement had increased numbers of null cells. The percent B cells has been variably reported to be increased when measured by EAC rosettes and surface Ig (Mellbye et al, 1972), normal by surface Ig (Vernon-Roberts et al, 1974) and decreased using surface Ig (Winchester et al, 1973; Brenner et al, 1975).

Lymphocytes bearing FcR are of particular interest in R.A. These have usually been detected by using erythrocytes coated with specific IgG antibody and have been called 'rheumatoid rosettes',

suggested to be characteristic of R.A. (Bach, Delrieu and Delbarre, 1970; Sany et al, 1975). However, these cells are found in normal peripheral blood (Brain and Marston, 1973; Froland, Wisloff and Michaelson, 1974). Although most B cells have FcR some FcR +ve cells do not have surface membrane immunoglobulins, complement C_3 receptors or bind to nylon fibre columns (Froland et al, 1974) and hence usually form part of the null cell fraction. Sharpin and Wilson (1977 a, b) have investigated the nature of EA- rosettes formation in R.A. and found that varying the titre of erythrocyte coating antibody showed up big differences of percent rosettes between R.A. and control.

R.A. patients formed more rosettes than healthy subjects at all concentrations of erythrocyte-coating antibody, particularly at the low concentrations. The authors explained this as being due to a substantial increase in the density or avidity of FcR on the lymphocyte surface. The binding of both antigen-antibody complexes and aggregated IgG has been extensively studied (Dickler, 1976) and increased uptake of immune complexes, as found in R.A. joints and peripheral blood, may affect their clearance or the activity of this cell population. Also, this population seems to contain the effector cells of antibody dependent cell mediated cytotoxicity (ADCC) - the K cells which may be cytotoxic for the numerous tissues for which specific antibodies are found in R.A. FcR may be important for B cell recognition of immune complexes and be required for complete antigenic stimulation and antiglobulin production (Johnson, Watkins and Holborow, 1975).

III ABNORMAL LYMPHOCYTE RESPONSES IN VITRO

In vivo cellular immunity has been evaluated using skin-test

reactions to a panel of common antigens (i.e. PPD, SK-SD, mumps, *Candida albicans*) and immunization with compounds such as DNCB or KLH. Although not unanimous, most studies of responses to memory antigens suggest an impairment of delayed hypersensitivity (Waxman et al, 1973; Bitter, 1971; Whaley et al, 1971). The reason for this disturbance is unclear. However, it has been observed that prolonged stimulation by a single antigen (i.e., in this instance the putative antigen of R.A.) can lead to poor responses to other antigens, resulting from antigenic competition (Adler, 1964).

In vitro function of rheumatoid T cells has been evaluated by mitogen - and antigen-induced blastogenesis and mediator production. Some workers have found that R.A. lymphocytes respond normally to PHA (Reynolds and Abdou, 1973; Paty et al, 1974; Georgescu, Georgiu and Gociu, 1975), but the majority of evidence seems to support diminished responses to PHA (Ivanyi, Lehner and Burry, 1973; Rawson and Huang, 1974 a; Froebel et al, 1975; Lockshin et al, 1975; Rawson, 1975; Clot et al, 1975; Hepburn, McDuffie and Ritts, 1976; Silverman et al, 1976; Menard, Dion and Richard, 1977). Responses to Con A have been reported as being both unchanged (Rawson and Huang, 1974b) and diminished (Clot et al, 1975; Lockshin et al, 1975; Silverman et al, 1976) in R.A. Responses to PWM (transforms predominantly B cells) are usually reported as being the same in R.A. and healthy controls (Ivanyi et al, 1973; Silverman et al, 1976).

The MLC test was developed as an in vitro model for the recognition phase of the homograft reaction. R.A. lymphocytes usually respond and stimulate appropriately in an MLC reaction with normal lymphocytes (Astorga and Williams, 1969; Stastny, 1974). The lymphocyte surface antigens are coded for by genes in the major

histocompatibility complex (MHC) which also code for HLA antigens.

A number of human diseases are correlated with an increased frequency of HLA antigens; the most notable in rheumatic diseases being the association between ankylosing spondylitis and HLA B-27. Recently, an association has been found between R.A. and HLA-Dw4 (Stastny, 1978) and related to patients with a family history of R.A. (Wooley, Panayi and Batchelor, 1978) and serum RF levels (Roitt et al, 1978).

Due to the difficulty of assaying immunoglobulin production by antigenic stimulation of human B cells in vitro, attempts to identify B cell defects can only be explored indirectly. Most studies are concerned with circulating immunoglobulin levels or antibody responses to immunization. However, deficiencies or excesses may be due to T cell abnormalities in the modulation of antibody production. Total serum Ig levels in patients with R.A. are increased (Barden, Mullina and Waller, 1967; Bandilla, Pitto and McDuffie, 1970), but this may well represent the non-specific stimulatory effect of chronic inflammation, rather than specific hyperreactivity of the humoral immune system. Naturally occurring antibodies, such as the isohaemagglutinins (anti-A and anti-B) are not elevated in rheumatoids, but possibly decreased (Rawson, Abelson and McCarty, 1961). Antibody responses to exogenous antigens, such as Brucella abortus or tetanus toxoid are generally reported as being normal in R.A. (Waller, Ellman and Toone, 1966; Rhodes et al, 1969), but in Bristol Bucknall et al (1978)

found decreased response, both 1° and 2°, to Ø x, an antigenic bacteriophage not previously met by normal subjects. On the other hand, the response to endogenous antigens (i.e., autoantibodies to

self antigens such as thyroglobulin) appears to exceed that of age - and - sex - matched controls (Muller, 1971).

The role of helper or suppressor cells in R.A. has not been evaluated, the lack of human assay systems being the main reason, particularly for suppressor cells. Also the role of such cells in vivo could only be guessed at.

Recently the cytotoxic potential of rheumatoid lymphocytes has been investigated. PHA induced cytotoxicity is reduced in R.A. (Wisloff, Froland and Natvig, 1975), this may be related to the reduced thymidine incorporation of R.A. lymphocytes cultured with PHA. Increased ADCMC has been reported for R.A. lymphocytes and related to FcR bearing cells (Sany et al, 1977). However, Diaz-Jouanen et al (1976) and Panayi and Corrigan (1977) reported unchanged ADCMC in R.A.. Isturiz et al (1976) found decreased ADCMC in R.A. The possibility of cytotoxicity towards an antigen in the rheumatoid synovium has been investigated and been found to be increased (Person, Sharp and Lidsky, 1976) and normal (Griffiths et al, 1976) in R.A. Panayi, (1976) has found increased cytotoxicity towards Chang human liver cells in R.A. and suggested a possible antigenic similarity between rheumatoid synovial membrane and Chang cells.

The identification of an antigen (G) responsible for the intra-articular immune response is obviously one of the most important steps in solving the mystery of R.A. Many animal models have been used in those studies to assay the effects of a range of antigens (Cooke and Jasin, 1972). These studies have a limited usefulness, but they have shown that antigen persistence is not essential for the maintenance of chronicity

in R.A., the initial antigen possibly inducing the formation of a second antigen that persists in the absence of the first. R.A. lymphocytes produce MIF if cultured with aggregated or complexed IgG (Eibl and Sitko, 1975). Rheumatoid factor has mitogenic properties in inducing lymphocyte transformation (King, Messner and Williams, 1968). Antigen-antibody complexes also stimulate lymphocytes, particularly in the presence of complement, (Moller, 1969; Bloch-Schtacher et al, 1968) and may be involved in maintenance of the inflammation by providing an altered IgG structure which acts as an autoimmunogen.

..... To counteract the involvement of lymphocytes in R.A. a number of drug treatments have been tried. Glucocorticoids may work by being immunosuppressive and cytotoxics, such as azathioprine and cyclophosphamide have been used with some success (Bacon et al, 1978). Antilymphocyte serum has been used in systemic lupus erythematosus (SLE), (Nossa et al, 1969). Drainage of the thoracic duct lymph has been used as immunosuppressive treatment in glomerulonephritis (Bonomini et al, 1970), myasthenia gravis (Bergstrom et al, 1975), multiple sclerosis (Brendel, Ring and Seifert, 1975) and in preparation for allotransplantation. Thoracic duct drainage (TDD) of R.A. patients has been used by Paulus et al (1977) and produced significant clinical improvement, as shown by a number of parameters, supporting the hypothesis that some lymphocytes in thoracic duct lymph are essential for the continued activity of the inflammation associated with rheumatoid arthritis. 51-Chromium-tagged reinforced autologous lymphocytes were demonstrated in the inflamed synovium and synovial fluid and had a slower rate of disappearance from these sites than from the liver or spleen. If

'homing' was occurring only a few cells were participating. Edgren et al (1976) found that extracorporeal irradiation of thoracic duct lymph in R.A. patients produced a marked clinical improvement and found that peripheral blood T lymphocytes were reduced after reinfusion of the irradiated lymph. Indeed, both groups of workers suggest that removal of thoracic duct lymphocytes may cause depletion of suppressor T cells.

IV LYMPHOCYTE ACTIVATION IN JOINTS

The lymphocyte functions described above depend upon certain lymphocytes being activated at one time or another. Some of them will be activated in joints and others, possibly, in lymph nodes, spleen or peripheral blood. Very few studies have been undertaken, into the occurrence and nature of activated lymphocytes in the rheumatoid patients. Loewi, Dorling and Howard (1974) found increased numbers of large lymphocytes in R.A. synovial fluid; plasma cells were also found and it was suggested that these account for the increased numbers of null cells found in R.A. synovial fluid. Traycoff, Pascual and Schumacher (1978) found lymphoblasts in R.A. synovial fluid but rarely in synovial fluid from patients with crystal-induced synovitis or bacterial infections. Patients with ankylosing spondylitis and psoriatic arthritis had small numbers of lymphoblasts which were non-T cells. Both groups suggested that the identification of lymphoblasts in synovial fluid may be a useful marker for rheumatoid arthritis. Electron microscopic examination of perivascular infiltrates of the rheumatoid synovium showed large numbers of plasma cells grouped together in certain areas and transitional areas containing both plasma cells, immunoblasts

and small lymphocytes (Ishikawa and Ziff, 1976). Most of the immunoblasts contained rough endoplasmic reticulum, well-developed Golgi areas and mixtures of mono - and polyribosomes in the cytoplasm, indicating that they were plasmablasts (i.e., destined to become plasma cells). About a third of the blasts had T blast characteristics. Macrophages were seen to be intimately associated with both T and B blasts suggesting release of mediators of cell mediated immunity and cell-cell interaction. The authors suggested that plasma cells were common in R.A. synovium because of their relative immobility compared with recirculating T cells. So, although most of the small lymphocytes in R.A. synovium are T cells, most of the blasts are of B cell origin. Isolation of lymphoid cells from rheumatoid synovium by de Vries et al (1977) showed that a minority of blasts (30-40%) formed E-rosettes, but that no blasts formed EAC-rosettes and interpreted this as a lack of B-blasts in rheumatoid synovium. Their assumption that B blasts retain C_3 receptors has been shown to be incorrect (Jondal, 1974 a,b) and thus many B blasts may be present due to the low number of T blasts shown.

V. PERIPHERAL BLOOD IMMUNOBLASTS

The demonstration of immunoblasts in rheumatoid synovium does not mean that the lymphocytes are activated and/or transformed there. Horwitz, Stastny and Ziff (1970) found increased numbers of proliferating lymphoid cells in the peripheral blood of patients with inflammatory diseases, including rheumatoid arthritis. Increased thymidine incorporation was found in acute polyarthritis,

the highest levels being in acute juvenile R.A. The authors were not sure if these cells were haematopoietic stem cells or lymphocytes stimulated to divide by a specific antigen.

Bacon, Sewell and Crowther (1975) showed that these proliferating mononuclear cells were immunoblasts and were raised in R.A.

by comparison with healthy subjects, particularly in early R.A.

and in acute exacerbation of disease activity. These cells were identical to those found in idiopathic thrombocytopenic purpura, other auto-immune disorders, infections and following immunization.

The authors suggested that the circulating immunoblasts in R.A.

may account for the changes in lymphocyte function and sub-populations

and may be related to events within the synovial membrane. Delbarre,

Le Go and Kahan (1975) also found increased numbers of immunoblasts

in R.A. and again found a positive correlation between numbers and

disease activity. Immunoblast numbers were not related to

inflammation, patients with psoriatic arthropathy having normal immuno-

blast levels. Increased numbers of immunoblasts have also been found

in polymyalgia rheumatica (Eghtedari, Esselinckx and Bacon, 1976) and

ankylosing spondylitis (Eghtedari, Davis and Bacon, 1976).

The immunoblast may have a central role in the pathogenesis of rheumatoid arthritis. It may be responsible for the spread of the disease or merely a non-specific response to immune complexes or a host of other functions between these two. Once the occurrence and nature of immunoblasts has been determined more work can be carried out to find out the origin and fate of these cells.

The work described was carried out with the aim of enumerating immunoblasts in R.A. and to characterize them by

means of surface marker techniques with a view to investigating their role in the disease pathogenesis.

CHAPTER TWO

MATERIALS AND METHODS

SECTION 1MATERIALS

All chemicals and reagents were of analytical grade whenever available and, unless otherwise stated, came from normal commercial sources such as British Drug Houses Ltd., Poole, Dorset.

All tissue culture media and fluorescein conjugated anti-Ig were obtained from Wellcome Reagents Ltd, Beckenham, Kent or Flow Laboratories, Irvine, Scotland.

..... All Sephadex and Sepharose gels, Ficoll and Ficoll/Paque were from Pharmacia, Uppsala, Sweden.

Radiolabelled (^3H -) thymidine was from the Radiochemical Centre, Amersham, Berks.

Preservative-free Heparin was from Evans Medical Ltd., Liverpool.

Physiological solutions and tissue culture media1. Phosphate buffered saline (P.B.S.) - DulbeccoDulbecco AFormula:

	<u>mg/litre</u>
Sodium chloride	8000
Potassium chloride	200
Disodium Hydrogen phosphate	1150
Potassium Di-hydrogen phosphate	200
in distilled water (D.W.)	

Dulbecco BFormula:

	<u>mg/litre</u>
Calcium chloride	100
Magnesium chloride	100
in distilled water (D.W.)	

Five ml of Dulbecco B was slowly added, with stirring to 1 litre of Dulbecco A to make the complete Dulbecco phosphate buffered saline (pH 7.4) which was filter sterilised into sterile glass bottles and stored at +4°C.

2. TC 199Formula:

TC199 (10X concentrate)	100 ml
Sterile water	830 ml
penicillin/streptomycin	20 ml
4.4% Sodium bicarbonate	C.50 ml

Sodium bicarbonate varied to adjust to pH 7.4 at 20°C.

Filter sterilised into sterile glass bottles. Stored at +4°C.

3. TC199 + 10% new born calf serum

Formula:

TC199 (10 X concentrate)	100 ml
Heat inactivated new born calf serum	100 ml
Penicillin/streptomycin (5000 iu/ml and 5000 mcg/ml)	20 ml
Glutamine (200 mM)	10 ml
Distilled water	720 ml
4.4% Sodium bicarbonate	C. 50 ml

pH adjusted to 7.4 with sodium bicarbonate. Filter
sterilised into sterile glass bottles and stored at +4°C.

SECTION 2

METHODSa. Venous Blood - Defibrination and anti-coagulation

Blood was drawn from a **cu** bital vein into a sterile disposable syringe and immediately expelled into 10 ml sterile bottles, each containing 300 iu heparin (preservative-free) and gently shaken for one minute. Alternatively, the blood was expelled into 20 ml bottles, each containing two large steel **paper** clips and gently, but constantly, shaken until the blood was completely defibrinated. The fibrin coated paper clips were then removed from the bottle.

b. Lymphocyte preparationsi) Methyl Cellulose and Carbonyl Iron

Materials 1% Methyl Cellulose (BDH) in 0.9% saline.

Carbonyl Iron (BDH)

Method

20 ml blood was mixed with 2.5ml methyl cellulose solution and 50 mg carbonyl iron in a 30 ml McCartney bottle. This was shaken gently at 37°C for 30 minutes.

The bottle was then allowed to stand on a strong magnet at 37°C for 30 minutes. During this period the erythrocytes were clumped by the methyl cellulose and sedimented and the polymorphonuclear leucocytes and monocytes, which had phagocytosed the iron, were pulled to the bottom of the bottle by the magnetic field.

The lymphocyte rich supernatant was aspirated with a pasteur pipette and collected in glass centrifuge tubes.

ii) Ficoll/Triosil and Ficoll-Paque

Materials Ficoll/Triosil - 10 ml 9% Ficoll in distilled water, 4ml 50% Triosil in distilled water mixed, filter-sterilised and stored at -20°C .

Ficoll-Paque - (Pharmacia)

Method

20 ml of blood was diluted 1:1 with PBS, mixed and carefully layered on to Ficoll/Triosil in glass centrifuge tubes at the ratio of 6ml of diluted blood to 4ml of Ficoll/Triosil. The tubes were then centrifuged at 400g maximum at 18°C for 30 minutes. During this period the Ficoll caused the erythrocytes to agglutinate and sediment, the Triosil was phagocytosed by the dense polymorphonuclear cells, which increased in density and sedimented. The interface between the upper plasma layer and the lower Ficoll/Triosil compartment became a site for accumulation of lymphocytes and some monocytes.

Taking care to avoid contamination with Ficoll/Triosil the lymphocyte rich interface was carefully removed and ejected into a glass centrifuge tube.

On some occasions the lymphocyte rich supernatant from the methyl cellulose-carbonyl iron method was further purified on Ficoll/Triosil.

Ficoll-Paque was sometimes used instead of Ficoll/Triosil and the cell yield and contamination levels were similar enough to allow interchange of the two media as availability required.

c. Measurement of total cells and viability

Materials White cell counting fluid (WCCF) comprising:

0.1% Methylene Blue	} in distilled water
3.0% acetic acid	
Trypan Blue - 0.16% in PBS	

For total cell count an aliquot of the cell suspension was diluted with WCCF which lysed the red cells and stained the leucocytes a pale blue, and the concentration microscopically determined in a counting chamber.

..... To assess viability an aliquot of the cell suspension was diluted in Trypan blue stain and viability determined in a counting chamber. According to Boyse, Old and Chourovulinkov (1964) viable cells exclude the stain but non-viable cells cannot and stain dark blue. Hence, percentage viability may be recorded. Other dyes have been used, such as Eosin Y (Terasaki, Vredevoe and Mickey, 1967) and nigrosin (Kaltenbach, Kaltenbach and Lyons, 1958). Trypan blue is not a sensitive test of damage of lymphocytes (Tennant, 1964); cells that appear under phase contrast to be in an advanced state of degeneration may nevertheless exclude the dye. Although such cells may be alive at the time of testing there can be no confidence in their prolonged survival.

d. Cell smears

Permanent preparations of cell suspensions were made by smearing one drop of concentrated cell suspension on one glass slide with another of lesser width, to reduce surface tension effects.

In some instances a Shandon cytocentrifuge was used to prepare permanent slides.

All slides were left to air dry (≥ 10 mins), fixed by flooding

with methyl alcohol and then, when dry, stained. Most preparations were stained with Wright's stain (0.2% in methyl alcohol) in the following manner: the slide was flooded with Wright's stain for two minutes. After this period the stain was diluted 1:1 with a phosphate buffer (pH 5.8), and left to differentiate for eight minutes. The stain was washed off the slide with the phosphate buffer and left to air dry prior to microscopic examination.

e. E- rosettes

Materials Sheep Red blood cells (SRBC) (in Alsevers solution). (Oxoid-fresh weekly)
SRBC absorbed heat inactivated new-born calf serum.
Preparation: 1ml packed washed (X3) SRBC mixed with 10ml heat inactivated new-born calf serum.
Left overnight at 4°C. Centrifuged at 200g for 10 minutes. Supernatant aspirated and stored at -20°C in 1 ml aliquots.

Method

One ml sheep blood was mixed with a large excess of PBS with a pasteur pipette and centrifuged at 200g for 10 minutes to pellet the cells. The cells were washed twice by resuspension in PBS and centrifugation. The supernatant was carefully aspirated with a pasteur pipette and 50 µl packed cells removed, mixed with 1 ml SRBC-absorbed heat inactivated new-born calf serum and made up to 10 ml with PBS.

Lymphocytes from Ficoll/Triosil preparations were washed

three times with PBS by repeated sedimentation and resuspension in fresh media, and adjusted to 2×10^6 /ml.

Equal volumes (0.2ml) of lymphocyte suspension and SRBC suspension were mixed in small, stoppered, plastic, round bottomed tubes (LP3 Luckham). Immediate incubation at 37°C for 10 minutes was followed by centrifugation at 200g for four minutes. Refrigeration at 4°C was maintained for two hours. During this period a 0.2% crystal violet solution in PBS was prepared and filtered through cotton wool. Three drops of crystal violet dye were added to each tube and the contents resuspended in a controlled manner on a rotary turntable (20 revolutions/min - Baird and Tatlock) for one minute. A drop of the resuspended cells was placed in a counting chamber (Neubauer) and the % rosetting lymphocytes determined. The lymphocytes stain dark blue and a rosette was counted as a lymphocyte binding three or more SRBC.

E- rosette depletion

To deplete populations of E- rosetting lymphocytes the rosettes were formed as above, but mixing 10 ml volumes of both lymphocytes and SRBC instead of the 0.2 ml aliquots. The incubation and centrifugation steps were as described. The twenty ml. of lymphocytes plus SRBC was layered on to 4 X 4ml volumes of Ficoll/Triosil and centrifuged at 400g for 30 minutes, at 18°C . During this period the SRBC sedimented to the bottom of the tube, and any lymphocytes bound to the SRBC sedimented with them. Hence, the interface between the upper plasma layer and the lower Ficoll/Triosil compartment contained lymphocytes depleted of E- rosetting cells. This interface was aspirated with a pasteur pipette, ejected into a glass centrifuge tube and washed three times with PBS.

f. EA- rosettes

Materials Ox blood -fresh fortnightly courtesy of
Rabbit anti **ox** RBC antiserum Dr. J. Smith,
Regional Immunology
Service, Southampton.

The IgG fraction of the rabbit anti-ox RBC antiserum was purified by affinity chromatography on Protein A-Sepharose CL-4B (Pharmacia). The F_c portion of IgG specifically interacts with Protein A (Forsgren and Sjöquist, 1966).

The gel was poured into a small chromatography column (K9 Pharmacia) and washed with PBS. Rabbit anti-ox RBC antiserum (2.5ml) was loaded on to the column and eluted with PBS to remove all the serum proteins except IgG. The bound IgG was eluted with 1M acetic acid and the eluate concentrated to 2.0ml over an Amicon Diaflo PM10 membrane (nominal exclusion weight 10,000 daltons) at 4°C.

The protein content of the IgG fraction was assessed using the Folin- Ciocalteu method (Lowry et al, 1951) and the concentration adjusted to the normal concentration found in serum (8-16mg/ml).

Method

Ox blood (0.5 ml) was washed three times with PBS, pelleted, the supernatant carefully removed by aspiration and discarded. 100µl packed cells were made up to 5ml with PBS and mixed. Equal volumes (1 ml) of ox RBC and rabbit IgG anti-ox RBC antiserum, at varying dilutions in PBS, were mixed and incubated at 37°C for 60 minutes. The cells were sedimented at 200g, washed three times with PBS and made

up to 2 ml with PBS , resulting in a 1% suspension of IgG - coated ox RBC.

59

During this period lymphocytes were prepared by isopycnic centrifugation on ~~Ficoll~~/Ficoll/Trisil, washed three times with CMFSS and adjusted to 2×10^6 /ml.

Equal volumes (200 μ l) of 1% IgG coated ox RBC and lymphocytes were mixed in LP3 tubes, centrifuged at 200g for four minutes and incubated at 4°C for one hour.

After incubation the rosettes were stained with crystal violet, resuspended and % rosetting lymphocytes determined as for E- rosettes. (Section 2.e)

g. EAC- rosette depletion

This is the depletion of lymphocytes with receptors for the C3 component of complement.

Materials Sheep blood in Alsevers Solution

Rabbit anti-SRBC (Flow Ltd.) diluted in PBS

Human serum

Method

1ml packed washed SRBC was mixed with 5ml 1/10,000 rabbit anti-SRBC and incubated for one hour at 37°C. The SRBC were pelleted and washed three times with PBS. After the third sedimentation the supernatant was discarded and 0.5ml human serum added as a source of complement, incubation at 37°C for 30 minutes was followed by three washes with PBS. 50 μ l of packed cells were made up to 10 ml in PBS.

Lymphocytes from a Ficoll/Paque separation were washed three times and adjusted to 2×10^6 /ml.

Equal volumes (10 ml) of SRBC and lymphocytes were mixed, incubated at 37°C for 10 minutes, sedimented at 400g for 10 minutes and left to incubate at 4°C for 30 minutes. The pellet was resuspended and an aliquot taken for determination of % EAC- rosetting lymphocytes. The rosettes were determined as for E- rosettes. (Section 2.e)

The remainder of the resuspended cells were layered on to 4 X 4ml volumes of Ficoll/Triosil and centrifuged at 400g for 30 minutes at 18°C. The interfaces, containing lymphocytes depleted of cells with C3 receptors, were harvested and washed.

h. Surface Ig

This is a technique in which a fluorescein isothiocyanate (FITC) labelled anti-human Ig was used to enumerate cells expressing surface Ig.

Materials Sheep anti-human Ig (fluorescein conjugated)
diluted in PBS (1:5)

Method

Lymphocytes prepared by Ficoll/Paque separation were washed three times, counted and 10^6 cells pelleted. The supernatant was discarded and 0.2 ml of FITC anti-human Ig added to the pellet, mixed and incubated at 4°C for 30 minutes. The cells were washed three times with PBS and the final pellet resuspended in three drops of 40% glycerol in PBS. The cells staining for surface Ig were assessed by their fluorescence under a UV microscope. Positively staining cells had a bright yellow cell membrane.

i. Mitogen stimulation of lymphocytesMaterials

Phytohaemagglutinin (purified grade) (Wellcome Ltd.)

Micro-titre plates (sterilin) from Arnold R. Horwell, Ltd., Kilburn, London.

Oxford sample micropipetting system and polypropylene tips. Oxford Laboratories, International Corporation, Ireland.

Scintillant - PPO (2.5 diphenyl oxazole) : 12.5g

POPOP (14 Di(2-(4-methyl 5 phenyloxazolyl Benzene): 0.75g

Toluene : 2.5 l

Glass fibre filter papers (Whatman Ltd.) : 2.5cm.

GF/C

Method

Lymphocytes, obtained by aseptic Ficoll-Paque separation, were washed with sterile TC199 three times, resuspended in TC199 containing 10% heat-inactivated new-born calf serum at 10^6 /ml. The suspension was aliquoted in 200 μ l volumes into micro-titre plates and PHA added to give final concentrations within the range 0.2 - 100 μ l/ml. All this preparation was carried out in a Microflow sterile cabinet. The microtitre plates were covered and incubated in 5% CO₂/95% air at 37°C at high humidity for three days.

The contents of each well were resuspended by repeated withdrawals and expulsions from an Oxford pipette system and

$^{29}\text{Ci}/\text{mmol}$ ^3H -thymidine added at a strength of $1\mu\text{Ci}/\text{ml}$ cell suspension to each well. The cell cultures were returned to the 37°C , 5% CO_2 incubator for a further 60 minutes.

The contents of each well were again resuspended and transferred to a glass tube. The well was washed out with 200 μl distilled water and the washings added to the cell suspensions. Approximately 1ml distilled water was added to each tube which was mixed with the cell suspension by a whirlimixer. This lysed the cells to allow release of the cell contents. 2ml of ice-cold 10% trichloroacetic acid (TCA) was added to each tube which was whirlmixed again. Each tube was transferred to 4°C for one hour to allow the macromolecules to precipitate.

The tubes' contents were resuspended on a whirlimixer, and the precipitate filtered on to a glass fibre filter and washed with 20 ml of a mixture of three parts ether to one part ethanol and then with 20 ml of ether alone. The filters were dried at 37°C for two hours and placed in 2" X $\frac{1}{2}$ " glass vials covered with 3ml of the scintillant. Radio-activity incorporated into DNA was counted in a Phillips liquid scintillation analyser using an internal standard.

Whenever possible, all samples were tested in triplicate. The results were calculated as disintegrations per minute (d.p.m.) and used as a measure of DNA synthesis. Control values were for samples without PHA. Incorporation of test cultures was expressed as a percentage of the control.

CHAPTER THREE
FICOLL DENSITY GRADIENT SEPARATION
OF LYMPHOCYTES

A practical means of separating lymphocytes by size was needed so that functionally and morphologically different populations could be enriched, evaluated and characterised.

3.2 INTRODUCTION

The purification and separation of lymphocytes and sub-populations of lymphocytes from various tissues has been accomplished by filtration of cells through columns of glass wool, nylon fibres or glass beads (Cooper, 1968; Lewis and Robbins, 1972; Cudkowicz, Bennett and Shearer, 1964), by cannulation of lymphatic vessels (Koster and McGregor, 1970) and by electrophoresis (Thomas and Cater, 1972). The ability to fractionate a complete population of cells into sub-populations differing in one or more physical characteristic and still capable of normal function could be a very powerful tool for the study of many problems in cellular biology, particularly in the field of differentiation. Such a technique could be used preparatively, to obtain a purified sub-population of cells to be used in further experiments or analytically to study the properties of a particular sub-population of cells under varying conditions. The analogy with biochemistry is obvious, a field in which physically based techniques such as centrifugation, electrophoresis and chromatography are widely used, both preparatively and analytically.

Microscopic examination of most cell populations show that different cell types vary greatly in size, and that cell size is often correlated with cell function. Thus a technique producing fractionation of different sized cells might also produce

fractionation of functionally different cells. There are a number of techniques available for the separation of lymphocytes on the basis of size.

Glass-bead column filtration, originally described by Shortman (1966) is based on the simple (and, as it turns out, oversimplified) physical proposition that if beads of sufficiently small size are packed into a column, so that the pores between them become comparable in size to the diameters of cells, the flow of large cells will be hindered more than that of small cells. The method is simple, rapid and can yield very pure small lymphocytes. Dead and damaged cells are trapped, so that a high proportion of the passed cells are viable. Using this technique, the immunological potential of small lymphocytes has been investigated by Shortman and Szenberg (1969); Shortman et al (1971); Lewis, Mitchell and Nossal (1969); and Hunt, Ellis and Gowans (1972). However, the yields are very low if large lymphocytes are to be eliminated effectively, often a second passage through the column being necessary. Physical adherence to the glass beads also tends to remove plaque-forming (Plotz and Talal, 1967) or erythrocyte-binding (Wigzell and Makela, 1970) cells from spleen cell suspensions, indicating that thymus-independent lymphocytes are more adherent than thymus-dependent lymphocytes. The electrical charge of the lymphocyte is another factor which the columns may distinguish (Sela et al, 1970). This method was not used because of the total loss of large cells from the lymphocyte preparations.

Large lymphocytes are preferentially destroyed during incubation in unfavourable conditions, thus producing a relative enrichment of small lymphocytes. This is performed at a high cell concentration ($> 10^7$ /ml) shaking at 37°C overnight in serum-depleted medium,

(Gowans, 1962). It is simple and effective (contaminating large lymphocytes being reduced to less than 1 per cent) and leaves small lymphocytes still capable of immunological function (Lewis, Mitchell and Nossal, 1969; Ellis, Gowans and Howard, 1969; Gowans and Uhr, 1966). Its main drawbacks are the variable overall loss of small lymphocytes; there may thus be an undetected differential survival of sub-populations of small lymphocytes. Again, the large lymphocytes are not recoverable.

The main problems of the above techniques are the loss of cells in toto and the selective destruction of large lymphocytes. More desirable techniques allow the separation of cells by volume and density with recovery of all cell fractions at high yields.

Fulwyler (1965; 1970) has described an electronic cell sorter which separates cells on the basis of volume. Although significant separations can be achieved with this apparatus (Van Dilla et al, 1967) its complexity and the low rate at which it separates cells make it unsuitable for routine fractionations of heterogeneous populations of cells.

In theory it might be possible to use differences in sedimentation rate of separate cells on the basis of size. The sedimentation velocity of a spherical cell falling through a fluid under the influence of gravity is given by Stokes Law.

$$S = \frac{2}{9} \frac{(\rho - \rho')}{\eta} g r^2$$

Where η is the coefficient of viscosity, ρ and ρ' the density of the cell and fluid medium respectively, g the acceleration due to gravity and r the radius of the cell. If it can be arranged that

($\rho - \rho'$) and n remain practically constant, it is evident that the final sedimentation velocity becomes a sensitive function of the size of the cell being proportioned to the square of its radius. Variations in the density of the cells and of the medium become relatively unimportant if ($\rho - \rho'$) is substantially greater than variations in either ρ or ρ' . Thus, in the conditions of the normal experimental procedure, although density may exert a small influence, the major factor determining sedimentation velocity is size.

..... Théorétically, the layering of a band of cells over a medium of constant density should allow a separation of large from small lymphocytes, the larger cells sinking more rapidly. In practice, however, two complications interfere - the first concerns the stability of the suspending medium during the introduction and recovery of the cells; any turbulence due to mixing or convection would ruin the separation. To avoid this, a very shallow density gradient can be imposed on the medium, solely to stabilize it. The gradient should be as shallow as possible in order to retain the approximate constancy of ($\rho - \rho'$) and n . The function of this gradient is totally different from those employed in isopycnic centrifugation. The second complication is that Stokes Law is not obeyed by cells at high concentrations where, instead, characteristic 'streamers' of cells break away from the main band and sediment at an excessively fast rate. Streaming, which can be detected by careful visual inspection of the cell band, and by analysis of the final profile of the sedimented cells, can be overcome by lowering the cell concentrations to less than the streaming limit. To separate large numbers of cells the size of

the sedimenting chamber has to be increased. However, pools of enriched large lymphocytes have been obtained (Hunt, Ellis and Gowans, 1972) and these pools were shown to be enriched particularly in lymphocytes in their DNA synthetic ('S') phase, since it has been shown that, in common with many other cell types (MacDonald and Miller, 1970), 'S'-phase large lymphocytes sediment more rapidly than those not in 'S'-phase (Hunt, Ellis and Gowans, 1972; Yoshida and Osmond, 1971).

This technique has been used in analyses of mouse spleen (Phillips and Miller, 1970; Osoba, 1970; Miller and Phillips, 1970) and bone marrow (Edwards, Miller and Phillips, 1970) of human blood (Brubaker and Evans, 1968) rat bone marrow (McCool et al, 1970) and in a partial separation of thymus-derived and thymus-independent peripheral lymphocytes of the rat (Howard, Hunt and Gowans, 1972).

Functional studies using velocity sedimentation have included the separation of cytotoxic T lymphoblasts from mouse spleen (Greenberg 1973), transformed lymphocytes from mouse malignant neoplasm (Zettergren, Luberoff and Pretlow, 1973), cytotoxic effects on lymphocytes generated by human mixed leucocyte reaction (MLR) (Fradelizi et al, 1977), mouse helper and suppressor T lymphocytes (Tse and Dutton, 1976) and killer cells infiltrating allografts in mice (Roberts and Hayry, 1977).

There are a number of problems associated with velocity sedimentation; firstly, the time taken for sedimentation is usually in excess of four hours and the preparation of the gradient takes at least another hour, reducing the number of samples separable in one day. Second, it is virtually impossible to compare the size range recorded from one sample with another sample if fine differences are to be noted. Third, if the harvested fractions are to be cultured all

of the sedimenting chamber and gradient maker have to be kept aseptic, not a simple operation if the equipment has to be left standing for five or more hours on a laboratory bench. Fourth, the sedimenting chamber has to be large in diameter, and hence the gradient media is more susceptible to convection currents and physical vibration.

The third, more commonly used, technique for lymphocyte separation is density flotation (or isopycnic centrifugation). This has been developed from the original work by Leif and Vinograd (1964) who used centrifugation to equilibrium in an albumin density gradient to effect separation between young and old erythrocytes and to characterise an individual erythrocyte population with a high degree of precision and reproducibility. Other earlier workers (Hilal, et al, 1964) employed dextran density gradients to separate cells of the myeloid series from bone marrow.

The separation of cells by centrifugation to equilibrium in density gradients should reflect differences only in cell density, not cell size, since the cells come to density equilibrium in a gradient which covers the range of densities in the cell population. So, compared with velocity sedimentation the density difference between the density flotation media and the cell population is small and a much higher g force is used for density separation, often due to the high viscosity inherent in solutions with a density above 1.05 centipois (cps).


Although density gradient centrifugation does not separate cells by size there is reason to predict an indirect relationship between size and density. The nucleus of a lymphocyte should be more dense

than the cytoplasm. Thus, a small lymphocyte with proportionally more nucleus, should be more dense than a large lymphocyte with proportionally more cytoplasm. This was confirmed by Shortman (1971), using thymocytes, who compared the volumes of cells from gradient fractions of different densities. Using both Coulter Counter size distribution data and direct morphological examination of stained preparations it was possible to determine the density distribution of cells of a given size range (Fig. 3.0). This established that even a given size or morphological class of lymphocyte was heterogeneous. Shortman also noticed that not all large cells were light and not all small cells were dense, although this general rule was adequate to obtain enriched populations of lymphocytes of various sizes, if not pure ones. Such a distribution may reflect the presence of several independent lymphocyte populations, each with its own series of large, medium and small cells. Alternatively it may reflect a series of definite metabolic stages through which any one lymphocyte may pass in its development.

Recently evidence has shown that density flotation may be more sensitive than velocity sedimentation in separating newly activated lymphocytes. (Steinman et al, 1978). Velocity sedimentation separates large cells, mainly those in the S and G + M phases of the cell cycle, rather than blast transformed cells per se. This is also a problem showed by ^3H -thymidine labelling of lymphocytes; the recently transformed cells, in G_1 of the cell cycle are not identified or characterised (Splinter and Reiss, 1974). On the other hand, density flotation has been successfully used to identify and characterise in vitro stimulated lymphoblasts in the G_1 phase (Steinman et al, 1978). Low density blasts, identified morphologically by basophilic cytoplasm, readily discernable nucleolus and abundant

Fig.3.0 "The average volume of normal rat thymus lymphocytes banding in various regions of a density gradient. The results are from a single animal, but are typical of many experiments."

From: Shortman, K.(1971.) J. Cell. Physiol. 77: 319



polyribosomes, were found within 20-24 hours of stimulation of mouse spleen cells by mitogens. These cells were not markedly enlarged, but went on to enlarge and synthesize DNA after a further 18-20 hours in culture.

It is not known why or when, during its initial cell cycle, the transformed lymphocyte acquires a low density. Possibly changes in the condensation of chromatin or in the water content of the cell (? nucleus) are involved. It appears that the density shifts occur before DNA synthesis begins. Gunthe et al (1974) obtained information that stimulated cells in G_1 are irreversibly committed to entering S phase many hours before they actually synthesize DNA. They noted that removing the stimulus (Con A) after 20 hours of stimulation did not prevent the cells from entering S phase after a day of additional culture.

Shortman et al (1967) showed that Vinblastine, a mitotic inhibitor, when administered to antigen stimulated mouse lymphocytes in vitro, caused an increase in the number of less dense cells in a density gradient and a marked depletion of others. As it is a drug that causes a build-up of cells in the process of division and a relative depletion of product cells, this lends further support for the usage of density gradient techniques to separate and characterize activated and non-activated cells. The ability to separate a number of samples concurrently is another advantage over velocity sedimentation, as well as the separation taking less time and media. Linked with the separation of newly activated cells, not possible with velocity sedimentation, the merits of density flotation are clear.


Density gradient centrifugation has been used to identify

antigen sensitive cells (Haskill, 1967; 1969) and cells forming 19S haemolytic antibody in the rat (Haskill, Legge and Shortman, 1969) to enrich chicken blood lymphocytes initiating a graft-versus-host reaction (Shortman and Szenberg, 1969), to demonstrate several classes of killer cells found in allograft rejection (Roberts and Hayry, 1977), to characterize cells involved in the immune response to sheep erythrocytes (Haskill, Byrt and Marbrook, 1970) and to recover mouse spleen lymphoblasts generated in vitro (Steinman et al, 1978).


There are two ways of using gradient centrifugation techniques; continuous gradients and discontinuous gradients. The continuous gradient is prepared in such a manner so that there is an uninterrupted range of concentrations within the centrifuge tube as illustrated in Fig. 3.1a. This requires a gradient generator to mix a light solution with a heavy solution and pump it into the centrifuge tube. The discontinuous gradient is prepared by carefully layering sequentially lighter solutions upon each other in a centrifuge tube (Fig. 3.2a). Upon centrifugation of a continuous gradient the cells find their density equilibrium at the relevant point within the gradient. To harvest the separated cells a needle, with a tube attached, is inserted into the bottom of the centrifuge tube and the contents drained off in as many fractions as required. In a discontinuous gradient the separated cells collect at the interfaces between the layers of media. This gives a limited number of fractions, but the harvesting is easily accomplished by aspirating off the successive layers.

Loos and Roos (1974) compared the two methods and found that the discontinuous system required less preparation, was easier to

Fig. 3.1.

The generation of density gradients

a) Continuous gradient



b) Discontinuous gradient

harvest, and gave a better yield of viable lymphocytes (approximately 50% more). They recommended its use for preparative techniques and I employed it routinely for lymphocyte separation.

The separation of lymphocyte sub-populations on the basis of density differences places particular demands on the materials to be used for the preparation of the gradients. Obviously, the material must be dense enough to produce gradients covering the normal lymphocyte density range (1.055 - 1.095 g/ml). Besides, the material must allow generation of gradients which are iso-osmotic and with a constant pH through the whole gradient. Moreover, the material should not influence the cells with respect to aggregation, density or viability, nor should it be antigenic. Additionally, its viscosity should be low in order to allow short centrifugation times and easy measurement of its physical properties. Finally, the gradients should give high cell yields and reproducible patterns.

Several materials have been used for the generation of density gradients, continuous as well as discontinuous, among them sucrose, gum acacia (Kinoshita et al, 1970), Renograffin (Grdina et al, 1973), albumin (Shortman, 1968; Williams, Kraft and Shortman, 1972; Haskill, 1969), Ficoll-Hypaque (Day, 1972), Ficoll-isopaque (Loos and Roos, 1974) and Ficoll (Bach and Brashler, 1970; Gorczynski, Miller and Phillips, 1970). However, none of these materials is completely without disadvantages. Sucrose permeates easily through lymphocyte membranes and exerts a relatively higher osmotic pressure at high concentrations. Gum acacia shows a non-linear relationship between concentration and osmolarity and its viscosity is too high for easy handling (Loos and Roos, 1974). Renograffin is not isotonic throughout the gradient (Grdina et al, 1973). Bovine Serum Albumin (BSA) and Ficoll are the two materials most widely used for lymphocyte separations., Ficoll having superseded BSA to some extent in recent

years.

BSA has been strongly advocated for lymphocyte separation by Shortman (Shortman, 1966; 1968; 1971; Shortman and Szenberg, 1969; Shortman et al, 1971) and Haskill (Haskill, 1967; 1969; Haskill, Byrt and Marbrook, 1970) whereas Ficoll has been favoured by Bach and Brashler (1970), Gorczynski, Miller and Phillips (1970), Glinski et al, (1976) and Yu et al (1973).

Shortman (1968) noted a marked heterogeneity in the density distribution of lymphocytes from a number of tissues in continuous BSA gradients. Many discrete bands or peaks were noted, suggesting a far greater heterogeneity than expected from biological considerations such as cytological criteria. Haskill, Legge and Shortman (1969) also noted a heterogeneity of antibody producing cells from mouse spleen. These authors thought that such distribution patterns may reflect the presence of several independent lymphocyte populations, each with its own series of large, medium and small cells. Alternatively, they believed it may reflect a series of definite metabolic stages through which any one lymphocyte may pass in its development.

Gorczynski, Miller and Phillips (1970) pointed out that velocity sedimentation of antibody-forming cells indicated that by size, those cells are a homogeneous population (Phillips and Miller, 1970). Although there was some heterogeneity in sedimentation velocity this could almost entirely be explained in terms of cell volume changes associated with cell growth through the division cycle. If there was heterogeneity of the magnitude reported by Haskill et al (1969), the distribution of sedimentation velocity for antibody-producing cells should have been much broader than was observed

and should have shown some fine structure.

Worton, McCulloch and Till (1969) were disturbed about the lack of reproducibility of the reports on BSA density gradient analyses of mammalian cells. This could be caused by the nature of the gradient material. BSA is a natural product and its isolation may vary from one batch to another, variations in different preparations can generate different density profiles for the same population of cells (Shortman, 1968). In addition, because BSA is a charged molecule, it will bind ions and decrease the osmolality of the solution (Scatchard, Batchelder, and Brown, 1946). Changes in osmolality are known to radically alter the density distribution of some classes of cells (Shortman 1968; Gorczynski et al, 1970). Also, to use BSA successfully the pH has to be quite low, e.g., Shortman (1968) used pH 5.1. However, the heterogeneity he reported for lymphocyte density distribution is almost certainly due to the osmotic gradient formed within the density gradient. The inability of a peak of harvested cells to reband at the same point in a second BSA gradient indicates that the BSA affects the density of the cells in the gradient.

In contrast, Ficoll, a large polymer of sucrose, is not a natural product and can be synthesized in large amounts to a required standard. It contains no ionized groups and hence does not need the rigorous desalting that must be applied to BSA before its use in a density gradient. Ficoll has a very low permeability towards cell membranes and hence cannot change cell density in the same way that, for instance, unpolymerized sucrose may do.

Cells will associate or aggregate in Ficoll unless the divalent cations calcium and magnesium are omitted from the media in which the

Ficoll is dissolved (Walder and Lunseth, 1963; Yu et al, 1973).

Alternatively, a dispersing agent, 2-naphthol-6,8-disulphonic acid (dipotassium salt), has been used successfully (Shortman, 1968).

The pH may be maintained at a physiological level in a Ficoll gradient whereas with BSA aggregation can only be reduced by lowering the pH to 5.1, which increases lymphocyte density (Legge and Shortman, 1968).

Gorczynski et al (1970) found that continuous Ficoll gradients are isotonic throughout the density range used for lymphocyte separation and that Ficoll is neither toxic nor stimulatory for small lymphocytes, a report confirmed by Loos and Roos (1974) and Yu et al (1973). The basic test to determine the validity of a density distribution profile of rebanding a single peak of lymphocytes in a second gradient has been used to evaluate Ficoll gradients, both continuous and discontinuous (Loos and Roos, 1974; Bach and Brashler, 1970; Yu et al, 1973; Glinski et al, 1976; Williams, Kraft and Shortman, 1972). This was successful in all cases, but depends upon the media being isotonic to the plasma osmolarities, varying between 260 and 320 milli-osmoles (Williams, Kraft and Shortman, 1972).

Ficoll appears to be fully established and acceptable as a density gradient material and in some ways preferable to BSA, not least being its relative cheapness. Ficoll was used throughout for the analysis and preparation of lymphocyte populations in discontinuous density gradients.

3.3 DISCONTINUOUS FICOLL DENSITY GRADIENT METHODOLOGY

The method described was that routinely employed for lymphocyte density separation. During the design of this method a number of variables were examined and these will be described in the next section.

a. Lymphocyte preparation

Lymphocytes were prepared from peripheral blood by isopycnic centrifugation on Ficoll/Triosil as described in chapter 2. 2b.

Calcium-and-magnesium-free media (CMFSS) was used throughout for blood dilutions and lymphocyte washings.

Calcium-,and-magnesium-free salt solution (CMFSS)

Sodium chloride	8000 mg l^{-1}
Glucose	1600 mg l^{-1}
Potassium chloride	400 mg l^{-1}
Potassium di-hydrogen phosphate	60 mg l^{-1}
Di-sodium hydrogen phosphate	48 mg l^{-1}

pH 7.3 at 18°C

Occasionally phenol red (17mg l^{-1}) was added to enable a constant visual check on the pH during storage.

The media was usually made up as a 10 X concentrate, filter sterilized and stored in 125ml bottles at 4°C. Each bottle was diluted with distilled water and the pH adjusted with concentrated KH_2PO_4 or Na_2HPO_4 .

b. Discontinuous Ficoll gradient generation


Ficoll 400 was dissolved with stirring overnight at 4°C in CMFSS. A stock solution of 30% Ficoll was prepared in CMFSS and filter sterilized into sterile 30ml McCartney bottles for storage at -20°C.

Prior to use the Ficoll was thoroughly thawed, mixed and left to stand to allow air bubbles to escape. Dilutions were made in test tubes as follows:-

<u>%Ficoll</u>	<u>30% Ficoll (ml)</u>	<u>CMFSS (ml)</u>
16	2.13	1.86
17	2.26	1.73
18	2.40	1.60
19	2.53	1.46
20	2.66	1.33
21	2.80	1.20
22	2.93	1.06
23	3.06	0.93
30	4.00	0

The tubes were thoroughly mixed on a whirlimixer and left to stand to allow air bubbles to escape.

Eighteen ml. glass centrifuge tubes (110 X 15mm) were used for the density gradient centrifugation. Consequently lighter 2ml volumes of Ficoll solution were carefully layered above each other as shown in Fig 3.2, excepting the 16% Ficoll. Two density gradients were always prepared for strict control in sample



comparisons of density distribution.

c. Monocyte depletion

The cell pellet, consisting of washed lymphocytes and monocytes, was resuspended in 2ml of 16% Ficoll in a glass tube and incubated at 37°C for 30-60 minutes. This lymphocyte rich supernatant was decanted and carefully layered at the top of the density gradient.

d. Density gradient centrifugation

..... The separation was in the swing-out head of a refrigerated centrifuge (MSE Mistral 2L) at 10°C. A fixed temperature was used to prevent convection currents and the changes in viscosity and density which occur with temperature variations (Shortman, 1968). The centrifuge speed was increased slowly to prevent mixing of the gradient and a force of 3,000g, at the tip of the tube, maintained for one hour. The brake was not used for running down the centrifuge motor, again to prevent mixing of the gradient.

The lymphocytes localised at the interfaces between the gradient layers. To harvest the cells the layers were sequentially withdrawn using a pasteur pipette, rotating the pipette around the inside wall of the centrifuge tube while aspirating. In this manner the contamination by adjacent interfaces was minimal.

Each interface suspension was ejected into a graduated tapered glass centrifuge tube, made up to approximately 10ml with CMFSS, thoroughly mixed by repeated withdrawals and expulsions in a pasteur pipette and the cells sedimented by centrifugation at 200g for 10 minutes. The cell-free supernatant was decanted and discarded and the cell pellet made up to 0.5ml with CMFSS and thoroughly mixed. The

lymphocyte count was determined by diluting an aliquot in white cell counting fluid (WCCF) and counting the cells in a haemocytometer using the low power (X10) of a binocular microscope.

3.4 EXPERIMENTAL DESIGN - INVESTIGATION OF PARAMETERS

A range of techniques for density gradient centrifugation using Ficoll have been described (Zettergren, Luberoff and Pretlow, 1973; Loos and Roos 1974; Yu et al, 1976; Bach and Brashler, 1970; Glinski et al, 1976; Glinski, Gershwin and Steinberg, 1976; Minderhoud and Smith, 1972; Williams, Kraft and Shortman, 1972; Gorczynski, Miller and Phillips, 1970).

a. Density gradient material

Ficoll is a synthetic high polymer (MW 400,000) made by the co-polymerization of sucrose and epichlorohydrin. The molecules have a branched structure and have a high content of hydroxyl groups leading to a very good solubility in aqueous media. No ionized groups have been found in Ficoll (Pharmacia Information Services).

Ficoll has been used in a mixture with Isopaque (sodium-, calcium-, and magnesium-N-methyl-3,5-diacetamido-2,4,6-tri-iodobenzoate) to form density gradients for lymphocyte separation (Loos and Roos, 1974). This was used successfully with both continuous and discontinuous gradients, showing similar density distributions and with good cell yields, particularly in the discontinuous gradients.

My attempts to repeat these experiments were fruitless. The Ficoll/Triosil mixture detailed in Chapter 2.2b for lymphocyte isolation was used diluted with phosphate buffered saline to produce a density range of 1.040 - 1.080 g/ml in six steps as a discontinuous gradient. Upon applying the lymphocyte sample to the surface of this gradient and centrifuging at 3,000g the cells aggregated to a large extent and formed a monolayer at one of the gradient interfaces. Possibly Triosil is unsuitable by virtue of it having a slightly different formulation to Isopaque, even though both

are mainly sodium-, calcium- and magnesium salts of metrizoic acid.

Ficoll/Triosil was rejected as a density gradient medium and Ficoll alone in solution was the next medium investigated. Initial attempts used a full PBS media but the divalent cations, calcium and magnesium, caused aggregation of lymphocytes in the gradient and the cells formed a monolayer at the first gradient interface, i.e. did not distribute according to density. The calcium-, and magnesium-free salt solution was based on the Flow Catalogue


Formulation for Hanksbalanced salt solution and prevented lymphocyte aggregation to a great extent. Similar results were found by Bach and Brashler (1970) and Yu et al (1973) who remark on the importance of eliminating cell aggregation. Aggregation of lymphocytes of varying densities may seriously distort the density distribution pattern. An association between a large light cell and a small dense cell would result in a doublet with a buoyant density less than the arithmetic mean of the two densities. The mean cell volume of cells in the less dense region of the gradient is usually greater than cells in the more dense region. Thus, an increasing proportion of doublets and larger aggregates with higher cell concentrations would be expected to cause a small decrease in the apparent average buoyant density. More importantly, preparative techniques would be rendered impractical due to the contamination of the density gradient fractions and further analysis of gradient fractions would be of only limited value.

The density range of various Ficoll concentrations is shown in Fig. 3.3 The range of Ficoll concentrations used here produces a similar range of specific gravities to those used by other authors to separate lymphocytes into different density classes (Shortman, 1968;

Fig. 3.3.

Density of Ficoll as a function of
concentration.

86



Haskill, Legge and Shortman, 1969; Haskill, Byrt and Marbrook, 1970; Loos and Roos, 1974) although upon close examination the mean density of lymphocytes varies by as much as 0.020g/ml from one group to another.

My initial experiments with the density gradient had a wider range of Ficoll concentrations than finally employed. After several lymphocyte separations it became clear that the lymphocytes were only separating in the 16-23% Ficoll layers. This was determined by visual examination of the intact gradients to detect the presence or absence of cells. The final gradient means that many lymphocytes were found in fraction G and so produced a wider, better density distribution analysis of the less dense cells than of the dense cells.

Fortunately, Ficoll solution is viscous enough to allow solutions varying only by 0.004g/ml to be layered above each other by hand with a pasteur pipette and bulb. The process is slow however, and requires a certain amount of practice to ensure a clear interface Schlieren between layers.

The Schlieren line between the layers disappears after a few minutes but the interfaces and layers stay intact much longer. This was demonstrated by colouring, with a small amount of 1% Trypan blue, the 17%, 19%, 21% and 23% Ficoll layers before forming a gradient. After the gradient had been made it was left to stand at 4°C for 36 hours during which time the coloured layers had not mixed at all with the adjacent clear layers, confirming the stability of the gradient.

Centrifugation up to 3,000g maximum with gentle acceleration and deceleration, did not disturb the layers.

In many published studies on density gradients the cells were

suspended in the most dense solution of the gradient because suspending cells in the light layer may cause distortion of the density profile by the "wall effect" (Anderson, 1955). This was examined by Leif and Vinograd (1964) who found that a certain proportion of cells were lost by sedimentation against the cylindrical walls of the tube. More recently, Yu et al (1973) examined this phenomenon in discontinuous gradients of Ficoll and found the density distribution profiles very similar whether the cells were resuspended in 5% or 30% Ficoll. Because of the differences in viscosity, it is easier to disperse cells in a low density Ficoll solution than a high density Ficoll solution. This prevents aggregation which has been shown to seriously distort the density profile. Similar results to those of Yu et al (1973) with Ficoll gradients have been shown with discontinuous BSA density gradients (Adler, Peavy and Smith, 1970). For these reasons I always suspended the cells in the lowest density Ficoll in my experiments.

b. Duration and force of centrifugation

The duration of centrifugation required for equilibrium was tested. The duration finally employed (60 minutes) was more than sufficient for equilibrium and is similar to that used by Raidt, Mischell and Dutton (1968) in their discontinuous albumin gradients.

For confirmation of this lymphocytes were prepared from 50ml venous blood from a healthy person, (S.C.), divided into two equal amounts and separated in two Ficoll gradients. One was centrifuged at 3,000g for 40 minutes and the other at 3,000g for 60 minutes. Cells were seen only at the interfaces of both gradients, suggesting

equilibrium had been attained. The density distribution profiles were virtually identical (Fig. 3.4), suggesting that 60 minutes at 3,000g was more than adequate to permit equilibration.

There are several factors to consider when determining the optimum centrifugation speed. The force applied needs to be high enough to bring the system to equilibrium in a reasonable time, and in addition, high enough to overcome the forces causing light and dense cells to associate together and thus to band in a region of intermediate density. However, to reduce the possibility of high forces damaging cells, the speed should be as low as practicable. The force used (3,000g) is at the low end of the range of force used by a number of workers which extends up to 55,000g yet, as it appears to allow equilibrium, there is no need to go any higher.

c. Cell load

Cell to cell association affects density distribution, and possibly the cell load affects cell interaction. Shortman (1968) and Yu et al (1973) have investigated this parameter to see if cell load changes caused variations in lymphocyte density distribution profile. They have suggested upper limits of at least 5×10^8 cells and 3×10^8 cells per tube respectively before the profile changes. The maximum number of cells loaded in any experiments described here is 1×10^8 /tube, below either reported maximum.


d. Rebanding of a gradient fraction

A basic test to determine the validity of a density distribution profile is to reband a single peak or fraction of cells in a second gradient. This fraction should then reform only one fraction, at the same density position. In tests with rat thymus cells

Fig. 3.4

The effect of centrifugation time on the
density separation of healthy lymphocytes.

60 ml of peripheral blood was taken from a healthy subject and the lymphocytes prepared (Chapter 2.2.b ii). The monocytes were depleted by adherence (Chapter 3.3.c) and the lymphoid cells separated by density in a Ficoll gradient (Chapter 3.3.d.) at 3,000g for either 40 minutes or 60 minutes. The interfaces were harvested and counted and the results plotted as the percentage of the total recovered lymphocytes found in each fraction.

Fig. 3.4.

(Shortman, 1968) an isolated fraction formed only one peak on a second gradient, but with a shift to the more dense regions.

This was shown to be due to stress caused by the high force used in the second centrifugation. The relatively low centrifugal force used in these experiments should not cause any change in optimal rebanding of a fraction of cells.


Rebanding experiments have been used to validate density gradients of BSA (Shortman, 1968; Adler, Peavy and Smith, 1970; Williams, Kraft and Shortman, 1972), Ficoll-Isopaque (Loos and Roos, 1974) and Ficoll (Bach and Brashler, 1970; Gorczynski, Miller and Phillips, 1970; Yu et al, 1973; Gliniski et al, 1976). Two purposes are fulfilled by **successful rebanding**. First, the reproducibility of the gradients and the degree of resolution may be shown. Second, cell to cell association in the first gradient would show itself in the second gradient by the appearance of cells in more fractions than that being rebanded. If no major spreading of the rebanded fraction occurred it would show that large and medium lymphocytes were banding at their own buoyant density and were not just non-specifically trapped by their association with small lymphocytes.

Lymphocytes from a healthy donor were separated on a discontinuous Ficoll gradient, the fractions harvested and washed with CMFSS. The cells from fraction G were pelleted, resuspended in 16% Ficoll and separated on a second Ficoll gradient. The fractions were **harvested**, washed and cell counts performed. The results are shown in Fig. 3.5. The majority of the cells (84%) from fraction G from the first gradient reisolated

Fig. 3.5

Re-centrifugation of lymphocytes from a Ficoll density
gradient fraction in a second gradient.

Lymphocytes were harvested from 30 ml venous blood (Chapter 2.2.b.ii) and the monocytes depleted by adherence (Chapter 3.3.c.). The lymphocytes were separated by density in a Ficoll gradient at 3,000g for one hour. The fractions were harvested and counted and fraction G re-centrifuged in a second Ficoll gradient. The fractions were again harvested for lymphocyte counts. The results are plotted as percentage of total recovered cells in each fraction.

Fig. 3.5.

at G in the second gradient. This degree of enrichment in the reisolation is enough to suggest that the cells are separating by their buoyant density. This is a better re-isolation than some authors have obtained, particularly so with the narrow density range employed here. The second gradient separation was performed as soon after the first as possible, because other authors have reported that prolonged exposure to gradient material may affect lymphocyte density (Shortman, 1968; Williams, Kraft and Shortman, 1972; Bach and Brashler, 1970; Loos and Roos, 1974), particularly by increasing the density of the reisolated lymphocytes. This was not found to be a problem in this experiment.

e. Reproducibility

To test the reproducibility of replicas a sample of healthy peripheral blood lymphocytes was divided into two equal aliquots of 2×10^7 cells and put through two separate density gradient tubes. Fig. 3.6 shows that the differences between the two tubes are small and that comparisons of density profiles of cells in identical gradients, prepared and centrifuged together, are valid.


If serial studies of density distribution profiles from the same subjects are to be attempted, the day to day, week to week and month to month variation should be considered. The various cells involved in the immune response are constantly being turned over, lymphocytes with varying functions have different life spans (Gowans, 1966). It is important to know if the density distribution profile of various healthy subjects lymphocytes is representative of a status quo in the lymphocyte dynamics or a phase in an ever fluctuating pattern.

During the course of experimentation the lymphocytes from some

Fig. 3.6

Reproducibility of Ficoll density
gradients

60ml of blood was taken from a healthy subject and the lymphocytes separated (Chapter 2.2.b ii). The monocytes were depleted by adherence (Chapter 3.3.c.) and the sample separated in two Ficoll gradients (Chapter 3.3.d.). The interfaces were harvested the lymphocytes counted and the results compared.

Fig. 3.6.

healthy individuals, notably laboratory personnel, have been used several times for comparison with patients suffering from rheumatic diseases. One example is shown in Table 3.1. The density distribution profiles are for the lymphocytes from one person determined at approximately three monthly intervals and show little variation within each fraction. All blood samples were taken at the same time of the day to eliminate any diurnal variation. The stable pattern shows that in a healthy person there is an equilibrium in the range of lymphocytes as detected in this system, in the peripheral blood. This pattern is expected to change during immunization or infection as shown by cytological examination (Crowther, Hamilton-Fairley and Sewell, 1969).

f. Yield and viability

The yield of lymphocytes from the gradient was found to vary between 30% and 60%. This is similar to the results of other authors. Higher yields have been reported for BSA (Shortman, 1968; Haskill, Legge and Shortman, 1969) and the addition of human albumin (1%) to Ficoll solutions has also been reported to increase the yield from a Ficoll gradient (Loos and Roos, 1974).

The cell viability in each fraction was assessed by the Trypan Blue dye exclusion test. Incubating the lymphocytes in the 16% Ficoll layer to remove monocytes by adherence had a beneficial side effect. The Ficoll conferred a degree of protection to the lymphocytes during the gradient centrifugation, particularly to the less dense cells which are more susceptible to damage. (Shortman, 1968). This phenomenon of cell protection by concentrated solutions of Ficoll has been reported previously

TABLE 3.1 DENSITY DISTRIBUTION PROFILES OF A HEALTHY .
SUBJECT'S (S.C.) PERIPHERAL BLOOD LYMPHOCYTES
(AT APPROXIMATELY THREE MONTHLY INTERVALS)

<u>FRACTION</u>	<u>CONSECUTIVE SAMPLES</u>					<u>$\bar{X} \pm 1 \text{ S.D.}$</u>
	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	
A	0	0	0	0.5	0.2	0.1 ± 0.19
B	0	0	2.9	1.5	0.3	0.9 ± 1.12
C	5.2	2.2	2.1	0.8	1.2	2.3 ± 1.54
D	3.6	3.2	2.7	4.5	3.5	3.5 ± 0.59
E	4.4	3.2	6.2	4.1	6.0	4.8 ± 1.15
F	12.3	9.2	10.9	9.0	7.8	9.8 ± 1.58
G	74.5	79.0	72.2	77.0	75.0	75.5 ± 2.31
H	0	0.9	3.0	2.7	6.0	2.5 ± 2.07

(All figures % of total recovered cells
in each fraction)

(Stuart and Simpson, 1970) and was found, in this work, to radically improve both yield and viability of density gradient fractions. The viabilities of lymphocytes prior to and after including pre-incubation in Ficoll as a medium for monocyte depletion is shown in Table 3.2. The Ficoll pre-incubation increases the cell viability, particularly in the less dense fractions. The poor viability in fraction H is due to the fact that most of the cells in this fraction are there because they are dead. When a lymphocyte lyses one of the first events is the appearance of holes in the cytoplasmic membrane. Then the cytoplasm is lost through these enlarging gashes leaving only the nucleus, which is denser than cytoplasm, and hence the cell becomes dense and sinks within the density gradient to equilibrate in a lower position than most of the viable cell population. The poor viabilities in the less dense cell fractions have been previously reported by Shortman, (1968) and he noted that most of the damaged cells were large lymphocytes. The fragility of large lymphocytes has been reported by other authors (Hall, 1966; Sewell, R. L., personal communication) although no reasons for this have been put forward.

g. Cytological examination of density gradient fractions

The cell suspensions of each fraction were washed in CMFSS, centrifuged to a pellet and smears made on duplicate glass slides. These were air dried and stained with Wrights stain (chapter 2.2d.). Under a microscope the less dense fractions contained more cells with the cytological appearance of activated lymphocytes and plasma cells.

Table 3.2

VIABILITY OF LYMPHOCYTES IN FICOLLDENSITY GRADIENT FRACTIONS

<u>Fraction</u>	<u>Pre-incubation in</u> <u>CMFSS</u>	<u>Pre-incubation in</u> <u>16% Ficoll</u>
A	12	64
B	10	61
C	34	60
D	35	75
E	81	91
F	88	92
G	89	90
H	15	30

All figures % viable cells as assessed by
dye exclusion

The dense fractions were relatively depleted of activated lymphocytes and plasma cells compared with the original population. Quantitative analysis was difficult because many of the cells in the low density fractions were damaged, proportionally more than shown in the viability data, indicating that at least some cells were damaged during the making of a smear, providing more evidence that large lymphocytes are relatively more fragile than small lymphocytes. In addition, all the damaged cells in the low density fractions had typical nuclear features of activated lymphocytes. The degree of enrichment of these large cells was thus difficult to determine accurately by normal microscopical means although inspection suggested that large lymphocytes were less dense than small lymphocytes.

To reduce cell damage some slides were made from cell suspensions using a cytocentrifuge. The cell morphology was distorted but more of the large lymphocytes were undamaged. Accurate measurement of cell diameter was impractical, but a visual assessment of lymphocyte activation on Wrights stained preparations was made on the basis of cell size, nuclear characteristics, nuclear: cytoplasm ratio, number of nucleoli and basophilia of cytoplasm. The results (Table 3.3) are from a patient with rheumatoid arthritis and a chronic chest infection who had large numbers of less dense cells. Increased numbers of large cells are shown in the less dense fractions and decreased numbers in the denser fractions. Contaminating granulocytes were usually found in fraction H, but a few (<5%) were occasionally found in fraction G. Granulocytes density (> 1.085 g/ml) is more than that of lymphocytes (1.050 - 1.080 g/ml) and so the result is expected. This is the means of

Table 3.3

DIFFERENTIAL COUNT OF LYMPHOCYTES IN FICOLLDENSITY GRADIENTLymphocyte Size

<u>Fraction</u>	<u>Small</u>	<u>Medium</u>	<u>Large</u>
A	51	32	17
B	42	35	23
C	39	36	25
D	40	35	25
E	46	39	15
F	78	22	0
G	96	4	0
H	91	9	0

All figures % of total in each fraction

separating lymphocytes and granulocytes on Ficoll/Triosil or Ficoll-Paque (isopycnic centrifugation).

Erythrocytes were only a minor problem as contaminants and only then if present in such large numbers that auto rosettes formed and lymphocyte density was affected by cell to cell association. This was detected before cell separation on density gradients and the sample was then rejected.

h. ^3H -thymidine incorporation

Proliferating lymphocytes are actively involved in DNA production and as such will incorporate labelled thymidine. The degree of proliferation may be determined from the rate of incorporation of the labelled thymidine. The incorporation of ^3H -thymidine by density gradients fractions was investigated to compare proliferation within the separated sub-populations.

Lymphocytes (30×10^6), from a healthy donor, were re-suspended in 10ml TC 199 containing 10% heat-activated newborn calf serum and 2mmol glutamine. ^3H -thymidine (29 Ci/mmol) was added at $2\mu\text{Ci}/\text{ml}$. Incubation at 37°C for one hour was followed by two washes with CMFSS. The cells were resuspended in 2ml of 1% Ficoll, incubated at 37°C for 30 minutes and then separated on the Ficoll density gradient. The harvested fractions were washed, counted, adjusted to $10^6/\text{ml}$ with CMFSS and 200 μl aliquots taken. The ^3H -thymidine incorporation of each aliquot was determined as in Chapter 2.2i. The results (Table 3.4) show that fractions C, D and, to a lesser extent, E incorporate more ^3H -thymidine than the denser fractions, indicating that these low density fractions contain the bulk of the proliferating cells in

³H-THYMIDINE INCORPORATION OF A HEALTHY SUBJECT'S
FRACTIONS FROM A FICOLL DENSITY GRADIENT.
(MEAN OF TWO EXPERIMENTS)

<u>Fraction</u>	<u>³H-thymidine incorporation (d.p.m)</u>
A	290
B	409
C	930
D	905
E	581
F	352
G	208
H	71

(All figures for incorporation of
2 X 10⁵ cells for one hour)

peripheral blood. Similar results have been published by August et al (1970) on their studies of human tonsil lymphocytes fractionated on BSA density gradients. This assay has been called the To test by Dicke, Tridante and van Bekkum (1969) in their studies of haematopoietic stem cells from mouse spleens and August et al (1970) reported that the distribution of human tonsillar lymphocytes with high To tests was almost identical to that obtained with mouse spleen. Low density lymphocytes, incorporating ^3H -thymidine at a high rate, have also been found in human thoracic duct lymph by discontinuous Ficoll density gradient analysis (Yu et al, 1973). In vitro generated blasts, using mouse spleen cells stimulated by mitogens, have been shown to have a low density and incorporate ^3H -thymidine. (Steinman, et al, 1978).

i. Density distribution of mitogen-stimulated lymphocytes

Mitogens, such as phytohaemagglutinin (PHA), concanavalin A (con A) and Pokeweed mitogen (PWM) have been used to activate lymphocytes non-specifically in vitro. Once activated those cells then transform into large lymphocytes which morphologically resemble those found as a result of in vivo antigen stimulation. Many of the morphological (Douglas, 1971; Hirschhorn et al, 1963; Janossy and Greaves, 1971) and biochemical changes (Naspitz and Richter, 1968; Ling, 1968) have been investigated and shown to have parallels with antigen stimulated cells.

From a practical point of view, mitogens are of value because large numbers of lymphocytes may be stimulated compared to specific antigens. Low dose PHA has been shown to preferentially stimulate a sub-population of T cells (Hirschhorn et al, 1971) and recent

work by Morgan (1978) has shown that in excess of 30% large lymphocytes may be found after three days of culture with this mitogen. Mouse spleen lymphoblasts stimulated to divide in vitro by lipopolysaccharide (LPS) Foetal Calf Serum (FCS) and Con A, acquire a low density and may be separated by isopycnic techniques (Steinman et al, 1978). Human lymphoblasts stimulated in vitro, have a high cytoplasmic:nucleus ratio (Ling, 1968) and should have a low density. Thus, this system was used to prepare large numbers of activated lymphocytes to assay the ability of the Ficoll density gradient to separate activated from non-activated cells.


Lymphocytes (30×10^6) from a healthy subject, isolated by isopycnic centrifugation on Ficoll Paque were incubated with purified PHA at $1\mu\text{g/ml}$ (this contains relatively more mitogenic activity and less agglutinating activity than the crude preparation) as described in chapter 2. After three days in culture the cells were sedimented, washed twice in CMFSS, resuspended in 16% Ficoll and separated on a Ficoll gradient. Simultaneously, more lymphocytes from the same donor were prepared fresh and separated on an identical gradient. The fractions were harvested, washed, counted and cell smears made on a cytocentrifuge.

The density distribution profiles are shown in Fig. 3.7. The peak of cells in fraction G is much reduced in a PHA activated population and a new peak appears at C. The intermediate fractions D, E, and F, show that PHA activation is not entirely synchronous, as new cells are continuously being recruited (Steinman et al, 1978). The large number of less dense cells shows that this technique of density gradient separation is sensitive to activation of lymphocytes.

Fig. 3.7.

The effect of culturing with PHA
on the density of lymphocytes in a
Ficoll gradient

30 ml of blood was taken and the lymphocytes prepared aseptically (Chapter 2.2.b.ii). The lymphocytes were cultured in 5% CO₂ at 37°C at 1×10^6 ml with PHA ($1\mu\text{gml}^{-1}$) for three days. The cells were washed X2 with CMFSS and separated on Ficoll gradient. Freshly prepared autologous lymphocytes were also separated in a parallel gradient. The gradient fractions were harvested and the lymphocytes counted. The results are expressed as percentage of total recovered cells in each fraction.

Fig. 3.7.

The cytocentrifuged smears were examined after staining with Wrights stain. Again, a cytocentrifuge was used to prepare the slides as the shearing forces resulting from a slide over slide smear cause cell damage particularly to large lymphocytes.

The results of differential counts are shown in Table 3.5. The larger cells are primarily of low density and the smaller cells are of high density. The density distribution is related to the size of the cell, but quantitative measurements were not possible because although cytocentrifuged slides help to keep cells intact they are distorted in the process and micrometric measurement cannot be accurate. The cells were, instead, graded on gross histology. Plasma cells were not seen in any preparations, neither were polymorphonuclear leucocytes or monocytes/macrophages.

In this experiment the Ficoll density gradient method described has been shown to be able to enrich and separate large lymphocytes from small lymphocytes.

j. Monocytes

Ficoll-Paque and Ficoll/Triosil are good media for separating lymphocytes from whole blood, almost all red cells and polymorphonuclear leucocytes are removed. However, monocytes are still present in the lymphocyte preparation and need to be removed before density gradient analysis. This is because the density of human monocytes (S.G. ~~1.055~~ 1.055) is within the range of lymphocyte densities, (S.G = 1.050 - 1.080) (Dacie and Lewes, 1963). Investigations of allograft rejection in mice by Roberts and Hayry (1977) have shown monocytes to distribute within the lymphocyte density range but not in the low density fractions.

Table 3.5

DIFFERENTIAL COUNTS OF DENSITY GRADIENT FRACTIONS OF
PHA-STIMULATED LYMPHOCYTES

<u>Fraction</u>	<u>Lymphocyte Size</u>		
	<u>Small (%)</u>	<u>Intermediate (%)</u>	<u>Large (%)</u>
A	2	21	77
B	5	21	74
C	5	28	67
D	12	27	61
E	29	37	34
F	60	29	11
G	86	14	0
H	100	0	0

Elliot, Haskill and Axelrad (1975) have shown that mouse macrophages sediment at a rate (10-15mm/h) within the lymphocyte range as shown by velocity sedimentation. Monocytes are actively phagocytic, a property not shared by lymphocytes. Monocytes also adhere readily to glass, which lymphocytes do not, although B cells adhere to a certain extent. These properties have been used to separate monocytes from lymphocytes in order to purify the lymphocytes prior to density gradient analysis to check on any interference of monocytes in the density profile.

..... Lymphocytes were obtained from peripheral blood by two means.

In the first method carbonyl iron was incubated with whole blood (chapter 2.2b) for 30 minutes at 37°C. The suspension was then left to sediment on a magnet. The supernatant was applied to Ficoll/Triosil (chapter 2.2b). The lymphocytes were harvested from the interface after centrifugation, washed twice with CMFSS and separated on a Ficoll density gradient.

In the second method lymphocytes were prepared from whole blood solely on Ficoll/Triosil and washed twice with CMFSS. 60% of the cells were pelleted resuspended in 10ml of Eagles medium containing 10% heat inactivated mouse serum and incubated in Falcon flat plastic flasks (50ml capacity) for three hours at 37°C. The flask was gently swirled and the cell suspension decanted. This contained non-adherent cells. Adherent cells were removed by adding 5ml CMFSS to the flask and gently rubbing the flask's interior surface with a polypropylene-covered spatula prior to decanting. This washing and rubbing was repeated to harvest as many adherent cells as possible. These two populations were spun down, washed twice with CMFSS, resuspended in 16% Ficoll and immediately isolated on Ficoll density gradients. The remaining lymphocytes from the Ficoll/Triosil

separation were resuspended in 16% Ficoll in a glass tube and incubated for 30 minutes at 37°C prior to separation on a Ficoll density gradient.

The four density gradients were centrifuged simultaneously, the fractions harvested, washed and counted. The results are shown in Table 3.6. The adherent cells equilibrated predominantly in fractions A, B, C, and D. The relatively high numbers in fraction G are probably contaminating small lymphocytes which had not been removed in the initial decanting of non-adherent cells. These are probably B cells which are believed to adhere by 'physical' means as opposed to the active adherent process of monocytes. It has been shown that by reducing the temperature to 4°C to inhibit cell metabolism, B cells still adhere, whereas monocytes will not (Shortman et al, 1971). This principle has been used to enrich plaque-forming (Plotz and Talal, 1967) and erythrocyte-binding (Wigzell and Makela, 1970) cells.

The three methods of monocyte depletion produce very similar density gradient profiles on the resulting population. Stained smears from those three populations showed no identifiable monocytes (1,000 cells were examined in each sample).

To confirm this observation a monocyte-specific stain was used, usually referred to as non-specific esterase. The staining is due to the specific classes of enzymes in the cytoplasm of monocytes. The method used was based on the original work of Yam, Li and Crosby (1971).

METHOD

The stain was applied to cytocentrifuged-prepared smears,

table 3.6

MONOCYTE AND LYMPHOCYTE DENSITY DISTRIBUTION(leucocyte source - healthy subject)Means of monocyte depletion

<u>Fraction</u>	<u>3 hours incubation in Eagles</u>		<u>30 minutes inc. carbonyl</u>	<u>iron</u>
	<u>Non-adherent</u>	<u>Adherent</u>	<u>in 16% Ficoll</u>	
A	0.1	18.2	0.2	0.2
B	0.1	31.8	0.3	0.3
C	1.0	4.5	1.2	0.3
D	2.4	9.1	3.5	1.0
E	8.6	0	6.0	4.0
F	11.6	0	7.8	8.3
G	75.0	27.3	75.0	79.9
H	1.2	9.1	6.0	6.0

Adherent cells \bar{x} 9.6% of total leucocytes

air dried and fixed in buffered formal acetone.

Stain Preparation

- A) 50mg. α -naphthyl acetate was dissolved in 3ml methoxy ethanol and added to 40ml phosphate buffer pH 7.6.
- B) 1ml 4% Sodium nitrite was added to 1ml pararosanilin and shaken for one minute.

Solutions A and B were mixed, adjusted to pH 6.2 using 2N HCl, filtered into a coplin jar, the slides immersed for one hour and then counter-stained with methyl green (1%) for 45 seconds.

The α -naphthyl acetate acts as a substrate for the specific enzymes and monocytes have a diffused brick red cytoplasm upon staining. The only other cells which may stain are some T cells which sometimes have a small dense staining body, but not the characteristic diffuse staining of the monocyte.

Samples of cell suspensions were tested before and after adherence and density gradient fractions were also analysed. (table 3.7). Adherence adequately removed the monocytes, but the adherent population had only a few contaminating non-myeloid cells, a point to be remembered as the loss of some cells at this stage would not have allowed representative density distribution analyses of lymphocytes. Non-specific esterase staining of the Ficoll gradient fractions showed that very few monocytes contaminated the gradients (less than 1% of the cells). As expected, these few cells which have a high cytoplasm/nucleus ratio, were found in the low density fractions B and C. The maximum contamination (3% of fraction B) was regarded as being minimal and not high enough to distort either the lymphocyte density distribution

Table 3.7

MONOCYTE DEPLETION FROM MONONUCLEAR
CELL SUSPENSION BY ADHERENCE TO GLASS
AND PLASTIC

a) 30 minutes incubated in flask

	<u>Pre-incubation</u>	<u>Non-adherent</u>	<u>Adherent</u>
%NSE +ve *	4	< 1	> 95

b) 30 minutes incubation prior to
Ficoll gradient separation

Fraction	Pre-inc.	A	B	C	D	E	F	G
%NSE +ve * per fraction	5	0	3	< 1	0	0	0	0

* NSE +ve = Non-specific esterase staining positive

data or the surface marker studies.

Carbonyl iron depletion produced a relative depletion of cells in the low density fractions, C. D. and E. This is a more physically demanding method of separation than adherence and the more fragile cells may well be damaged during the 30 minutes agitation with the carbonyl iron. Large lymphocytes are fragile so this may explain the relative loss of cells from the low density fraction. Incubation with carbonyl iron does not produce such a high lymphocyte yield from blood as Ficoll/Paque or Ficoll/Triosil (chapter 2.).

Adherence, thus, seems to be an adequate means of monocyte removal and 30 minutes at 37°C in 16% Ficoll in a glass tube seems adequately effective.

CHAPTER FOUR

ENUMERATION OF CIRCULATING

ACTIVATED LYMPHOCYTES IN

RHEUMATOID ARTHRITIS

ENUMERATION OF CIRCULATING ACTIVATED LYMPHOCYTES IN RHEUMATOID ARTHRITIS

4.1 INTRODUCTION

Increased numbers of immunoblasts have been shown in peripheral blood during infection (Wood and Frenkel, 1967) and immunization (Crowther, Fairley and Sewell, 1969). Previous workers have shown increased numbers of immunoblasts in the peripheral blood in R.A. (Bacon, Crowther and Sewell, 1975), ankylosing spondylitis (Eghtedari, Davis and Bacon, 1976) and polymyalgia rheumatica (Eghtedari, Esselinckx and Bacon, 1976). In R.A. the high incidence of immunoblasts correlated with increased disease activity and early onset of the disease. No studies on the effect of drugs on immunoblast numbers in R.A. have been reported. In almost all cases of lymphoid cell examination the blood (or a lymphoid cell-enriched preparation) was smeared, air dried and then differentially stained. Identification of immunoblasts is relatively easy; they are large (12-25 μ m diameter) have a high cytoplasm: nucleus ratio, prominent nucleoli and intensely basophilic cytoplasm. Occasionally, methyl green/pyronin has been used to highlight the pyroninophilic cytoplasm, mainly to eliminate cells of the monocyte/macrophage series.

Enrichment of the mononuclear cell population helps to prevent confusion of PMN precursors as immunoblasts. Some workers have removed monocytes/macrophages from mononuclear cell preparations by using adherence (Panayi, 1976) or by ingestion of iron particles (Eghtedari, Davis and Bacon, 1976).

DNA synthesis has been used to identify proliferative cells, usually uptake of tritiated thymidine. However, only about 30%

of immunoblasts are in the DNA synthetic phase of cell growth (Crowther, Fairley and Sewell, 1969). Autoradiography of smeared labelled cells is particularly useful for identification of the proliferating population, while scintillation counting of labelled cell suspensions is purely quantitative.

Lymphocyte size distribution has been used to compare healthy controls and patients with immunodeficiency status (Heyn, Tubergen and Althouse, 1973) and juvenile rheumatoid arthritis (Brungard, Moser and Althreya, 1975). In both reports the patients had more large cells than the controls.

Density gradients have been used for lymphocyte fractionation, the density of small lymphocytes being invariably higher than the density of larger cells (Gorczynski, Miller and Phillips, 1970). Systemic ^{lupus} erythematosus (SLE) patients peripheral blood lymphocytes have been compared with healthy controls on Ficoll density gradients (Glinski et al, 1976). To date, no report has appeared concerning the density distribution of R.A. peripheral blood lymphocytes.

A study was therefore undertaken to determine immunoblast numbers, lymphocyte size distribution and density distribution analysis in rheumatoid patients and to relate these parameters to disease activity and chrysotherapy.

4.2 MATERIALS AND METHODS

4.2.a. PATIENTS AND CONTROLS

All patients were receiving only analgesic or non-steroidal anti-inflammatory drug (NSAID) treatment unless otherwise stated. Patients receiving gold, penicillamine, cortico-steroid or cytotoxic therapy were excluded, as were all patients with infections. Controls were healthy hospital staff, age and sex matched for R.A. patients wherever possible, particularly for the density gradient analysis.

4.2.b. LYMPHOCYTE PREPARATION AND MICROSCOPIC EXAMINATION

Venous blood was collected and defibrinated or heparinized (Chapter 2.2.a.). Lymphocytes were prepared by separation with methyl cellulose/Carbonyl iron and/or Ficoll-Triosil (Chapter 2.2.b) cell smears were made and differentially stained with Wrights stain (Chapter 2.2.d.) and examined under a binocular microscope.

Immunoblasts, identified as outlined in 4.1, were expressed as a percentage of lymphoid cells, at least 200 cells being counted.

Lymphocyte diameters were measured using an eyepiece graticule which had been calibrated with a stage graticule, 100 cells being measured in each instance.

In any method of purifying lymphocytes from whole blood there is cell loss. More important, there may be a selective loss of certain sub-populations. Johnson and Madsen (1978) found no difference in human lymphocyte sub-populations in heparinized and defibrinated blood. I had a greater total loss of lymphocytes in defibrinated blood and eventually used heparinized blood for all experiments.

Thomson, Bull and Robinson (1966) found that activated lymphocytes were particularly susceptible to damage and death during lymphocyte preparation. Loos and Roos (1974) found a loss of activated lymphocytes from mouse spleen cell preparations when purifying the lymphocytes. This problem was investigated in this study by assaying the loss of mitogen-stimulated cells in a Ficoll-Paque separation so that I could enumerate any loss of activated cells.

Peripheral blood lymphocytes prepared by Ficoll/Paque isopycnic centrifugation (Chapter 2.2.b.) from four healthy donors were incubated with PHA at $1\mu\text{g}/\text{ml}$ for three days (Chapter 2.2.i.). An aliquot of these cells (1×10^6) was taken for ^3H -thymidine incorporation. The remainder (approx. 10×10^6) were subjected to isopycnic centrifugation on Ficoll-Paque at $2 \times 10^6/\text{ml}$. The interface cells were washed and an aliquot (1×10^6) taken for ^3H -thymidine incorporation. The relative incorporation was determined by equal cell numbers before and after Ficoll-Paque purification. The results (Table 4.1) show that although there was a large cell loss associated with Ficoll-Paque separations, this did not cause a relative depletion of proliferating lymphocytes as estimated by ^3H -thymidine incorporation. In fact there was an increase in the incorporation per cell. This is probably due to the decreased cell density of the mitogen stimulated population. The unstimulated cells, which are denser, pass through the Ficoll-Paque and are not recovered. Thus, there is no selective loss of proliferating lymphocytes with Ficoll-Paque. This conclusion has its limitations if extended to the separation of in vivo stimulated blasts from blood. The slight leucoagglutinating activity of the purified PHA may prevent the stimulated cells from passing into the Ficoll-Paque during centrifugation; also the agglutination may be protective

TABLE 4.1

RECOVERY OF PROLIFERATING CELLS ATFICOLL-PAQUE INTERFACES

Subject	Lymphocyte numbers (10^6)			Radioactivity (^3H) (dpm)		
	Pre F-P	Post F-P	% recovery	Pre F-P	Post F-P	% recovery
PM	7.9	2.1	27	31031	39375	126.9
NM	7.4	3.6	48	31263	39729	127.1
BS	9.7	4.0	41	44813	50850	113.5
JC	11.2	4.2	38	35284	39492	111.9
					Mean	119.9

to the proliferating cells. However, these are imponderables and it would seem that it is reasonable to use Ficoll-Paque to prepare samples of lymphocytes which contain activated cells.

4.2.c. LYMPHOCYTE VOLUME ANALYSIS

Lymphocytes were prepared by isopycnic centrifugation on Ficoll-Trisil (Chapter 2.2.b2) and monocytes depleted by adherence at 37°C for 30 minutes (Chapter 3.3.c.)

Contaminating erythrocytes were removed by repeated washing in 0.83% ammonium chloride. The cells were washed three times with PBS which had been rendered particle-free by two passages through 0.45µm Millipore filters. During these washings the cells were pelleted at 200g to reduce contamination by platelets. Cell viability assessed by dye exclusion was >95%. Cell volume was determined with a celloscope model B. A probe with a pore diameter of 60µm was immersed in the washed lymphocyte suspension (C. 5×10^4 lymphocytes/ml). The suspension was sucked into the pore and as each cell passed through the pore it caused a fluctuation in the movement of the mercury manometer used to draw in the suspension by negative pressure. The small movement of the mercury column was recorded as a change of electrical potential between two electrodes. This was detected by a mercury manometer and the results were accumulated by a multi-channel data analyser (Nuclear Data, Reading) and were displayed as a volume distribution curve on a cathode ray tube. Once a sufficient number of cells had been measured to produce a large distribution curve this data was printed on to graph paper via a flat bed pen recorder. Approximately 5,000-8,000 lymphocytes were measured for each sample.

4.2.d DENSITY DISTRIBUTION ANALYSIS

The methodology for this is described in Chapter 3 Section 3. Gradient tubes were run in parallel in every case, taking as much care as possible in age matching the control (who was always sex matched) for the patient. The patients disease activity was clinically assessed by Dr. P. A. Bacon (Consultant Rheumatologist) as being broadly "active" or "inactive". A large number of the active patients were new admissions to the R.N.H.R.D. who had received no previous "second-line" drugs, i.e. had never had any other drugs apart from analgesics or NSAIDs. Inactive patients were those whose disease was well controlled by NSAID or were in remission.

A number of patients (8) receiving gold therapy were also used. One patient (P.H) who had not responded to any drug therapy was subjected to whole body irradiation (500 rad) and the effect of this treatment on his peripheral blood lymphocytes density distribution was monitored. A patient with sarcoidosis was also tested: he had been diagnosed as an early rheumatoid at the time of testing.

4.3 RESULTS


a) IMMUNOBLAST NUMBERS

Lymphoid cells with the morphology of immunoblasts in my preparations were rarely of the great size reported by other authors (Horwitz et al, 1970; Delbarre et al, 1975). Many lymphoid cells which had the cytological characteristics of immunoblasts were only 10 μ m diameter, cells above 20 μ m were only seen in a few exceptional cases, even in patients with chest infections and infectious mononucleosis. Nearly all authors characterize immunoblasts as being above 15 μ m diameter, in fact Delbarre et al (1975) judged immunoblasts to be 20-30 μ m diameter. In this study immunoblasts were defined as being above 10 μ m and possessing a kidney-shaped, eccentrically placed nucleus with prominent nucleoli, a high cytoplasm/nucleus ratio and hyperbasophilic cytoplasm. However, to include an analysis of increased cell size, lymphocyte diameters were measured (see next Section). During the preparation of cell smears surface tension can result in large cells being carried towards the edges and tail of the smear. Thus, when examining the stained smears, each was systematically scanned to ensure that the data was truly representative. The percent incidence of immunoblasts in controls and active and inactive R.A. is shown in Fig. 4.1. Increased immunoblasts are shown predominantly in active disease with more than half of these patients having immunoblasts raised above the normal level (0.5%). Only four of 29 patients with inactive R.A. had raised levels of immunoblasts and then only to 1% of the lymphoid cell population.

Fig. 4.1

The incidence of immunoblasts
in the peripheral blood lymphoid population
of healthy and rheumatoid subjects

Peripheral blood lymphocytes were prepared from 10 ml venous blood (chapter 2.2.b.ii) and stained smears examined microscopically. The immunoblast was recorded as a mononuclear cell about 10 μ diameter with a large eccentrically placed, kidney-shaped nucleus which possessed two or more nucleoli. A high cytoplasm: nucleus ratio and hyperbasophilic cytoplasm were also used as positive criteria. The results are expressed with the immunoblasts as a percentage of the lymphocyte population. Each point is for one subject.

Fig. 4.1.

b) LYMPHOCYTE DIAMETERS

The eyepiece graticule allowed accuracy of $0.7\mu\text{m}$ in the measurement of lymphocyte diameters. Again, the slides were systematically scanned and only intact cells were measured. The results are shown in Fig. 4.2.a. In both rheumatoid (22 patients) and healthy subjects (17 patients) most lymphocytes (90%) are in the normal range. The main difference between the two subject groups is at the high end of the range (4.2.b). R.A. patients have more lymphocytes above $10\mu\text{m}$ diameter, 3.1% compared with 0.7% of $11\mu\text{m}$ diameter for healthy subjects. Healthy subjects did not have lymphocytes above this size whereas R.A. patients had 0.75% circulating lymphocytes of $11.8\mu\text{m}$ diameter. Lymphocytes above $12.5\mu\text{m}$ were exceptional ($\leq 0.1\%$).

The results do not show a great increase in the numbers of large lymphocytes in R.A. However, this method of measurement is long and subjective and measures cell volume as found on a stained, dried smear. As a result distortion occurs and the larger cells in R.A. did not always have the morphological characteristics of blasts.

c) ELECTRONIC SIZING

The equipment employed (a celloscope) measures the volume of cells in suspension. For this experiment it was assumed that lymphocytes in a physiological medium are spherical. Apart from measuring more cells than can practically be measured by microscopy, this system measured the cells alive and undistorted by the various forces inherent in the making of cell smears. The number of lymphocytes measured is indeterminate and hence renders statistical analysis difficult. For this reason the y axis is marked as percentage peak height.

A total of eight R.A. patients and seven healthy subjects were used in this experiment. The results shown in Fig.

4.3. are representative of five of the six patients with active R.A. In the other three patients the distribution profiles for R.A. and control were similar. The abnormal pattern shown for the R.A. patient shows an increase in the number of cells of above average volume. In fact, up to 10% of the peak number are found in channels 265-293. This pattern of results is similar to the more qualitative figures shown in Fig. 4.2.

The pattern of lymphocyte size was not considered to demonstrate an adequate number of large lymphocytes to permit their separation by velocity sedimentation. With the knowledge that lymphocyte density is changed before a significant increase in cell size is recorded (Steinman et al, 1978) the separation of lymphocytes by density was performed.

d) LYMPHOCYTE DENSITY DISTRIBUTION

The results of a comparison between 23 healthy subjects and 23 R.A. patients are shown in Fig. 4.4. In healthy subjects most of the lymphocytes (67%) were found in the dense fraction (G), with decreasing incidence in the less dense fractions.

Compared with healthy subjects the R.A. patients, selected from the whole spectrum of clinical activity, had a depletion of dense cells and increased numbers of low density cells. Fraction G contained only 50.5% of the lymphocytes in R.A. by comparison with 67% in healthy subjects. Fraction F contained 21% and 11%, fraction E 12.5% and 8%, fraction D 6.5% and 4% and fraction C 4% and 3% respectively.

Fig. 4.2

The mean diameter of peripheral blood lymphocytes
in healthy and rheumatoid subjects

Peripheral blood lymphocytes were prepared from venous blood (Chapter 2.2.b. ii) and stained smears examined microscopically. 100 lymphocyte diameters were determined with an eyepiece graticule whilst systematically scanning the slide.

The results are expressed as percentage lymphocytes of a given diameter.

a) shows the full range of results

b) highlights the difference between R.A. and healthy subjects at the higher end of the range.


Fig. 4.2.a


Fig. 4.2.b

Fig. 4.3

Lymphocyte volumes of healthy and
R.A. subjects (measured with a Celloscope).

Lymphocytes were prepared from peripheral blood
(Chapter 2.2.b.ii) and washed twice with filtered
PBS. Cell volume was determined using a Celloscope (model
B) (4.2c). The results are expressed as percentage
of peak height in each channel.


Fig. 4.3.


Fig. 4.4

Density distribution profiles of peripheral blood
lymphocytes from healthy and R.A. subjects

Peripheral blood lymphocytes were prepared from 30 ml venous blood (chapter 2.2.b.ii). Monocytes were depleted (Chapter 2.3.c.) and the lymphocytes separated by density in a Ficoll gradient. The fractions were harvested and counted (Chapter 3.3.d.). R.A. patients were analysed in parallel with age and sex matched healthy subjects. The results are expressed as the mean percentage of total recovered lymphocytes in each fraction \pm 1 standard deviation (S.D.).

The differences between R.A. and healthy subjects were statistically significant (Student t-test) in the following fractions:

<u>Fraction</u>	<u>p</u>
A	< 0.02
B	NS
C	< 0.01
D	< 0.001
E	< 0.01
F	< 0.001
G	< 0.001
H	NS

Fig. 4.4.

The proportions in fractions A and B were similar for R.A. and healthy subjects. The differences between R.A. and healthy lymphocyte percentages were highly significant in fractions D, E, F and G ($P < 0.001$).

Upon comparing a separate group of patients with clinically active R.A. (n=11) with healthy subjects (n=23) the results were more strikingly different (Fig. 4.5). The number of lymphocytes in fraction G was further reduced to 37% and fractions C, D, E and F increased to 5.5%, 8.0%, 16.0% and 27.5% respectively. From this result it can be seen that it is mainly the active patients who have the increased number of low density cells. The comparisons between active R.A. and healthy subjects are highly significant in fractions C, D, E, F and G ($P < 0.001$).

The density distribution profiles of a separate group of 12 clinically inactive patients peripheral blood lymphocytes are shown in Fig. 4.6 and shown to be dissimilar to patients with active R.A. In fact fraction G contains 63% of the lymphocytes, almost as many as in healthy subjects (67%). The fractions E and F each contain less than 15% of the lymphocytes and C and D are less than 5% each of the total. The differences with clinically active patients are significant in fractions D ($P < 0.01$), C and E ($P < 0.02$) and F, G and H ($P < 0.001$).

Conversely, eight patients receiving gold therapy had a mean density distribution pattern similar to that for healthy subjects

Fig. 4.5

Density distribution profiles of peripheral
blood lymphocytes from healthy subjects
and clinically active R.A. patients.

Peripheral blood lymphocytes from healthy subjects and clinically active R.A. patients were separated by density as described in the legend to Fig. 4.4 and the results presented accordingly.

The differences between clinically active R.A. and healthy subjects were significant in the following fractions:

<u>Fraction</u>	<u>p</u>
A	NS
B	NS
C	< 0.001
D	< 0.001
E	< 0.001
F	< 0.001
G	< 0.001
H	NS



Fig. 4.5.

Fig. 4.6

Density distribution profiles of peripheral
blood lymphocytes from R.A. patients with
clinically active and inactive disease.

The lymphocytes were prepared and separated by density as described in the legend to Fig. 4.4. The results are presented in the same format, as Fig. 4.4, and statistically significant differences between the patient groups were recorded as follows:

<u>Fraction</u>	<u>p</u>
A	NS
B	NS
C	< 0.02
D	NS
E	< 0.02
F	< 0.001
G	< 0.001
H	< 0.001

Fig. 4.6.

(Fig. 4.7). Differences between the fractions for the two subject groups were not significant. Compared with active R.A. patients (Fig. 4.8), the patients on gold had fewer cells in fractions C and D (5%) E (7%) and F (12.5%). Fraction G contained 68% of the lymphocytes - a very similar figure to the healthy value (67%).

Total body irradiation of a clinically active R.A. patient (P.H) in two half-body treatments caused a shift of the lymphocyte density distribution profiles towards the low density fractions (Table 4.2). The peak of lymphocytes on day 0 (48%) was in fraction G, but at three days (two days post-irradiation) was in fraction F (78.3%) and after the second irradiation, at 14 days, was in fraction E (48.7%). At 31 days and 93 days (17 and 79 days after the second irradiation) the distribution profile was shown to be slowly resuming to that recorded before treatment, at 93 days the peak (59.1%) being in fraction F.

The patient with sarcoidosis had the most abnormal density distribution profile of the study. The majority of the lymphocytes (58%) were found in fraction E in contrast with only 9% for healthy subjects. Fraction G was reduced to 5% of the total compared with 67% for healthy subjects. Fractions C and D had increased cell numbers relative to healthy subjects (Fig. 4.9.).

Fig. 4.7

Density distribution profiles of peripheral
blood lymphocytes of healthy subjects and
R.A. patients receiving gold therapy.

The lymphocytes were prepared and separated by density as described in the legend to Fig. 4.4 and the results presented in the same format. There was no statistically significant difference between the two subject groups.


Fig. 4.7.

Fig. 4.8

Density distribution profiles of peripheral
blood lymphocytes from R.A. patients with
clinically active disease or receiving gold
therapy.

The lymphocytes were prepared and separated by density as described in the legend to Fig. 4.4. The results are expressed in a similar manner and statistically significant results were shown in the following fractions:

<u>Fraction</u>	<u>p</u>
A	NS
B	NS
C	< 0.05
D	< 0.05
E	< 0.01
F	< 0.001
G	< 0.001
H	NS


Fig. 4.8.

TABLE 4.2

DENSITY DISTRIBUTION PROFILES OF A CLINICALLY
ACTIVE RHEUMATOID ARTHRITIS PATIENT (P.H.). THE EFFECT OF
WHOLE BODY IRRADIATION.

<u>Fraction</u>	Days Post-irradiation				
	<u>0</u>	<u>3</u>	<u>19</u>	<u>31</u>	<u>93</u>
A	0.8	0.5	0.4	0	0.2
B	0.7	1.1	2.1	0.7	0.7
C	2.5	1.4	6.3	4.4	0.7
D	6.2	1.6	28.4	9.9	1.0
E	14.8	10.5	48.7	37.8	2.0
F	22.3	78.3	11.1	34.6	59.1
G	48.5	5.5	2.8	12.7	35.5
H	4.1	1.1	0.2	0	0.8


Two half body irradiations (500 rad) on day 1 and day 14.

Figures = % total recovered cells in
each fraction

Fig. 4.9

Density distribution profile of a patient
with sarcoidosis. Comparison with healthy
subjects.

The peripheral blood lymphocytes were prepared from
..... venous blood and separated by density as in the legend
to Fig. 4.4

Fig. 4.9.

4.4 DISCUSSION

The activation of peripheral blood lymphocytes investigated in this chapter will be discussed in relation to the surface marker studies in Chapter 6. This discussion will deal only with the results of the patient who was subjected to whole body irradiation and the patient with sarcoidosis.

Whole body irradiation resulted in an increase in the numbers of low density lymphocytes from two days up to 18 days post-irradiation. Bond et al (1961) found a large decrease in the percent DNA synthesizing cells 24 hours after irradiation of man at a similar level (500 rads). However, the number of those proliferating cells were increased above the normal level from two days onwards and were still supra-normal after 50 days. Normal levels were attained at 141 days. These workers also found a drop in DNA synthesizing cells in irradiated dogs for 24 hours, followed by a sharp increase which peaked after seven days and had returned to normal levels after 14 days. The mitotic index of bone marrow also rises after an initial fall (Fleidner et al, 1959). For irradiation to have such a lasting effect, as shown by Bond et al (1961) and in this study, more than circulating cells were being affected. The bone marrow is certainly affected and the increased numbers of the proliferating population and the low density population may be due to increased bone marrow output. Edgren et al (1976) showed that T lymphocytes, irradiated extra-corporeally, were destroyed or incorporated into the intravascular lymphoid pool upon re-infusion. This was also reflected by a decrease in the number of circulating T cells, suggesting that T cells are more radiosensitive than B cells.

Whilst the density distribution profile was not, in this study, determined within 24 hours of irradiation, it is the proliferating population of lymphocytes which are believed to be the most radiosensitive (Waithe and Hirschorn, 1973). However, the shift in the densities of the circulating lymphocytes is such that this cannot be totally explained by an increase in bone marrow output. The change in density is more sensitive than any noted change in lymphoid morphology with respect to activation.

Sarcoidosis gave the most abnormal density distribution profile found in any disease state analysed in this study. This disease is characterized by a hypereactivity of the antibody producing system and a depression of delayed hypersensitivity. Both of these abnormalities could be due to antigen induced activation of both populations (Cole, 1977), an antigen having been found in sarcoid spleen (Kvein, 1941). Polyclonal activation could explain the density distribution profile.

CHAPTER FIVE

THE NATURE OF THE LYMPHOCYTE
SUB-POPULATIONS SEPARABLE BY
DENSITY IN HEALTH AND R.A.

5.1 INTRODUCTION

The previous chapter has shown a significant difference in the density distribution profiles of circulating lymphocytes from healthy and rheumatoid subjects. This difference was greater than the size differences shown by microscopy and electronic sizing, particularly for clinically active R.A. patients. Steinman et al (1978) believe that a lowering of density is one of the first indications of lymphocyte activation, which is reflected in an increased cytoplasmic refractive index (Metcalf et al, 1976). The preparative properties of discontinuous density gradient analysis allow further studies on the nature of the lymphocytes in the various fractions.

If the increased numbers of less dense lymphocytes shown in R.A. (Chapter 4) are activated cells it would be of great importance to find out their origin and function. Surface marker studies may help to do this, and this has been examined here.

Some lymphocyte sub populations retain their markers when activated, some may lose them. The literature shows that T cells do not lose their receptors for sheep erythrocytes upon activation and B cells do not lose integral surface Ig, but lose their C3 receptors and possibly their Fc receptors also (Jondal, 1974 b, c; Miyama et al, 1978). Conversely, activated T cells may acquire Fc receptors (Van Boxel and Rosenstreich, 1974), notably suppressor T cells (Moretta et al, 1977). The expression of surface markers by newly activated lymphocytes is unknown. Preparative continuous Ficoll gradient analysis allows examination of any such changes as a result of activation.

The data for T and B cell numbers in peripheral blood in R.A. is equivocal (Hoyerall, Froland and Wisloff, 1975; Brenner, Scheinberg and Cathcart, 1975; Froland, Natvig and Wisloff, 1975). The variations in results may well be due to technical differences in the assaying of both T and B lymphocytes.

Recent evidence has shown an increase in EA- rosette formation by circulating lymphocytes in rheumatoid arthritis (Sharpin and Wilson, 1977 a). This has been shown to be due to the binding of the Fc region of IgG, which is not due to cell bound rheumatoid factor. The enhanced EA- rosette formation by peripheral blood lymphocytes from R.A. patients occurred through a range of concentrations for sensitizing antibody, suggesting that the Fc receptors on R.A. lymphocytes are more avid for EA than receptors on lymphocytes from healthy subjects (Sharpin and Wilson, 1977 b). A similar EA- rosette system, using sensitized rabbit (Fawaz-Duhaut et al, 1975) or human erythrocytes (Bach, Delrieu and Delbarre, 1970; Sany et al, 1975) has detected increased EA- rosette formation in R.A. A number of authors (Williams et al, 1973; Brenner, Scheinberg and Cathcart, 1975) have commented on the increased numbers of circulating non T, non B (null) cells in R.A.

This chapter describes the results of studies designed to characterize the lymphocyte surface markers in Ficoll density gradient fractions. In both health and R.A., clear trends were shown in the incidence of surface markers within the lymphocyte density range.

5.2 MATERIALS AND METHODS

a) DENSITY DISTRIBUTION OF ENRICHED SUB-POPULATIONS

i) DENSITY DISTRIBUTION OF E-ROSETTE DEPLETED CELLS

T cells were depleted from the peripheral blood lymphocyte population of four rheumatoid patients as described in chapter 2.2e and the remaining cells subjected to Ficoll density gradient analysis (Chapter 3.2), in a comparison with autologous unseparated lymphocytes.

ii) DENSITY DISTRIBUTION OF EAC-ROSETTE DEPLETED CELLS

Cells bearing receptors for the C3 component of complement were depleted from the peripheral blood lymphocyte population of one R.A. patient by EAC- rosette depletion (Chapter 2.2g). The remaining cells were separated in a Ficoll density gradient (Chapter 3.2) in a comparison with autologous unseparated lymphocytes.

b) SURFACE MARKERS ON LYMPHOCYTES IN FICOLL DENSITY

GRADIENT FRACTIONS

Peripheral blood lymphocytes from R.A. patients and age and sex matched healthy subjects were fractionated, in parallel, in Ficoll density gradients (Chapter 3.2). The cells from each fraction were counted and washed three times with CMFSS.

i) E- ROSETTES

The lymphocytes from each fraction were pelleted and adjusted to 2×10^6 /ml using PBS. Equal aliquots (0.2ml) of lymphocytes and washed sheep erythrocytes were mixed and the percentage E- rosettes determined (Chapter 2.2e).

ii) EA- ROSETTES

The lymphocytes from each fraction were adjusted to

2×10^6 with CMFSS. Ox red blood cells were washed, coated with IgG antibody and rewashed and mixed with the lymphocytes to form EA- rosettes as described in Chapter 2.2f.

The concentration of coating antibody to be used in these experiments was determined by a number of titration experiments. The range of concentrations used was 1/200, 1/400, 1/800, 1/1600 1/3200. The highest concentration (1/200) was the strongest sub-agglutinating concentration. Whole lymphoid populations, both healthy and R.A., were tested in this way. From this range of titres two concentrations were chosen with the aim of determining high avidity EA- rosette formation and low + high avidity EA- rosette formation. These two concentrations were used for the determination of EA- rosette formation in Ficoll density gradients.

iii) SURFACE IMMUNOGLOBULIN

Surface immunoglobulin was assayed using fluorescein isothiocyanate (FITC) conjugated rabbit anti-human Ig as described in Chapter 2.2.h. To simplify the labelling and washing of large numbers of cell fractions the whole mononuclear cell population prepared on Ficoll/Paque was labelled with the FITC anti-Ig for 30 minutes at 4°C . However, only as much of the labelled conjugate was added to 30×10^6 lymphocytes as usually added to 1×10^6 cells. To assay the validity of this approach doubling dilutions of a lymphocyte suspension were prepared in PBS, pelleted at 200g and 0.2ml of the conjugate (1:5 in PBS) added to each pellet, mixed and incubated for 30 minutes at 4°C before reading under a UV microscope. The results (table 5.1) show that cell number does not appear to affect the efficiency of staining with the labelled conjugate.

TABLE 5.1

THE EFFECT OF CELL NUMBER ON THE LABELLING
OF LYMPHOCYTES WITH FITC- ANTI HUMAN Ig.

<u>Cell number (X10⁶)</u>	<u>% Surface Ig positive</u>
1	19
2	20
4	21
8	20
16	23
32	22

The lymphocytes were thus labelled before density gradient fractionation (chapter 3.2). The number of cells per fraction was determined and then the cells were pelleted for estimation of percent FITC- labelled cells in each fraction.

5.3 RESULTS

The R.A. patients examined in this chapter included the whole range of disease activity, patients with the more clinically active disease being chosen where possible.

a) ROSETTE DEPLETION

i) E- ROSETTE DEPLETION

The effect of E- rosette depletion on the lymphocyte density distribution profiles of four R.A. patients is shown in Fig. 5.1. In comparison with the autologous whole lymphoid population the E- rosette depleted population had a decrease in the number of dense cells and a proportional increase in the number of less dense cells. The E- rosette depleted population had a reduction of lymphocyte percentage from 52.5% to 21.5% in fraction G. Fraction F was reduced from 23% to 11%. The profile of E- rosette depleted cells showed a shift to the left and the peak was found in fraction E (28%), compared with the non E- rosette depleted cells. Fractions A, B, C and D, were also proportionally increased, fraction D being the most affected by the depletion, with the percentage of cells in this fraction rising from 6% to 20%. The differences in fractions B to G are highly significant. (C, D, E: $P < 0.02$; B, F, G: $P < 0.01$).

The inference from these results is that the low density R.A. lymphocytes in the Ficoll density gradient are predominantly non-T. The depletion of lymphocytes from Fraction G was great, only 21.5% of the remaining lymphocytes were found in this fraction. This may be a result of a loss of denser lymphocytes on reapplication to a second Ficoll-Paque separation, the step used to remove E-rosetted

Fig. 5.1.


The effect of depletion of E- rosetting
cells on the density distribution profiles
of peripheral blood lymphocytes in R.A.

60 ml of venous blood was taken and the lymphocytes separated (Chapter 2.2.b.ii). The E- rosetting cells were depleted from half of the lymphocyte sample (Chapter 2.2.e.) and the non E- rosetting cells (from the Ficoll/Paque interface) and the whole lymphoid population were separated in parallel Ficoll gradients (Chapter 3.3.d.).

The gradient interfaces were harvested, counted and the results expressed as a percentage of the total recovered lymphocytes in each fraction.

The differences between the separated samples were statistically significant (Students t-test) in the following fractions:

<u>Fraction</u>	<u>p</u>
A	0.002
B	0.01
C	0.02
D	0.02
E	0.02
F	0.01
G	0.01
H	NS

Fig. 5.1

lymphocytes. This possibility may be overcome by the direct testing of density gradient fractions for E- rosette formation. However, the rapidity of the former method allowed relatively quick assessment of the surface markers in the density gradient fractions; a relevant point considering the fragility of large lymphocytes.

ii) EAC- ROSETTE DEPLETION

The effect of EAC- rosette depletion on the lymphocyte density distribution of one R.A. patient is shown in Fig. 5.2. By comparison with autologous unseparated lymphocytes there was a depletion of lymphocytes from fractions F, G and H. Quantitation of this can only be of limited value as only one patient was tested. The implications are that relatively few of the low density lymphocytes have receptors for the C3 component of the complement. Conversely, many of the cells with C3 receptors would be found in fractions F and G. Due to the need to use two Ficoll-Paque separations, as described in the previous section, this approach was discarded. The relative lack of C3 receptors on activated B cells also reduced the value of this data to some extent.

b) DIRECT SURFACE MARKERS

The lack of sufficient numbers of lymphocytes, particularly in the low density fractions, necessitated restricting the analysis to one marker study per subject sample.

i) E- ROSETTES


The results for each patient are shown in Fig. 5.3 and the group mean values summarized in Fig. 5.4. In Fig. 5.3 the percentage of E- rosettes is seen to be increased in the denser fractions (fraction

Fig. 5.2

The effect of depleting EAC- rosetting cells
on the density distribution profiles of peripheral
blood lymphocytes in R.A.

60 ml of venous blood was taken and the lymphocytes separated (Chapter 2.2.b.ii). The EAC- rosetting cells were depleted (Chapter 2.2.g.) from half of the lymphocyte sample and the non EAC- rosetting cells (from the Ficoll/Paque interface) and the whole lymphoid population were separated in parallel Ficoll gradients (chapter 3.3.d.).

The gradient interfaces were harvested, counted and the results expressed as a percentage of the total recovered lymphocytes in each fraction. There was no statistical analysis.

Fig. 5.2

H excepted). The range of results is increased in the denser fractions, subjects varying by up to 38% in fraction G (40-78%). The mean of these results (Fig. 5.4) shows that both healthy and rheumatoid subjects display the trend of increasing E- rosette formation in the denser fractions. The difference between R.A. and healthy subjects is most noticeable in fraction G in which R.A. patients have a mean of 54% E- rosettes compared with 61% for healthy subjects. This difference is not statistically significant (Students t- test), neither are the small subject group differences in fractions B, C, D, E and F. The relative numbers of E- rosettes in the Ficoll gradient fractions are shown in Fig. 5.5. It can be seen that most of the E- rosettes are in fraction G, more so in healthy subjects. R.A. patients have relatively more E- rosettes in the low density fractions, suggesting some T cell activation. Accumulating the E- rosettes in the gradient fractions shows that healthy subjects' lymphocytes formed 51% E- rosettes compared to 47% for R.A. patients. Statistical analysis of this data is not possible as it is a function of two results, each with their own population deviations.

ii) EA- ROSETTES

The mean percentages of EA- rosettes formed by whole lymphoid populations from R.A. and healthy subjects at various erythrocyte antibody concentrations are shown in Fig. 5.6. At all concentrations the R.A. patients formed more EA- rosetting cells, suggesting more avid EA- rosette formation ($1/400$ $P < 0.02$; $1/800$ $P < 0.01$; $1/1600$, $1/3200$, $1/6400$ $P < 0.001$).

To analyse EA- rosette formation in the Ficoll gradient fractions two antibody coating concentrations were used.

Fig. 5.3

5.4

The density distribution of lymphocytes forming

5.5

E- rosettes in Ficoll density gradient fractions.

Comparison of healthy subjects and those with R.A.

Peripheral blood lymphocytes were prepared from 30 ml venous blood (Chapter 2.2.b.ii), the monocytes depleted (Chapter 3.3.c.) and the lymphocytes separated by density in a Ficoll gradient (Chapter 3.3.d.). The interface fractions were harvested, counted adjusted to $2 \times 10^6 \text{ ml}^{-1}$ and E- rosettes determined (Chapter 2.2.e.). The results (Fig. 5.3) are expressed as percentage E- rosetting lymphocytes in each fraction and each point represents the lymphocytes from one patient in each fraction.

The mean result (± 1 standard deviation of the mean) for each fraction is shown in Fig. 5.4. There were no statistically significant differences between R.A. and healthy subjects.

The E- rosetting lymphocytes of each fraction are presented with the mean total cell proportion in each fraction (Fig. 5.5.). This figure was obtained by multiplying the % incidence of E- rosettes by the proportion of lymphocytes in each fraction., e.g., in Fraction G for healthy subjects the E- rosettes are 41 of 67% (approx 2/3) of the total lymphocytes in the gradient.

Fig. 5.3

○ Healthy subjects

● R.A.





Fig. 5.4



Fig. 5.5.

Fig. 5.6

EA- rosette formation of healthy and
rheumatoid whole lymphocyte populations

Peripheral blood lymphocytes were prepared from venous blood (Chapter 2.2.b.ii). EA- rosette formation was determined using various titres of rabbit IgG anti ox red cell antisera (Chapter 5.2.b.ii). The results are plotted as the mean \pm 1 standard deviation. The differences between R.A. and healthy subjects were significant at the following concentrations:

<u>Concentration</u>	<u>Numbers tested</u>		<u>p</u>
	<u>R.A.</u>	<u>Healthy</u>	
1/200	8	8	NS
1/400	10	10	<0.02
1/800	10	10	<0.01
1/1600	34	19	<0.001
1/3200	26	18	<0.001
1/6400	7	7	<0.001

Fig. 5.6.

The 1/800 titre was used to detect medium (and high) avidity Fc receptor binding and the 1/3200 titre was used to detect only high avidity binding.

The results of the medium avidity analysis of Ficoll gradient fractions are shown in Fig. 5.7 and summarized in Fig. 5.8. EA- rosettes were formed at between 28% and 34% for both R.A. and healthy subjects in fractions B, C, D, E and F. The differences between the subject groups were not significant in these fractions, but were in fraction G in which R.A. had significantly reduced EA- rosettes ($p < 0.05$). The EA- rosette formation in fraction G was also reduced relative to other fractions, for both R.A. and healthy subjects, R.A. patients forming 18% EA- rosettes and healthy subjects 23%.

Upon relating this percentage to the lymphocyte density distribution profile the relative numbers of lymphocytes forming medium avidity EA- rosettes is shown (Fig. 5.9). Healthy subjects have most of their lymphocytes with medium avidity Fc receptors in fraction G whereas R.A. patients have relatively more in the lower density fractions D, E and F. However, R.A. patients (23.5%) do not form more medium avidity EA- rosettes than healthy subjects (29.1%).

The distribution of lymphocytes forming high avidity EA- rosettes is shown in Fig. 5.10 and Fig. 5.11. More EA- rosettes were formed in the denser lymphocyte fractions. R.A. patients formed more EA- rosettes than healthy subjects in fractions C, D, E, F and G, of which C and E are significant ($P < 0.05$). The trend shown here is different to that shown for medium avidity EA- rosettes. Relating these low rosette numbers to the cell numbers or fractions serves no useful purpose as the numbers are

Fig. 5.7.

- 5.8. Fc receptor bearing lymphocytes in Ficoll
 5.9. density gradient fractions. Comparison of
R.A. and healthy subjects.

30 ml of venous blood was taken and the lymphocytes prepared (Chapter 2.2.b.ii). Monocytes were depleted (Chapter 3.3.c.) and the lymphocytes separated by density in a Ficoll gradient (Chapter 3.3.d.). The interfaces were harvested and the lymphocytes adjusted to $2 \times 10^6 \text{ ml}^{-1}$. EA- rosettes were determined using the 1/800 titre of rabbit IgG anti ox red cell antisera (Chapter 5.2.b.ii). The results (Fig. 5.7) are expressed as the % EA- rosette forming cells in each fraction and each point represents the lymphocytes from one patient in each fraction.

The mean result (I 1 standard deviation) for each fraction is shown in Fig. 5.8. The only significant difference between R.A. and healthy subjects is in Fraction G ($p < 0.005$).

The EA- rosetting lymphocytes of each fraction are presented with the mean total cell proportion in each fraction (Fig. 5.9). This figure was obtained by multiplying the % incidence of EA- rosettes by the proportion of total lymphocytes in each fraction. E. G., in Fraction G for healthy subjects the EA- rosetting cells are 16 of the 67% (approx $\frac{1}{4}$) of the total lymphocytes in the gradient.


Fig. 5.7.


Fig. 5.8.


Fig. 5.9.

Fig. 5.10

High avidity Fc receptor-bearing lymphocytes
in Ficoll density gradient fractions. Comparison
of healthy subjects and R.A.

30 ml of venous blood was taken and the lymphocytes prepared (Chapter 2.2.b ii). Monocytes were depleted (Chapter 3.3.c.) and the lymphocytes separated by density in a Ficoll gradient (Chapter 3.3.d.). The interfaces were harvested and adjusted to $2 \times 10^6 \text{ ml}^{-1}$. EA rosettes were determined using the 1/3200 titre of rabbit IgG anti-ox red cell antisera (Chapter 5.2.b.ii). The results (Fig. 5.10) are expressed as the % EA- rosette forming cells in each fraction and each point represents the lymphocyte from one patients in each fraction.

The mean result (± 1 standard deviation) for each fraction is shown in Fig. 5.11. Statistically significant differences between healthy subjects and R.A. are shown only in fractions C and D ($p < 0.05$).

Fig. 5.10.





Fig. 5.11.

so low. Medium avidity rosettes also detected the high avidity ¹⁸¹
Fc receptor bearing population and hence to show the medium
avidity EA- rosette formation the latter set of results may
be subtracted from the former (Fig.5.12). This gives a
distribution pattern in which decreased proportions of rosettes
were shown in the denser fractions, particularly in R.A.

iii) SURFACE IMMUNOGLOBULIN

The percent of surface Ig bearing cells in the
Ficoll gradient fraction is shown in Fig. 5.13 and Fig. 5.14.

No statistically different results were recorded for R.A. patients
and healthy subjects. The results show a decrease in percentage
SIg bearing cells in the denser fractions. The dense fraction
(G) had 22% SIg positive cells whereas fraction B had 43% with a
more or less linear fall in numbers between these fractions.

Due to the similarities of these results for R.A. patients
and healthy subjects, relating SIg positive percentages to lymphocyte
density distribution merely becomes a function of the latter (Fig.
5.15).

5.4 DISCUSSION

In this study lymphocytes of varying densities were shown to
include different proportions of surface markers. Compared with
the high density cells of fraction G, the low density cells were
depleted of T cells, and those cells with high avidity Fc-
receptors. Moreover the less dense cell population was enriched
in B cells and those cells with medium avidity Fc- receptors.

Both E- rosette depletion and direct E-rosette studies on
gradient fractions suggested that low density cells were pre-
dominantly non-T cell. E- rosette depletion was used for
preliminary studies, but discarded due to the loss of lymphocytes

Fig. 5.12Medium avidity EA- rosette formation

The mean percentages of high avidity EA- rosette forming cells in Ficoll gradient fractions (Fig. 5.11) have been subtracted from the medium + high avidity EA- rosette percentages (Fig. 5.8). The result gives the calculated number of medium avidity EA- rosette forming cells in the different fractions.


Fig. 5.12.

Fig. 5.13
 5.14 Surface Ig bearing cells in Ficoll density
 5.15 gradient fractions. Comparison of R.A.
and healthy subjects.

30 ml of venous blood was taken and the lymphocytes prepared (Chapter 2.2.b.ii). The lymphocytes were labelled for SIg with 1/5 fluorescein isothiocyanate conjugated to rabbit anti human Ig (Chapter 2.2.h.). Monocytes were depleted by adherence (Chapter 3.3.) and the lymphocytes separated by density in a Ficoll gradient (Chapter 3.3.d.). The SIg positive cells were detected under a U.V. microscope and the results are expressed as percentage SIg positive cells in each gradient fraction (Fig. 5.13). Each point represents the lymphocytes from one patient in each fraction.

The mean result (± 1 standard deviation) is shown in Fig. 5.14. There was no statistical difference between R.A. and healthy subjects in any of the fractions. The surface Ig positive cells of each fraction are presented with the mean lymphocyte proportion in each fraction (Fig. 5.15). This figure was obtained by multiplying the % incidence of surface Ig positive cells by the proportion of total lymphocytes in each fraction, e.g., in Fraction G of healthy subjects the SIg positive cells are 15 of 67% (approx 1/4) of the total lymphocytes in the gradient.


Fig. 5.13.



Fig. 5.14.

Fig. 5.15.

upon using a second Ficoll-Paque separation. This favours the recovery of the lower density lymphocytes, as shown in Chapter 3.

The conditions used for the determination of E- rosettes satisfied the criteria suggested by the WHO report (1974). Divalent cations (Ca^{2+} and Mg^{2+}) were used in the medium, which included 10% human serum. Higher numbers for E- rosette percentages were only found by preincubating lymphocytes or SRBC with neuraminidase (unpublished). This is believed to stabilise the rosettes. The method employed in this study would also detect "active E rosettes" (Wybran and Fudenberg, 1973).

The distribution of E- rosettes in the gradient fractions suggests that the activated lymphocytes, both in R.A. and at low levels in healthy subjects, are predominantly non T cells. Another possibility is that T cells have a narrow high density range compared with B cells, if this were the case a large number would be lost during the initial isopycnic isolation of lymphocytes from whole blood. Even if T cells were denser than B cells this would not explain the increased numbers of low density cells in R.A. as the B cell numbers in R.A. peripheral blood are considered normal (Brenner, Scheinberg and Cathcart, 1975; Froland, Natvig and Wisloff, 1975).

The reduced numbers of T cells in the low density fractions could be explained by T cells losing their receptors for SRBC upon activation. Svedmyr et al (1974) found a high proportion of blasts in mixed leucocyte cultures lacking any surface marker attributable to B or T blasts. The authors found such null blasts were produced from almost pure SRBC - binding lymphocytes and considered the null blast to be of T origin. However, Jondal (1974, b) found that PHA and Con A induced T blasts retained their receptors for SRBC. Moreover, stimulation of purified T cells with allogeneic lymphocytes produced no change in the detection of receptors for SRBC (Jondal,

1974 b, c). The results of Svedmyr et al (1976) may be contradictory to those of Jondal (1974) because the stimulating cells used by the former were from a lymphoblastoid cell line. Thus, E- rosette formation studies of Ficoll gradient fractions were believed to detect resting and blast transformed T lymphocytes. Quantitative antigen-induced T cell proliferation has not been used to study T marker expression due to the small number of responding lymphocytes in such an in vitro test system.

The density distribution analysis of peripheral blood lymphocytes depleted of EAC- rosetting cells suggests that the low density cells in fractions C and D did not express C3 receptors. This is not conclusive, not just because only one patient was studied, but because C3 receptor bearing lymphocytes do not entirely overlap with surface immunoglobulin-positive cells (Ross et al, 1973). Caution must be taken when using C3 receptors as B cell markers as this marker is absent on PWM- induced surface immunoglobulin blasts (Jondal, 1974, b). Hence, the enrichment of cell numbers in fractions C and D after EAC- rosette depletion could be due to the presence of either B cells or T cells, particularly as activated lymphocytes are found in these fractions. Lymphocytes bearing receptors for C3 were the denser lymphocytes of fractions F, G and H, suggesting that non-active B cells, or T cells (Ross et al, 1978), are C3 receptor-positive.

The increased incidence of surface Ig positive cells in the low density fractions, 35-47% compared with 22% in fraction G, implies an increased number of B cells in these fractions. This would be in agreement with the relatively low incidence of T cells in these fractions. The absolute numbers of surface Ig positive cells in the fractions illustrates the low number of dense B cells

in R.A., even though the total B cell numbers in fractions B, C and D are not greater in R.A. than in healthy subjects.

Are the results representative of the B cell numbers in the density fractions? As the lymphocytes were labelled prior to gradient separation it is unlikely that the density equilibrium attained could affect the labelling quality due to a function of Ficoll concentration. Incubation at 37°C prior to Ficoll gradient separation should have removed any passively bound IgG (Winchester, Hoffman and Kunkel, 1975), and hence SIg negative cells with receptors for IgG should not be detected. The distribution of Fc receptor bearing lymphocytes, both medium and high avidity does not accord with the detection of passively bound IgG by the usage of FITC-labelled anti human Ig.

B lymphocytes activated in vitro with PWM give rise to B blasts which are surface Ig-positive (Jondal, 1974, b, c). With the knowledge that C3 receptors are not found on some activated B cells it seems that surface Ig detection is a good assay for B cells. If this holds true for in vivo produced blasts it may be possible to get information about activation of the immune system under different clinical conditions. However, there are difficulties in classifying malignant and non-malignant lymphoblasts. Acute lymphatic leukaemia is an example in which the present markers used for T and B cells are unable to classify the proliferating population (Seligman et al, 1974). Another example of cells lacking all markers seems to be that of lymphoblasts induced in vitro by stimulation with lymphoblastoid cell lines (Svedmyr et al, 1974).

Using SRBC receptors as a criteria for T cells, and surface

immunoglobulin as a B cell criterion, seems to allow classification of almost all lymphocytes and lymphoblasts with no overlapping of these markers (Jondal, 1974, b). Recently, Ross et al (1978) have suggested that the possession of Ia determinants is a better B cell marker than the presence of surface Ig as it detects the Ig-negative pre-B cells present in normal bone marrow. However, some monocytes have surface Ia and B cells ultimately lose detectable Ia determinants during the differentiation process into plasma cells. Plasma cells still possess detectable quantities of surface Ig, some expressing immunoglobulin on the surface of the same class that they secrete. It is unclear whether this is an integral membrane component or merely immunoglobulin trapped in the process of secretion. IgG- and IgA- secreting plasma cells fail to express surface IgD, but many retain surface IgM, particularly IgM secretory plasma cells (Halper et al, 1978). It is only in the terminal stages of B cell development that the surface Ig pattern changes and if such cells are of a low density the actual percentage B cells shown in the low density fractions of this study would be higher than estimated. Conversely, if dense cells, the B cell numbers shown for fraction G would be an underestimate. Plasma cells were seen predominantly in fractions C, D and E in this study (Chapter 3).

The major trend found for Fc receptor expression in the different gradient fractions was the distribution of lymphocytes with high avidity receptors. High avidity receptors were found mainly on denser cells. In R.A. increased numbers of these cells were found in fractions C, D, E, F and G. The differences in patterns for medium and high avidity Fc receptor bearing lymphocytes suggests that these may be detecting different lymphocyte populations. Fc receptors

are found on monocytes, neutrophils, B cells, some T cells (Van Boxel and Rosenstreich, 1974), L cells (Horwitz and Garrett, 1977) and K cells (Perlmann and Holm, 1969). Sharpin and Wilson (1977 b) believe the increased avidity of the R.A. lymphocytes for EA binding is due to increased numbers of K cells in this disease. K cells are believed to be non-B cell and this could account for the distribution of the high avidity receptors in the Ficoll gradient. The distribution of the medium avidity receptor bearing cells in the gradient is similar to that of B cells, particularly as this assay will also detect the high avidity population. Hence the high avidity receptor population are probably non-B and may be described as L cells, the cytotoxic potential uncharacterized. The high incidence of medium avidity Fc receptor bearing lymphocytes in fraction G in health may be related to the low number of high avidity cells in this fraction in comparison with R.A.

As T cells were shown predominantly in fraction G there is the possibility that these possess avid Fc receptors. The possession of Fc receptors by T cells is well documented (Yoshida and Andersson, 1974; Miyama et al, 1978) but no data on their binding avidity for IgG has been presented to date. The Fc receptors on T lymphocytes have been described as being limited to activated cells which would not normally be found in fraction G. However small effector cells, resulting from proliferation of a stimulated subpopulation, such as suppressor or cytotoxic T lymphocytes, could be found in this fraction. Small effector lymphocytes have been demonstrated with Fc receptors.

The interpretation of the Fc receptor data is open to more

speculation than for the T and B cell markers, mainly due to the overlap of cell populations resulting from the use of this cell marker. Certainly the lymphocytes in the low density gradient fractions in R.A. are more likely to be B cell than T cell, which composed the major part of the denser population.

CHAPTER SIX

DISCUSSION

6.1 LYMPHOCYTE ACTIVATION IN R.A.

This study describes an attempt to separate circulating lymphocytes by density in health and R.A. and to characterize these cells by surface marker studies. The density separation method is discussed first and the density distribution of R.A. lymphocytes will be related to spontaneous proliferation *in vivo*. The nature of the different lymphoid populations, as defined by density, is discussed next, with a view to elucidating their role in the pathogenesis of R.A.

In this study, the separation of peripheral blood lymphocytes by density has been successfully achieved in R.A. The discontinuous Ficoll gradient developed for this study has allowed the separation and recovery of lymphocytes of particular densities at high purity as shown by the rebanding of a single peak within a gradient. The lower density cells contain those which are proliferating, as shown by cytological examination and the density distribution of mitogen-stimulated lymphocytes. The lymphocyte density profiles obtained for healthy subjects were shown to be relatively constant and the method repeatable.

The low density fractions also included lymphocytes which morphologically were not activated. Steinman et al, (1978) showed that low density is acquired by lymphocytes soon after activation (18 hours) *in vitro* with mitogen. These small low density lymphocytes may be either recently activated or 'trapped' by non-specific association with larger cells. However, the rebanding of a peak of cells indicates that the latter is unlikely. To check the former possibility small low density lymphocytes could be prepared and

cultured in vitro and further checks on any density change or morphology performed.

To conclude that low density lymphocytes are activated requires that no other processes, in vivo or in vitro, affect lymphocyte density. Variations of osmolality and pH are known to change lymphocyte density (Shortman, 1968). Ficoll does not cause any change in toxicity and allows the use of a physiological pH (Gorczynski, Miller and Phillips, 1970) and hence should give a good estimate of the actual in vivo density of a cell. The effect of other factors which may affect lymphocyte density have not been reported.

The innumerable factors which a lymphocyte counters in the body may affect density without causing activation. All density distribution data relating to in vivo events must take this proviso into account, particularly when comparing patient sub-groups.

In any study in which peripheral blood is sampled it must be remembered that this sample represents one point in time. Activated lymphocytes are programmed to a set of functions and during the course of their development the nature of these cells will change. Steinman et al (1978) has demonstrated that recently activated cells acquire a low density, but has not commented on their density changes during development to blasts and to effector cells and/or memory cells. Plasma cells have a lower density than resting lymphocytes (Haskill, Legge and Shortman, 1969) but this has not been quantified with respect to other lymphoid cells. Dense lymphocytes will include small resting cells and recent progeny from activation. The lowest density cell will be the largest cell, the blast, due to its high cytoplasm/nuclear ratio. Plasma cells will have a density intermediate between blasts and small lymphocytes and hence will be found in the

same fractions as recently activated lymphocytes. It thus cannot be accurately predicted from a single sample whether an intermediate density cell is getting larger or smaller or is recently activated or maturing. Unfortunately there is no simple answer to this problem since serial assays of the different functions has not been attempted here but it is known that activated less dense cells do not survive well in culture (Moore and Hall, 1970).

The increased number of circulating immunoblasts found in R. A. is in general agreement with the findings of other authors (Bacon, Crowther and Sewell, 1975; Delbarre, Le Go and Kahan, 1975; Lalla et al, 1973). However, the large size reported by these authors was not confirmed. Certainly, cells up to a diameter of 30 μ m, as recorded by Delbarre et al (1975), were not observed. This is probably because of technical details. The numbers of immunoblasts, as recorded in this study, were related to disease activity. Raised numbers of immunoblasts were found particularly, if not exclusively, in patients with clinically active R.A. The presence of increased numbers of immunoblasts in the peripheral blood has been suggested as a useful indicator of increased intrinsic disease activity (Bacon et al, 1975). By comparison, the healthy subjects had low numbers of immunoblasts. This agrees with the findings of a number of authors (Bond et al, 1959; Crowther, Fairley and Sewell, 1969a; Bacon et al, 1975; Delbarre et al, 1975). The percentage of immunoblasts is often increased in pathological situations where inflammatory and immune reactions occur together, such as acute viral infections and connective tissue diseases. In acute viral infections this increase is transient, going on for 10-15 days, while in connective tissue diseases the immunoblast level may be high for long periods and grossly reflect the level of clinical activity. Inflammation

alone does not sufficiently explain these results, since patients suffering from psoriatic arthritis, even if associated with acute polyarthritis, have normal immunoblast levels (Delbarre et al, 1975). Gout patients also have low immunoblast levels (Eghtedari et al, 1976).

Thus the occurrence of immunoblasts in the peripheral blood is assumed to be related more to immune reactions than to inflammation and this hypothesis is supported by a number of investigations. The immunoblasts resemble the large, rapidly dividing hyperbasophilic blasts which are released into the efferent lymph by antigenically stimulated lymph nodes. (Hall, and Morris, 1963, 1965; Delbarre et al, 1969; Sprent and Miller, 1972b). Circulating immunoblasts are capable of producing immunoglobulins (Birbeck and Hall, 1967); they synthesize DNA as they circulate in the blood (Bond et al, 1959; Carter, 1965). Crowther, Fairley and Sewell (1969b) showed raised levels of immunoblasts during infection and immunization.

Evidence for the role of lymphocytes in the pathogenesis of R.A. is strengthened by the thoracic duct drainage studies of Paulus et al (1977). Prolonged continuous removal of thoracic duct lymphocytes through a surgical fistula resulted in significant clinical improvement of disease activity in patients with severe R.A. Cessation of lymph drainage, as well as reinfusion of autologous lymphocytes, resulted in exacerbation of disease activity.

The increased number of large lymphocytes recorded in this study in R.A. patients is supported by Whaley et al (1971) and was also found in juvenile R.A. by Brungard, Moser and Athreya (1975). The inference is that the increase is due to lymphocyte activation in vivo. However, the size of large lymphocytes in this study was not so great as that recorded by other investigators (Bacon et al, 1975; Delbarre et al, 1975).

The reported physical susceptibility of large lymphocytes (Hall and Morris, 1965) may help to explain this discrepancy. As immunoblasts do not recirculate from blood to lymph (Hall et al, 1967) and hence only exist in the circulating blood for a matter of hours, they must be produced in relatively large amounts by comparison to the production of small lymphocytes which do recirculate and have a life span measured in months or years. As they do not recirculate the immunoblasts are either sequestered in the tissues, divide to produce small lymphocytes in the circulation or die. The large number of activated lymphocytes in the rheumatoid synovium could explain either the source or the fate of the circulating immunoblasts.

The increased number of low density lymphocytes found in R.A., particularly in clinically active patients, is consistent with the activation of lymphocytes in vivo in this disease. The increased number of immunoblasts was also increased in active R.A. although no study was undertaken to compare lymphocyte density profiles and immunoblast levels sequentially in the same patients. The density profiles of R.A. lymphocytes were strikingly different to healthy subjects, in contrast to the small differences in lymphocyte diameter. Possibly, as suggested by Steinman et al (1978) this is a very sensitive assay for lymphocyte activation.

Yu, Peter, et al (1973), in their studies of prolonged thoracic duct drainage of R.A. patients, showed that during drainage the number of peripheral blood low density lymphocytes increased as the disease activity regressed. However, this is almost certainly due to an increased bone marrow output of lymphocyte precursors to compensate for the loss of lymphocyte during drainage, confirming the idea that less dense cells are recently activated ones.

Increased numbers of low density lymphocytes were found in the peripheral blood of clinically active SLE patients, whereas patients with inactive disease had cell density distributions similar to healthy controls. (Glinski et al, 1976). These investigators also found increased spontaneous transformation in SLE patients but upon investigating spontaneous ^3H -thymidine incorporation found maximum incorporation in the denser gradient fractions. However, the lymphocytes had been cultured in vitro for 72 hours prior to the adding of the radiolabelled DNA precursor and most large lymphocytes do not survive well in such culture conditions.

Hence the low density lymphocytes would have disappeared from the cultures before spontaneous transformation was determined. Otherwise, one would expect the low density lymphocytes to have the highest spontaneous proliferation, as found in this study.

It is easy to envisage the majority of the low density cells as precursors of morphological immunoblasts; the large number relative to morphological immunoblasts representing the long time span of the G1 and G2 phases of the cell cycle which culminate in the short lived blast cell which rapidly divides to produce memory or effector cells. However, as stressed earlier, this was a single point in time rather than a sequential study.

Most of the patients in this study were taking analgesics or non-steroid anti-inflammatory drugs. The effects of these drugs on lymphocyte function in vivo are largely unknown, there is no recorded evidence of lymphocyte activation by these agents. However, there is always the possibility that they may alter lymphocyte density in vivo. The low number of immunoblasts recorded in other inflammatory arthritides, where the patients were receiving a range of drug therapies (Delbarre, et al, 1975) indicates that this drug-induced

activation is unlikely. A number of 'second-line' drugs are known to affect lymphocyte function in vitro (Renoux et al, 1976; Markham et al, 1978), but not by acting as antigens. In this study the R.A. patients receiving gold therapy had a decreased number of low density lymphocytes relative to the patients with active disease. Yet the patients receiving the gold were doing so for previously active disease not controlled by NSAIDs. There are a number of possible explanations for these results. The gold could be lymphocytotoxic and be removing the cells with the greatest turnover i.e, activated lymphocytes. Secondly, the gold could cause a remission of disease activity by some other means which itself caused a decrease in spontaneous lymphocyte activation.

The widespread effects of corticosteroids and cytostatic drugs make their mechanism difficult to study. A longitudinal study of patients receiving various forms of these therapies is needed to examine changes in the density distribution pattern. This would help increase our knowledge as to the role of these cells in the disease state and aid in the search for effector mechanisms of these drugs.

The studies on cell markers showed that the low density cells had low numbers of T cells, were enriched with B cells and had few cells which bound IgG avidly. As to whether all the low density cells were in various stages of activation can be only conjecture due to the problems of morphological comparison. This must be borne in mind when ascribing numbers to the surface markers present in each fraction.

As B cells retain surface Ig upon activation and T cells retain their receptors for SRBC the use of these markers for B and T cells respectively, is justified for analysis of all cell fractions. Upon examination of the cell marker data for healthy subjects the immediate impression is that T cells are denser than B cells. For

this to be so the reduction of fraction G in R.A. would require a drastic T lymphopenia, particularly in the active patients. This does not appear to happen in R.A. (Brenner et al, 1975; Froland, Natvig and Wisloff, 1975). Hence, in R.A. more B cells than T cells have a low density. The low density B cells in healthy controls are probably part of the 'background immune response' which results in the low numbers of immunoblasts. These have been shown in rats to be part of gut defence, probably against bacteria, by the ability of syngeneic blasts to home to the gut of antigen-free neonates and many are precursors of IgA secreting cells. (Hall and Smith, 1970).

The increase in B cells in R.A. is unlikely to be due to an increase in gut defence mechanisms. Patients receiving analgesics or NSAIDs often suffer from gastro-intestinal bleeding, but as these materials are not antigenic it is unlikely that a rise in the rate of B blasts migrating to the gut would arise.

Lymphocytes in the peripheral blood and synovial fluid of R.A. patients have been characterized by surface markers in a number of studies (Brenner et al, 1975; Froland, Natvig and Wisloff, 1975; Utsinger, 1975; Holborow, Sheldon and Papamichail, 1975; Van de Putte et al, 1976). Comparisons of these reports are difficult due to the different T and B cell assays employed. The methodology for T cell determinations used in this study was recommended by the WHO/IARC Workshop on human B and T cells (1974), but differences in results can easily occur in the different methods of resuspension of the cell pellet prior to staining the rosettes. In this study standardization of the resuspension was attained by using a rotary turntable. The variations in B lymphocyte percentages could be due to a number of variables. Incubation at 37°C for 30 minutes has been used to remove cytophilic antibody from lymphocytes with Fc receptors (Lobo, Westervelt and Horwitz, 1975). Immunoglobulin

capping, endocytosis and loss of surface marker occur at 37°C (Ross et al, 1978). A depression of synovial fluid B lymphocytes has been shown by EAC rosette formation (Mellbye et al, 1972). However, this work has not taken into account the loss of C3 receptors by B lymphocytes upon activation (Jondal, 1974b). The study of C3 receptors in low density fractions in R.A. was touched on in this study, showing their relative absence in one case. This would support Jondal's work, but needs further experimentation to be conclusive.

The distribution of cells with Fc receptors showed that cells with high avidity receptors were predominantly dense cells. The results for medium avidity receptors will also include the detection of high avidity receptors and the subtraction of the two sets of results would indicate increased medium avidity receptor bearing lymphocytes in the low density fractions. Antibody dependent lymphocyte toxicity is mediated by a sub-population of lymphocytes which express Fc receptors but are neither B nor T lymphocytes, as defined by integral surface Ig or SRBC receptors respectively. These K cells (Perlmann and Holm, 1969) have high avidity Fc receptors, by comparison with B cells (Wooley and Panayi, 1978). Hence, the increased number of high avidity Fc receptor bearing lymphocytes in fraction G may reflect K cell activity and the medium avidity Fc receptor bearing lymphocytes, found mainly in low density fractions, may be the B cells. This hypothesis is consistent with the B cell distribution as determined by integral surface IgG. Monocytes also have Fc receptors, but were excluded from the sample prior to density gradient separation. The problems of a single time point study are illustrated again here, since the surface markers expressed by lymphocytes are, in some instances, dependent upon the stage of the cell cycle and this

is reflected by the difference of Fc receptor expression of lymphocytes in the density gradient. If any of the low density T cells are activated cells, they are not possessors of high avidity Fc receptors. Hence the possession of high avidity Fc receptors is limited to small lymphocytes, be they B, T, L or K sub-populations.

Upon accumulating the results for the Fc receptor positive populations in the gradient fractions it may be seen that the healthy subjects show a higher total EA- rosette formation than R.A. patients. This is contradictory to the results for the whole lymphocyte population data and could be due to two reasons. Firstly, the lymphocytes lost during the density gradient separation in R.A. may be mainly the Fc receptor positive cells. This preferential loss in R.A. would suggest that more of the Fc receptor positive cells in R.A. were activated than in healthy subjects. This would be in agreement with the physical susceptibility found for low density cells in this study. The other alternative is that the expression of Fc receptors by lymphocytes is altered by the passage through the Ficoll gradient. There is no evidence of this from other authors. The expression of integral surface Ig and receptors for SRBC does not appear to be so affected.

6.2 SIGNIFICANCE OF LYMPHOCYTE ACTIVATION IN R.A.

Having established the presence of both increased percentages of immunoblasts and low density lymphocytes in R.A., both in this study and in relation to other authors findings, I now turn to their possible significance in R.A. The role of activated lymphocytes in R.A. is discussed, particularly in the propagation of rheumatoid disease. The activation of lymphocyte sub-populations is particularly

considered to highlight the roles of both cellular and humoral immunity²⁰⁵ in the disease pathogenesis.

The raised numbers of immunoblasts and low density cells in clinically active arthritis suggests the importance of activated lymphocytes in the disease state. Immunological findings are not often shown to be temporally related to the clinical state of R.A. and the big question is whether blasts are a primary cause of increased disease activity or a secondary phenomenon. The possibility that they are bone marrow stem cells produced to compensate for a leucopenia must always be considered (Barnes and Loutit, 1967). However, an increase of R.A. disease activity is not always reflected by an increased severity of leucopenia. Secondly, the increased number of plasma cells is unlikely to be a result of increased bone marrow output.

The main possibility for the presence of these cells is antigen activation of circulating small lymphocytes. Generally, the search for a specific antigen in rheumatoid synovial tissue has proved unsuccessful (Runge, 1976; Morgan, 1978), although some success has been claimed (Bacon et al, 1973). However, one cannot exclude the existence of a putative antigen and the evidence for an infectious aetiology of R.A. has been extensively reviewed by Dumonde (1976). The possibility of antigens being sequestered in other tissues must always be considered particularly in view of the apparent systemic nature of the disease, which this study must support.

Antigens are often sequestered in lymphoid tissue, where they are exposed to a maximal number of circulating lymphocytes, and if this is the case in R.A. it may explain the lack of success at finding a synovial antigen. However, the rheumatoid synovium may

easily be considered a lymphoid organ, particularly with regard to the dense, follicular lymphoid infiltration.

Antigen stimulation of rat lymph nodes results in the appearance of large numbers of immunoblasts in the efferent lymph (Birbeck and Hall, 1967). The high percentage of these blasts which have been shown to migrate to the gut probably do so because of the variety of antigenic stimuli there (Parrott and Ferguson, 1974), although antigen is not necessary for migration to the gut (Moore and Hall, 1972). This may be related to a change in receptors for GALT expressed by immunoblasts, as suggested by the change in surface markers of activated cells (Jondal, 1974b). A change in the endothelial receptors on activated lymphocytes could explain their inability to recirculate from blood to lymph. Extending this theory to the immunoblasts in R.A. could account for the lymphocyte rich accumulations in the rheumatoid synovial membrane.

The investigations of Hall have centred largely on B blasts, but Jacobsson and Blomgren (1973) have produced evidence for a loss of recirculating capacity of T cells after antigenic stimulation. A large number of the lymphocytes in the rheumatoid synovium are T cells (Bankhurst, Husby and Williams, 1976) and may be a result of selective migration of T blasts to this tissue (Loewi, Lance and Reynolds, 1975). The mechanism of T cell attraction to the chronically inflamed synovium may be viewed in the light of the work of Asherson and co-workers (for review see Asherson, 1974). He showed that T cells obtained from lymph nodes draining sites of immunization in rats, or re-injection into other rats, moved to sites of inflammation in a non-antigen specific manner. Earlier North (1973) had shown that activated T cells appeared in the spleen after immunization with *Listeria*

monocytogenes, and these were considered to be the cells responsible for the passive transfer of community to *Listeria*. These cells have similarly been shown to accumulate at sites of inflammation such as an oil induced peritoneal exudate. Asherson states that such cells may be produced by procedures which give rise to cell-mediated immunity such as allogenic skin grafts, the injection of Freund's complete adjuvant, and skin painting with contact sensitizing agents.

In recent work, McGregor and Logie (1974) have shown that rats infected with *L. monocytogenes* contain immunoblasts in the thoracic duct which have the property of homing to peritoneal exudates induced either by injection of the same or another organism such as *Salmonella typhii*. This is most readily explicable in terms of a non-specific attraction by the endothelial cells of capillaries of inflamed tissues causing adherence of blast cells followed by their passage through the capillary wall into the inflammatory site. The blast cells may develop locally into T lymphocytes and they are able to arm locally accumulated macrophages, thus rendering them protective (Koster, McGregor and MacKanness, 1971). It is, of course, an open question whether a similar mechanism occurs in the rheumatoid synovial membrane or whether the lymphoid as well as other events are of a more local nature in the synovium.

Recently, Dumonde, Kelly and Morley (1976) have discussed the rheumatoid synovium as an ectopic lymphoid organ and suggested that alteration of lymphoid microvascular endothelia may result in abnormal lymphocyte function in R.A. Envisaging the rheumatoid synovium as a lymphoid organ could explain the lymphocyte activation and accumulation in this tissue. However, the alteration in lymphocyte function could still be ascribed to sequestration of an antigen(s) within this lymphoid tissue.

There are, as yet, no data on the nature of the cell traffic between the systemic pool and the local accumulation. Such data may be obtainable by suitable labelling techniques in rheumatoid patients. The traffic of small and large lymphocytes could be studied by labelling the various lymphoid populations from a Ficoll gradient and monitoring their localisation in the syngeneic host.

The intense infiltration of the synovium by lymphocytes includes many aggregates or follicles of plasma cells, which have been shown to contain deposits of immunoglobulin (Fish et al, 1966). The local synthesis of large amounts of immunoglobulins, both IgM and IgG, by rheumatoid synovium has been demonstrated (Sliwinski and Zvaifler, 1970) and Natvig and Munthe (1975) have interpreted the role of the many of the plasma cells to be the production of an IgG rheumatoid factor which combines with similar IgG molecules ("self-associating IgG") within the cell. However, only 20-60% of the IgG plasma cells showed anti-gammaglobulin fixing activity. The role of the remaining 40-80% and the IgM plasma cells requires better understanding, particularly as Ig M is found in lower amounts than IgG in the rheumatoid synovium and is not identified with complement (Zvaifler, 1973). The activated B cells demonstrated in this study could be related to this anti-immunoglobulin synthesis, and hence may be involved in propagating the disease in the body in this manner. Such an explanation could result in the sudden onset of polyarthritis noted clinically in some cases, by the propagation of activated B cells from regional lymphoid tissues which are sequestered in synovial tissue in many joints and become RF plasma cells secreting altered IgG or antiglobulin (IgG or IgM).

The thoracic duct drainage studies of Paulus et al (1977) suggest that lymphocytes leave the synovial space via lymphatics and the thoracic duct, rather than by directly entering the blood. This was determined by following the distribution of reinfused ^{51}Cr -labelled, living autologous lymphocytes. The disappearance of radioactivity was slower from arthritic joints, which had preferentially taken up some of the lymphocytes, than from the liver or spleen, indicating a prolonged retention of some lymphocytes in the inflamed joint. Support for this hypothesis comes from Whitehouse et al (1969) who reported that the removal of thoracic duct lymphocytes from rats that had been injected with Freund's complete adjuvant tended to prevent the subsequent development of adjuvant arthritis. However, if the drained lymphocytes were infused into untreated syngeneic littermates the recipient rats developed adjuvant arthritis, confirming the suggestion that thoracic duct lymphocytes were mediators of the disease.

The activated lymphocytes found in the synovial membrane in R.A. may give rise to the activated lymphocytes demonstrated by microscopy by Traycoff, Pascual and Schumacher (1976) in the synovial fluid. Again their presence was restricted to rheumatoid arthritis, being largely absent in gout, pseudogout or infective arthritis. Sebok, Talerman and Wouters (1977) found increased lymphocyte activation in R.A. synovial fluid by comparison with osteoarthritis and gout by the measurement of increased lymphocyte intra-nuclear birefringence. The authors interpreted their findings as being due to activation in the synovial membrane. The nature of these activated cells has not been determined. Similarly, Eghtedari et al (in press) found immunoblasts in the synovial fluid in R.A. as a higher percentage of lymphoid cells than in peripheral blood and the authors interpreted their presence as being a result of activation within the joint.

Studies on the T and B cell percentages in the synovial fluid of R.A. have yielded conflicting results. Utsinger (1975) reported normal T and B cell percentages in R.A. synovial fluid and was supported by Brenner et al (1975). Winchester et al (1973) reported an increased percentage of E-rosetting cells in R.A. synovial fluid compared with peripheral blood which Van de Putte et al (1976) explained as a general feature of inflammatory exudates. The T cells in synovial fluid have been shown to have impaired blastogenic responses to PHA (Ivanyi, Lehner and Burrey, 1973; Hepburn, McDuffie and Ritts, 1976) which may be interpreted as a prior commitment of lymphocytes due to activation *in vivo*.

Analysis of the mitogen and antigen responses of peripheral blood lymphocytes in R.A. has again produced equivocal results. The majority of data suggest that the response of R.A. lymphocytes to PHA is identical to control lymphocytes (Georgescu et al, 1975; Silverman et al, 1976). However, diminished responses have been recorded (Lockshin et al, 1975) and considered to be due to the blocking of PHA receptors by a factor present in autologous serum (Rawson, 1975). Thus the activation of peripheral blood T lymphocytes *in vivo* is not suggested. Accordingly R.A. lymphocytes respond to and stimulate normally with healthy lymphocytes in the MLC reaction (Stastny, 1974).

The range of T lymphocyte found in this study is similar to that found by other authors; this is not difficult as percentages have been reported between 30% (Keith and Currey, 1973) and 92% (Winchester et al, 1973) in the peripheral blood of R.A. patients. T and B cell percentages are generally reported as being normal in R.A. (Brenner et al, 1975). This is reasonable since the markers

employed, surface Ig and SRBC receptors are retained by lymphocytes²¹¹ even upon activation. Thus, enumeration of T and B cell ratios in peripheral blood and synovial fluid shows no drastic changes in R.A. and mitogen studies of T cell activity suggest there may be some in vivo cell activation.

The B cell activation in R.A. intimated in this study is difficult to relate to previous studies but is not surprising in view of the hypergammaglobulinaemia and circulating immune complexes found in R.A. Such increases have also been noted in other diseases, indeed, serum rheumatoid factor has been detected in sub-acute bacterial endocarditis and syphilis, indicating the non-specific nature of the classification or IgM anti-IgG anti-globulin. The study of PWM responses of peripheral blood lymphocytes in R.A. has shown no differences compared to controls (Rawson, 1975; Clot, Mathieu, Sany, Sèrre, 1975; Silverman et al, 1976). However, PWM activates both human B and T lymphocytes independently (Janossy and Greaves, 1971) and thus gives no clear indication of B cell function. Antigen responses by B cells result in the activation of so few cells that comparisons between R.A. and controls are difficult.

The presence of rheumatoid factor secreting plasma cells in the circulation and non-synovial tissues has been demonstrated in certain R.A. patients by a haemolytic plaque-forming cell assay (Moore et al, 1978). It has been shown recently (Male, Roitt, and Hay, 1978) that the immune complexes present in the synovial fluid in R.A. consist almost entirely of gammaglobulins which establish the importance of the IgG as an auto-immunogen.

Agglutination tests for IgM rheumatoid factor are positive in 60-70% of patients with R.A. The large variation in titre

does not necessarily reflect disease activity accurately.

The appearance of rheumatoid factor can precede clinical signs of disease or occur subsequently (Maini, 1977). As previously discussed (chapter 1) rheumatoid factors have been implicated as an important component of immune complexes in the synovial fluid of R.A., and it has been suggested that both IgG and IgM classes of rheumatoid factor are involved in activation of complement enhanced phagocytosis and consequent increased production and release of lysosomal enzymes.

The activated B cells in peripheral blood are probably precursors of plasma cells, although a memory function for at least some seems likely. An increased number of plasma cells would account for some of the B cell hyperactivity (Barden et al, 1967; Bandilla et al, 1970). However, this may represent the non-specific stimulatory effect of chronic inflammation rather than specific hyperactivity of the B cell system. The disturbed control of B cell function in R.A. may be due to a malfunction of immune regulation by T cell sub-sets. Certainly a deficiency of suppressor T cells could explain this. Patients with systemic lupus erythematosus have been demonstrated with a quantitative defect in suppressor T cells (Fauci et al, 1978). This disease is also characterized by a B cell hyperactivity resulting in a humoral auto-immune state characterized by hypergammaglobulinaemia and auto-antibodies against a variety of substances, particularly nucleoproteins and lymphocytes. T suppressor dysfunction has also been detected in the NZB/NZW mouse model which is felt to be the rodent equivalent of SLE (Krakauer, Waldman and Strober, 1970). It is of interest that Moretta et al (1978) recently demonstrated that human T lymphocytes with IgG receptors exposed in vitro to immune complexes irreversibly lost this Fc receptor. T lympho-

cytes with Ig G receptors have been shown to be T suppressors cells while those with IgM receptors were shown to be T helper cells. (Moretta et al, 1976). The immune complexes found in the synovial fluid in R.A. may cause the irreversible loss of T suppressor cell Fc receptors and ultimate depletion of these cells from the circulation. It is quite possible that T cell modulation of B cell function could be affected in this way, both in the joint and the peripheral blood. This could help to explain the chronicity of the disease, particularly as IgM immune complexes are more readily phagocytosed by polymorphs (Hurd et al, 1970). Thus the seropositivity of the patient, as assayed by classical IgM rheumatoid factor would not necessarily reflect the chronicity of the disease, IgG rheumatoid factor possibly accounting for this by the alteration of immune regulation. However, as IgM complexes are phagocytosed more avidly than IgG complexes these will cause much of the lysosomal enzyme release. Thus, both types of plasma cell play a role in the disease pathogenesis. Increased number of circulating B blasts would be necessary for B cell hyperactivity in the blood and the production of IgG and IgM as antiglobulins and IgG as antigen in the circulation and joint.

Envisaging rheumatoid synovitis as an extravascular immune complex disease is not difficult (Zvaifler, 1974) and the question must be asked; do the immune complexes activate the lymphocytes? Hall (1978) has shown specific migration inhibition in R.A. by immune complexes and shown (Hall, in preparation) specific binding by rheumatoid lymphocytes of an altered IgG structure (Facb). The inference is that not only is the autoimmunogen IgG altered but so are the lymphocyte receptors. This may be due to the increased avidity of Fc receptors recorded for R.A. in this study and these cells are predominantly small lymphocytes. Hence the activated population

may result from IgG activation of small B cells with altered receptors which upon activation lose these high avidity receptors.

This study has shown increased numbers of activated lymphocytes in the peripheral blood in R.A. These were shown to be predominantly B cell, although some T cell activation was noted. In all of the mechanisms discussed activation of peripheral blood lymphocytes is likely or indeed an essential component. It is the immunoblasts which appear to have the ability of progogate an immune response most effectively. Also it has been shown that spreading of inflammation to non-inflamed sites may be mediated by these cells. These properties could be most important in the immunopathogenesis of R.A.

The presence of a putative antigen, in the joint or sequestered in lymphoid tissue, must remain a strong possibility. Doubtless the role of circulating lymphocytes is complex; almost certainly both T and B populations are involved and activation, in the joint and in the circulation, a contributing part. Once we have found out the function of the activated population we will have isolated a major part of the altered immune mechanisms in R.A. The increase in knowledge of immunogenetics in recent years may help us unravel some problems by finding the susceptible subjects. For now we must content ourselves with the knowledge that many of the findings in R.A.- lymphocyte infiltration immune complex formation, autoantibodies, and impaired immuno-regulation-one common to a number of chronic diseases where an etiologic agent has been identified. Based on present information I propose that it will be shown that rheumatoid arthritis is caused by an infectious agent (or agents) and limited to individuals with a particular genetic constitution.

REFERENCES

ABDOU, N.L., ALAVI, J.B. and ABDOU, N.I. (1976)

Human bone marrow lymphocytes: B and T cell precursors and subpopulations. *Blood*, 47 : 423.

ABRAHAMSEN, T.G., FROLAND, S.S., NATVIG, J.B. and PAHLE, J. (1975)

Elution and characterisation of lymphocytes from rheumatoid inflammatory tissue. *Scand. J. Immunol.* 4 : 823.

ADLER, F.L. (1964)

Competition of antigens. *Prog. Allergy*, 8: 41.

ADLER, W.M., PEAVY, D. and SMITH, R.T. (1970)

The effect of PHA, PPD, allogeneic cells and sheep erythrocytes on albumin gradient-fractionated mouse spleen cell populations. *Cell. Immunol.* 1 : 78.

AIUTE, F. and WIGZELL, H. (1973)

Function and distribution pattern of human T lymphocytes. I. Production of anti-T lymphocyte specific sera as estimated by cytotoxicity and diminution of function of lymphocytes. *Clin. Exp. Immunol.* 13 : 171

ALEXANDER, E.L. and WETZEL, B. (1975)

Human lymphocytes: Similarity of B and T cell surface morphology *Science*, 188 : 732.

ANDERSON, L.C., HAYRY, P. and KONTAINEN, S. (1977).

Size distribution of in-vitro-induced antigen-specific helper and suppressor cells. *Scand. J. Immunol.* 6 : 141.

ANDERSSON, L.C., NORDLING, S. and HAYRY, P. (1973)

Fractionation of mouse T and B lymphocytes by preparative cell electrophoresis. Efficiency of the method. *Cell Immunol.* 8: 235

ASHERSON, G.L. (1974)

The role of T cells in inflammation: activated T cells which move to sites of inflammation and suppressor T cells which depress contact sensitivity, in "Future trends in inflammation".

Eds. Velo, G.P., Willoughby, D.A. and Giroud, J.P. Piccin Medical Books, London.

ASHERSON, G.L., ALLWOOD, G.G. and MAYHEW, B. (1973)

Contact sensitivity in the mouse: XI. Movement of T blasts in the draining lymph nodes to sites of inflammation. *Immunology*, 25 : 485 .

ASHERSON, G.L. and BARNES, R.M.R. (1973)

Contact sensitivity in mouse. XII. The measurement of DNA synthesis in vivo to determine the anatomical location of immunological unresponsiveness to picryl chloride. *Immunology*. 25 : 495.

ASKONAS, B.A. and ROELANTS, G.E. (1974)

Macrophages bearing hapten-carrier molecules as foci inducers for T and B lymphocyte interaction. *Eur. J. Immunol.* 4: 1

ASTÉRGA, G.P. and WILLIAMS, R.C. (1969)

Altered reactivity in mixed lymphocyte culture of lymphocytes from patients with rheumatoid arthritis. *Arth. Rheum.* 12:547.

AUGUST, C.S., MERLER, E., LUCAS, D.D. and JANEWAY, C.A. (1970)

The response in vitro of human lymphocytes to phytohaemagglutinin and to antigens after fractionation on discontinuous density gradients. Cell Immunol. 1 : 603.

AVRAMEAS, S. AND LEDUC, E.H. (1970)

Detection of simultaneous antibody synthesis in plasma cells and specialised lymphocytes in rabbit lymph nodes. J. Exp. Med. 131 : 1137.

BACH, M.K. and BRASHLER, J.R. (1970)

Isolation of subpopulations of lymphocytic cells by the use of isotonically balanced solutions of Ficoll. Exptl. Cell Res. 61 : 387.

BACON, P. A. (1979)

Peripheral blood and synovial fluid immunoblasts in rheumatoid arthritis. Ann. Rheum. Dis (in press)

BACON, P.A., BLUESTONE, R., CRACCHIOLO, A. and GOLDBERG, L.S. (1973)

Cell-mediated immunity to synovial antigens in rheumatoid arthritis Lancet (i): 699.

BACON, P.A., SEWELL, R.L. and CROWTHER, D. (1975)

Reactive lymphoid cells (immunoblasts) in autoimmune and haematological disorders. Clin. Exp. Immunol. 19 : 201.

BANDILLA, K.K., PITTS, N.C. and McDUFFIE, F.C. (1970)

Immunoglobulin M deficiency in the immune response of patients with RA . Arth. Rheum. 13 : 214.

BANKHURST, A.D., HUSBY, C. and WILLIAMS, R.C. (1976)

Predominance of T cells in the lymphocytic infiltrates of synovial tissues in rheumatoid arthritis. Arth. Rheum. 19 : 555.

BARDEN, J., MULLINA, F. AND WALLER, M. (1967)

Immunoglobulin levels in rheumatoid arthritis: Comparison with rheumatoid factor titres, clinical stage and disease duration. Arth. Rheum. 10 : 228

BARLAND, P. (1973)

Possible relationships between an infectious agent and the immune reactions of rheumatoid synovitis. Am.J. Med. 54: 143.

BARNES, D.W.H. and LOUTIT, J.F. (1967)

Haemopoietic stem cells in peripheral blood. Lancet (i):1138.

BAUR, P.S., THURMAN, G.B. and GOLDSTEIN, A.L. (1975)

Reappraisal of lymphocyte classification by means of surface morphology. J. Immunol. 115 : 1375

BERGSTROM, K., FRANKSSON, C., MATTELL, G. et al (1975)

Drainage of thoracic duct lymph in twelve patients with myasthenia gravis. Eur. Neurol. 13 : 19.

BHAN, A.K. and ROY, S. (1971)

Synovial giant cells in rheumatoid arthritis and other joint diseases

Ann. Rheum. Dis. 30: 204

BIRBECK, M.S. and HALL, J.G. (1967)

Transformation in vivo of basophilic lymph cells into plasma cells

Nature, 214 : 183.

BITTER, T. (1971)

A clinical investigation of the pathogenesis of autoimmunity. In

"Rheumatoid arthritis: Pathogenic Mechanisms and Consequences in therapeutics". Eds. Muller, W., Harwerth, H.G. and Fehr, K.

Academic Press. New York.

BLAND, J.H. and PHILLIPS, C.A. (1972)

Etiology and pathogenetics of rheumatoid arthritis and related multisystem diseases. Semin. Arth. Rheum. 1: 339.

BLOCH-SCHTACHER, N., HIRSCHHORN, K. and UHR, J.W. (1968)

The response of lymphocytes from non-immunized humans to antigen antibody complexes. Clin. Exp. Immunol. 3: 889.

BLOMGREN, H. (1975)

Role of B cells in the expression of the PPD response of human lymphocytes in vitro. Scand. J. Immunol. 4 : 499.

BOGGS, D.R. (1967)

Kinetics of neutrophil leucocytes in health and disease.

Semin. Haematol. 4 : 359.

BOND, V.P., FLIEDNER, T.M., CRONKITE, E.P. and ANDREWS, G. (1961)

Deoxyribonucleic acid synthesizing cells in the blood of man and dog exposed to total body radiation. J. Lab. Clin. Med. 57 : 711

BOND V.P., CRONKITE, E.P., RUBINI, J.R., BRECHER, G. and SCHORK, P.K. (1959)

Proliferative potentials of bone marrow and blood cells studied by in vitro uptake of ^3H -thymidine. Acta Haemat 21: 1.

BONOMINI, V., MIOLI, V., ALBERTAZZI, A. AND VANGELISTA, A. (1970)

Immunosuppressive drugs and lymphatic depletion by thoracic duct fistula in adult progressive glomerulonephritis. Nephron 7: 389

BOYCE, E.A., OLD, L.J. and CHOURLINKOV, I. (1964)

from "The lymphocyte response to activators" by Waite, W.I. and Hirschhorn In: Handbook of Experimental Immunology, Vol.2. Cellular Immunology, ed. D.M. Weir. Blackwell Scientific Publications, Oxford, 1973.

BRAIN, P. and MARSTON, R.H. (1973)

Rosette formation by human B and T lymphocytes. Europ. J. Immunol. 3: 6

BRAUNSTEIN, J.D., GOOD, R.A., HANSEN, J.A., SHARPLESS, T.K. and MEHAMED, M.R. (1976)

Quantitation of lymphocyte response to antigen by flow cytofluorimetry J. Histochem. Cytochem. 24 : 378

BRENDEL, W., RING, J. and SEIFERT, J. (1975)

Immunosuppressive treatment of multiple sclerosis with ALG and/or thoracic duct drainage. *Neurology* 25 : 490.

BRENNER, A.I., SCHEINBERG, M.A. and CATHCART, E.S. (1975)

Surface characteristics of synovial fluid and peripheral blood lymphocytes in inflammatory arthritis. *Arth. Rheum.* 18 : 297.

BREWER, D.B. (1972)

Activities of the neutrophil polymorph. *B.M.J.* 11 : 396.

BRITTON, M.C. and SCHUR, P.H. (1971)

The complement system in rheumatoid synovitis. II Intracytoplasmic inclusions of immunoglobulins and complement. *Arth. Rheum.* 14 : 87.

BRUBAKER, L.M. and EVANS, W.H. (1968)

Separation of granulocytes, monocytes, lymphocytes, erythrocytes and platelets from human blood and relative tagging with diisopropyl fluorophosphate (DFP). *J. Lab. Clin. Med.* 72 : 860

BRUNGARD, K., MOSER, G. and ATHREYA, B.M. (1975)

Size distribution of lymphocytes in juvenile rheumatoid arthritis. *J. Rheumatol* 2 : 359

CANTOR, H., SHEN, F. W., and BOYSE, E. A. (1976).

Separation of helper T cells from suppressor T cells expressing different Ly components. II. Activation by antigen: after immunization, antigen specific suppressor and helper activities are mediated by distinct T-cell sub-classes. J. Exp. Med.

143: 1391

CARTER, R. L. (1965).

The mitotic activity of circulating atypical mononuclear cells in infectious mononucleosis. Blood 26: 579

CARTWRIGHT, G. E., ATHENS, J. W., and WINTROBE, M. M. (1964).

The kinetics of granulopoiesis in normal man. Blood 24: 780

CASSIDY, M., YEE, C. and COSTA, J. (1976).

Automated analysis of antigen-stimulated lymphocytes. J. Histochem. and Cytochem. 24 : 373

CATOVSKY, D., GALLETTTO, J., OKOS, A., MILIANI, E. and GATTON, D. A. G. (1974).

Cytochemical profile of B and T leukaemic lymphocytes with special reference to acute lymphoblastic leukaemia. J. Clin. Path. 27: 767

CEROTTINI, J. C., NORDIN, A. A. and BRUNNER, K. T. (1970).

Specific in vitro cytotoxicity of thymus-derived lymphocytes sensitized to alloantigens. Nature 228: 1308

CHALMERS, D. G., COOPER, E. H., COULSON, A. S., INMAN, D. R. and

TOPPING, N. E., (1966).

Immunologically stimulated human peripheral blood lymphocytes in vitro. Int. Arch. Allergy 30: 83

CHAYEN, J. and BITENSKY, L. (1971).

Lysosomal enzymes and inflammation with particular reference to rheumatoid disease. *Ann. Rheum. Dis.* 30: 522

CLEMENTS, P. J., YU, D. T. Y., LEVY, J., PAULUS, H. E. and BARNETT, E. V. (1974).

Effects of cyclophosphamide on B and T lymphocytes in rheumatoid arthritis. *Arth. Rheum.* 17: 347

CLOT, J., MATHIEU, O., SANY, J. and SERRE, H. (1975).

Mitogens of B and T cells from peripheral blood in rheumatoid arthritis. *Rheum. Annual Rev.* 6: 199

COLE, P. (1977).

Respiratory disease. In: *Immunology in Medicine*. Eds. Holborow, E. J., and Reeves, W. G. Academic Press. London.

COOKE, T. D., and JASIN, H. E. (1972).

The pathogenesis of chronic inflammation in experimental antigen-induced arthritis. I. The role of antigen on the local immune response. *Arth. Rheum.* 15: 327

COOPER, H. L. (1968).

Ribonucleic acid metabolism in lymphocytes stimulated by phytohaemagglutinin. *J. Biol. Chem.* 243: 34

COULSON, A. S. (1969).

Recognition pathways in lymphocytes. *J. Theoret. Biol.* 25: 127

CRADDOCK, C. G. (1972).

Granulocyte kinetics. In: *Haematology*. Eds. Williams, W. J., Beutler, E., Erolow, A. J., and Rundles, R. W., McGraw-Hill. New York.

CRISWELL, B.S., RICH, R. R., DARDANO, J. and KIMZEY, S. L. (1975).

Scanning E.M of normal and mitogen-stimulated mouse lymphoid cells.

Cell Immunol. 19: 336

CROWTHER, D., HAMILTON-FAIRLEY, G. and SEWELL, R. L. (1969a).

Significance of the changes in the circulating lymphoid cells in

Hodgkins disease. B. M. J. 2: 473

CROWTHER, D., HAMILTON-FAIRLEY, G. and SEWELL, R. L. (1969b).

Lymphoid cellular responses in the blood after immunization in

Man. J. Exp. Med. 129: 849

CUDKOWICZ, G., BENNETT, M. and SHEARER, G. M. (1964).

Pluripotent stem cell function of the mouse marrow lymphocyte.

Science 144: 866

CUNNINGHAM, A. J., SMITH, J. B., and MERCER, E. H. (1966).

Antibody formation by single cells from lymph nodes and efferent

lymph of sheep. J. Exp. Med. 124: 701

DACIE, J. V., and LEWES, S. M. (1963).

Practical haematology. J & A Churchill. London.

DAMESHEK, W. (1963).

'Immunoblasts' and 'Immunocytes' - an attempt at a functional

nomenclature. Blood 21: 243

DAVID, J. R., AL-ASKARI, S., LAURENCE, H. S., and THOMAS, L. (1964).

Relayed hypersensitivity in vitro. I. The specificity of inhibition

of cell migration by antigens. J. Immun. 93: 264

DAY, R. P. (1972).

Basophil leucocyte separation from human peripheral blood:

a technique for their isolation in high purity and high yield.

Clin. Allergy. 2: 205

DELBARRE, F., LE GO, A. and KAHAN, A. (1975).

Hyperbasophilic immunoblasts in circulating blood in chronic inflammatory rheumatic and collagen diseases. Ann. Rheum.

Dis. 34: 422

DELORME, E. J., HODGETT, J., HALL, J. G. and ALEXANDER, P. (1969).

The cellular immune response to primary sarcomata in rats. I. The significance of large basophilic cells in the thoracic duct lymph following antigenic stimulation. Proc. Roy. Soc. 174: 229

DENHAM, S., HALL, J. G., WOLF, A. and ALEXANDER, P. (1969).

The nature of the cytotoxic cells in lymph following primary antigenic challenge. Transplantation 7: 194

DENMAN, A. M. (1975).

The viral theory of connective tissue diseases: A review.

Med. Biol. 53: 61

DE VRIES, E., VAN LEUWEN, A. W. F. M., VAN DE PUTTE, L. B. A.

et al. (1977).

Atypical T cells in rheumatoid synovial membranes. Virchows Arch.

B. Cell Path. 24: 19

DIAZ-JOUANEN, E., BANKHURST, A. D., and WILLIAMS, R. C. Jnr. (1976).

Antibody-mediated lymphocytotoxicity in rheumatoid arthritis and systemic lupus erythematosus. *Arth. Rheum.* 19: 133

DICKLER, H. B. (1976).

Lymphocyte receptors for immunoglobulin. *Adv. Immunol.* 24: 167

DITTRICH, H. (1962).

Physiology of neutrophils In: *The physiology and pathology of leucocytes.*

Eds: Braunstainer, H., and Zucker-Franklin, D. Grune and Stratton, New York.

DOUGLAS, S. D. (1971).

Human lymphocyte growth in vitro: morphologic biochemical and immunologic significance. *Int. Rev. Exptl. Path.* 10

DOWNNEY, H. and MCKINLAY, C. A. (1923).

Acute lymphadenosis compared with acute lymphatic leukaemia. *Arch. Int. Med.* 32: 82.

DUMONDE, D. C. (1976).

Ed. 'Infection and immunology in the rheumatic diseases'. Blackwell, London.

DUMONDE, D. C., KELLY, R. H. and MORLEY, J. (1976).

Lymphoid and microvascular dysfunction in experimental models of rheumatoid inflammation. In: *Infection and Immunology in the Rheumatic Diseases.* Ed. Dumonde, D. C. Blackwell, London.

DUMONDE, D. C., WOLSTENCROFT, R. A., PANAYI, G. S., MATTHEW, M., MORLEY, J. and HOWSON, W. T. (1969).

'Lymphokines' non antibody mediators of cellular immunity by lymphocyte activation. *Nature* 224: 38

DUNLAP, B., BACH, F.H. and BACH, M.L. (1978)

Cellsurface changes in alloantigen activated T lymphocytes.

Nature 271 : 253.

DUTTON, R.W. and EADY, J.D. (1964)

An in vitro system for the study of the mechanism of antigen stimulation in the secondary response. Immunology 7 : 40.

EDGREN, J., KLOCKARS, M. WEBER, T. ~~et~~ al (1976)

Extracorporeal irradiation of thoracic duct lymph as immunosuppressive treatment in rheumatoid arthritis. Scand.J.Rheumatol.5 : 108.

EGHTEDARI, A.A., BACON, P.A. and COLLINS, A.J. (1979)

Immunoblasts in synovial fluid and blood in the rheumatic diseases. Ann. rheum. Dis. (In Press).

EGHTEDARI, A.A., DAVIS, P. and BACON, P.A. (1970)

Immunological reactivity in ankylosing spondylitis - circulating immunoblasts, autoantibodies and immunoglobulins. Ann. rheum. Dis. 35 : 155.

EGHTEDARI, A.A., ESSELINCKX, W. and BACON, P.A. (1976)

Circulating immunoblasts in polymyalgia rheumatica. Ann. rheum.Dis. 35 : 158.

EIBL, M.A. and SITKO, C. (1975)

MIF production of lymphocytes from patients with rheumatoid arthritis with antigen-antibody complexes. Ann. rheum. Dis. 34 : 117.

ELLIOTT, B.E. and HASKILL, J.S. (1975)

Rosette-forming ability of thymus-derived lymphocytes in humoral and cell-mediated immunity II. Helper cell activity. J. Exp. Med. 141:600.

ELLIOTT, B.E., HASKILL, J.S. and AXELRAD, M.A. (1975)

Rosette-forming ability of thymus-derived lymphocytes in cell-mediated immunity I. Delayed hypersensitivity and in vitro cytotoxicity. J. Exp. Med. 141 : 584.

ELLIS, S.T., GOWANS, J.L. and HOWARD, J.C. (1969)

The origin of antibody forming cells from lymphocytes. Antibiotica et chemotherapia 15 : 40.

ELVES, M.W., ISRAELS, M.C.G. and COLLINGE, M. (1963)

The response of lymphocytes to antigen challenge in vitro. Lancet. i : 806.

ENGLISH, L.S., MORRIS, B. and ADMAS, E.P. (1976)

The synthesis of immunoglobulins by lymphoid cells in the sheep. Cell. Immunol. 27 : 274.

ENGLISH, L.S., MORRIS, B. and ADAMS, E.P. (1977)

The modulating effects of lymph on the synthesis of immunoglobulins by lymphoblasts. Cell. Immunol. 28 : 90.

EPSTEIN, L.B. and BRECHER, G. (1965)

DNA and RNA synthesis of circulating atypical lymphocytes in infectious mononucleosis. Blood. 25 : 197.

FAUCI, A.S., STEINBERG, A.D., HAYNES, B.F. and WHALEN, G. (1978)

Immunoregularity aberrations in systemic lupus erythematosus.

J. Immunol. 121 : 1473.

FELDBUSH, T.L. and STEWART, N. (1978)

Antigen modulation of the immune response V. Generation of large memory cells in antigen draining lymph nodes. Cell. Immunol.

37 : 336.

FELL, H.B. and DINGLE, J.T. (1963)

Studies on the mode of action of excess of vitamin A6. Lysosomal protease and the degradation of cartilage matrix. Biochem.J. 87:403.

FELL, H.B. and THOMAS, L. (1960)

Comparison of the effects of papain and vitamin A on cartilage II.

The effects on organ culture of embryonic skeletal tissue.

J. Expt. Med. 111 : 719.

FELSBURG, P.J. and EDELMAN, R. (1977)

The active E-rosette test: a sensitive in vitro correlate for human delayed type hyper sensitivity. J. Immunol. 118 : 62

FISCHER, D.B. and MUELLER, G.C. (1969a)

N-acetyl-D-galactosamine inhibits the phospholipid response by lymphocytes to phytohaemagglutinin. Nature 221 : 566.

FISH, A.J., MICHAEL, A.F., GEWUGG, H. and ROOD, R.A. (1966)

Immunopathologic changes in rheumatoid arthritis synovium.

Arth. Rheum. 2 : 267.

FLIEDNER, T.M., CRONKITE, E.P., BOND, V.P., RUBINI, J.R. and
ANDREWS, G. (1959)

The mitotic index of human bone marrow in healthy individuals and
irradiated human beings. Acta. haemat. 22 : 65.

FORD, W.L. (1974)

Lymphocyte migration and humoral responses. A Review. Progr.Allergy.

FORD, W.L. and GOWANS, J.L. (1969)

The traffic of lymphocytes. Semin. Haematol. 6 : 67.

FORSQREN, A., SVEDJELUND, A. and WIGZELL, H. (1976)

Lymphocyte stimulation by protein A of Staphylococcus aureus.

Eur. J. Immunol. 6 : 207.

FRADELIZI, D., CHARMOT, D., CROSIER, P.S., CONROY, A., MAWAS, C.E.
and SASPONTES, M. (1977)

Cellular origin of cytotoxic effectors and secondary educated
lymphocytes in human mixed leukocyte reaction. Cell. Immunol. 29: 6.

FRANCO, A.E. and SCHUR, P.H. (1971)

Hypocomplementemia in rheumatoid arthritis. Arth. Rheum. 14 : 231.

FROEBEL, K., STURROCK, R.D., DICK, W.C. and MACSWEEN, R.N.M. (1975)

Cell mediated immunity in the rheumatoid diseases I. Skin testing of
mitogenic responses in seronegative arthritides. Clin. Exp. Immunol.
22 : 446.

FROLAND, S.S., NATVIG, J.B. & WISLOFF, F.. (1975)

Lymphocyte Subpopulation in Rheumatoid Arthritis. Rheum.
Annual Rev. 6: 231.

FROLAND, S.S., WISLOFF, F. & MICHAELSEN, T.E.. (1974)

Human Lymphocytes with Receptors for IgG. A population
of Cells Distinct from T and B Lymphocytes. Int. Arch.
Allergy. 47 : 124.

FULWYLER, M.J.. (1965)

Electronic Separation of Cells by Volume. Science. 150: 910.

FULWYLER, M.J.. (1970)

Electronic Cell Sorting by Volume. In Wied, G.L. and Bohn, G.F.
eds., Automated Cell Identification and Cell Sorting. 97.
New York: Academic Press.

GABBIANI, G., TRENCH, P. and HOLBOROW, E.J.. (1975)

Increase of Contractile Proteins in Human Cancer Cells.
Lancet (ii). 796.

GARDNER, D.L. (1972)

The Pathology of rheumatoid arthritis. Edward Arnold, London.

GEORGESCU, C.M., GHEORGHIU, M and GOCIU, M (1975).

Investigations of delayed type hypersensitivity in rheumatoid arthritis. Scand. J. Rheumatol. Suppl 8 (Abstract) 14-

GERGELY, P., VANKY, F and KLEIN, E. (1976)

Spontaneous & PHA-induced rosetting of human blood, tonsil lymphocytes & MLC blasts with sheep, human & horse erythrocytes.

Clin. Exp. Immunol. 25: 23-

GESNER, B.M. and GINSBURG, V (1964).

Effect of glycosidases on the fate of transfused lymphocytes.

Proc. Nat. Acad. Sci., 52, 750.

GLINSKI, W., GERSHWIN, M.E., BUDMAN, D.R. and STEINBERG, A.O. (1976).

Study of lymphocyte subpopulations in normal humans and patients with SLE by fractionation of peripheral blood lymphocytes on a discontinuous Ficoll gradient.

Clin. Exp. Immunol. 26: 228.

GLYNN, L.E. (1969)

Aetiology of rheumatoid arthritis with regard to its chronicity.

Ann. Rheum. Dis. 28. Suppl. 3.

GORCZYNSKI, R.M., MILLER, R.G. & PHILLIPS, R.H. (1970)

Homogeneity of antibody-producing cells as analysed by their buoyant density in gradients of Ficoll. Immunology 19: 817-

GOWANS, J.L. (1962)

The fate of parental strain small lymphocytes in F1 hybrid rats.

Ann. N.Y. Acad. Sci.

GOWANS, J.L. (1966)

Lifespan, recirculation of transformation

Int. Rev. Exp. Pathol. 5: 1

GOWANS J.L. & KNIGHT, E.J. (1964).

The route of re-circulation of lymphocytes in the rat.

Proc. Roy. Soc. London Ser.B. 159: 257

GOWANS, J.L., MCGREGOR, D.D., COWEN, D.M. (1962) and FORD, C.E.

Initiation of immune responses by small lymphocytes.

Nature 196: 651

GOWANS, J.L. & UHR, J.W. (1966)

The carriage of immunological memory by small lymphocytes in the rat.

J. Exp. Med. 124: 101

GRODINA D.J., MILAS, L, HEWITT, R.R. and WITHERS, HR. (1973).

Buoyant density separation of human blood cells in renografin

gradient. Expt. Cell 81: 250

GREENBERG, A.H. (1973)

Fractionation of cytotoxic lymphoblasts on Ficoll gradients by velocity sedimentation.

Eur. J. Immunol. 3: 793

GRIFFITHS, M.M., SMITH, C.B., WARD, J.R. and KLAUBE, M.R. (1976).

Cytotoxic activity of rheumatoid & normal lymphocytes against allogeneic and autologous synovial cells in vitro.

J. Clin. Invest. 58: 613

GUNTHER, G., WARD J.L. and EDELMAN, G.M. (1974).

The kinetics of cellular commitment during stimulation of lymphocytes by lectins. J. Cell. Biol 62: 366

HALL, J.G. (1967)

Studies of the cells in the afferent and efferent lymph of lymph nodes draining the site of skin homografts.

J. Exp. Med. 125: 737

HALL, J.G. (1969)

Effector cells in Immunity.

Lancet (1): 25

HALL, J.G. (1970)

Demonstration of antibody on lymph - borne immunoblasts by bacterial adherence.

Rev Eur. Etud. Clin. Biol. 15: 783

HALL, J.G. (1974)

Observations on the migration and localization of lymphoid cells.

Progress in Immunology II Vol 3

(Eds L. Brent and J. Holborrow) North Holland Publishing Company.

HALL, J.G. and MORRIS, B (1963)

The lymph-borne cells of the immune response.

Quart. J. Exp. Physiol 48: 235

HALL, J.G. and MORRIS, B (1965)

The origin of the cells in the efferent lymph from a single lymph node.

J. exp. Med. 121: 901

HALL, J.G., MORRIS, B., MORENO, G.D. and BESSIS, M.D. (1967).

The ultrastructure and function of the cells in lymph following antigenic stimulation.

J. Exp. Med. 125: 91

HALL, J.G. PARRY, M.D. & SMITH, M.E. (1971)

Antibody production by lymph-borne immunoblasts following subcutaneous injection into Xenogeneic recipients.

Immunol. 20: 625

HALL, J.G. and SMITH, M.E. (1970)

Homing of lymph-borne immunoblasts to the gut.

Nature 226: 262

HALL, N.D. (1978)

Leucocyte migration inhibition with IgG- IgG complexes and rabbit IgG fragments in patients with rheumatoid arthritis.

Clin. exp. Immunol. 34: 219.

HALPER J., FU, S.M., WANG, C.Y., WINCHESTER, R. and KUNKEL, H.G. (1978)

Patterns in expression of human 'Ia-like' antigens during the terminal stages of B Cell development.

J. Immunol. 120: 1450

HALSTEAD, T.E. and HALL, J.G. (1972)

The homing lymph-borne immunoblasts to the small gut of neonatal rats.

Transplantation. 14: 339

HAMERMAN, D., BARLAND, P and JANIS, R (1969)

In 'The Biological basis of Medicine' Eds: Bittar, E.E. and Bittar, N.
Vol 3: 269.

HAMERMAN, D., JANIS, R, and SMITH, C.J. (1967)

Cartilage matrix depletion by rheumatoid synovial cells in tissue culture.

J. Exp. Med. 126: 1005.

HASKILL, J.S. (1967)

Density distribution analysis of antigen sensitive cells in the rat.

Nature 216: 1229.

HASKILL, J.S. (1969)

Density distribution analysis of antigen-sensitive cells in the rat.

J. Exp. Med 130: 877.

HASKILL, J.S., BYRT, P, and MARBROOK, J. (1970)

InVitro and in vivo studies of the immune response to sheep erythrocytes using partially purified cell populations.

J. Exp. Med. 131: 57.

HASKILL, J.S., LEGGE, D.G. and SHORTMAN, E. (1969)

Density distribution analysis of cells forming 19S haemocytic antibody in the rat.

J. Immunol. 102: 703.

HEDBERG, H (1963)

Studies on the depressed hemolytic complement activity of synovial fluid in adult rheumatoid arthritics.

Acta. Rheumatol. Scand. 9: 165.

HELLSTROM, U., ZEROMSKI, J and PERLMANN, P (1971)

Immunoglobulin light chain determinants on unstimulated and stimulated human blood lymphocytes, assayed by indirect immunofluorescence.

Immunol. 20: 1099.

HEPBURN, B., McDUFFIE, F, and RITTS, R.E. (1976).

Impaired blastogenic response of lymphocytes from synovial fluid and peripheral blood of patients with rheumatoid arthritis..

J. Rheumatol 3:(2): 118.

HEYN, R.M., TUBERGEN D.G. & ALTHOUSE, N.T. (1973).

Lymphocyte size distribution.

Amer. J. Dis. Child 125: 789.

HILAL, S.K., MOSSER, D.G., LOKEN, M.K. and JOHNSON, R.W. (1964)

A new technique for high resolution density gradient separation of bone marrow cells.

Ann. N.Y. Acad. Sci. 114: 661.

HIRSCHHORN, K., BACH, F., KOLODNY, R.L. FIRSCHHEIN, I.L. and HASH^{EM}, N

(1963). Immune response and mitosis of human peripheral blood lymphocytes in vitro.

Science 142: 1185.

HIRSCHHORN, R., BRITTINGER, G., HIRSCHHORN K and WEISSMAN, G (1968).

Studies on lysosomes: XII. Redistribution of acid hydrolases in human lymphocytes stimulated by photohaemagglutinin.

J. Cell. Biol. 37: 412.

HIRSCHHORN, K and HIRSCHHORN, R (1965)

Role of lysosomes in the lymphocyte response

Lancet i: 1046.

HIRSCHHORN, R, TROLL, W., BRITTINGER, G and WEISSMANN, G (1969).

Template activity of nuclei from stimulated lymphocytes.

Nature 222: 1247.

HOLBOROW, E.J., SHELDON, P.J. and PAPAMICHAEL, M (1975).

Studies on synovial fluid lymphocytes in rheumatoid arthritis.

Rheum. Annual Rev. 6: 215

HOLLANDER, N., GINSBURG, H and FELDMAN (1974).

In vitro generation of memory lymphocytes reactive to transplanation antigens.

J. Exp. Med. 140: 1057.

HOLLINGWORTH, J.W., SIEGEL, E.R. and CREASEY, W.A. (1967)

Granulocyte survival in synovial exudate of patients with rheumatoid arthritis and other inflammatory joint diseases.

Yale J. Biol. Med. 39: 289.

HOLT, P.J.L. (1971)

Ultrastructural changes in rheumatoid arthritis articular cartilage in rheumatoid arthritis. ed: Muller, W., Harwerth, H.G. and Fehr, K Academic Press, London & N. York.

HOPKINS, J. and HALL, J.G. (1976)

Selective entry of immunoblasts into gut from intestinal lymph.

Nature 259: 308.

HIRSCHHORN D.A. and GARRETT M.A. (1977)

Distinctive functional properties of human blood L lymphocytes.

A comparison with T lymphocytes, B lymphocytes and monocytes.

J. Immunol. 118: 1712.

HIRSCHHORN D.A. and LOBO, P.I. (1975).

Characterization of two populations of human lymphocytes bearing easily detectable surface immunoglobulin.

J. Clin. Invest. 56: 1464.

HORWITZ, D.A. STASTNY, P and ZIFF, M. (1970)

Circulating DNA synthesizing mononuclear leukocytes. I. Increased numbers of proliferating mononuclear leukocytes in inflammatory disease.

J. Lab. Clin. Med. 76: 391

HOWARD, J.G. HUNT, S.V. GOWANS J.L. (1972).

Identification of narrow derived and thymus derived small lymphocytes in the lymphoid tissue and thoracic duct lymph of normal rats. J. Ex. Med. 135: 200.

HOYERALL, H.M., FROLAND, S.S., AND WISLOFF, F. (1975).

Lymphocyte populations and cellular immune reactions in juvenile rheumatoid arthritis.

Scand.J. Immunol 4: 801.

HUBER, B., CANTOR, H., SHEN, F.W. and BOYLE, E.A. (1976).

Independent differentiative pathways of Ly 1 and Ly 23 subclasses of T cells.

J. Exp. Med. 144: 1128

HUNT, S.V., ELLIS S.T. and GOWANS, J.L. (1972)

The roles of lymphocytes in antibody formation IV.

Proc. Roy. Soc B 182: 211

HURD, E.R. SPALLUTO J, and ZIFF, M. (1970)

Formation of leukocyte inclusions in normal polymorphnuclear cells incubated with synovial fluid.

Arth.Rheum. 13: 724.

INMAN, D.R. and COOPER, E.H.. (1965)

The Relation of Ultrastructure to DNA Synthesis in Human

Leucocytes. I Atypical lymphocytes in Phytohaemagglutinin

Cultures and Injections Mononucleosis. Acta. Haemat. 33: 257.

ISHIKAWA, Y. and ZIFF, M.. (1976)

Electron Microscopic Observations of Immunoreactive Cells in the

Rheumatoid Synovial Membrane. Arthritis & Rheumatism 19: 1

ISTURIZI, M.A., De BRACCO, M.D.M., PIZZI, A.M. and MANNI, J.A..
(1976).

Antibody-dependent Cell-mediated Cytotoxicity in Rheumatoid
Arthritis. *Arth. Rheum.* 19: 725.

IVANYI, L., LEHNER, T. and BURRY, H.C.. (1973).

The Response of Synovial FLuid Lymphocytes to T and B
Stimulants in Vitro. *Immunology* 25: 905.

JACOBSSON, H. and BLOMGREN, H.. (1973)

Evidence for a Loss of Recirculating Capacity of T Cells
After Antigenic Stimulation. *Clin. Exp. Immunol.* 13: 439.

JANÓSSY, G. and GREAVES, M.F.. (1971)

Lymphocyte Activation I. Response of T and B Cells to
Phytoimitogens. *Clin. Exp. Immunol.* 9: 483.

JOHNSON, P.M., WATKINS, T. and HOLBORROW, E.J.. (1975)

Hypothesis: Antiglobulin Production to Altered IgG in
Rheumatoid Arthritis. *Lancet* (i) 611.

JONDAL, M.. (1974a)

Surface Markers on Human B & T Lymphocytes. III. A marker
for Lymphoid Adherence. *Scand. J. Immunol.* 3: 269.

JONDAL, M.. (1974b)

Surface Markers on Human B & T Lymphocytes. IV. Distribution
of Surface Markers on Resting & Blast-transformed Lymphocytes.
Scand. J. Immunol. 3: 739.

JONDAL, M. (1974c)

Surface markers on human B and T lymphocytes v. Characterization of the lymphoproliferative response to three different lectins and allogeneic lymphocytes by surface markers. Scand.J.Immunol.

3 : 749

JONDAL, M., HOLM, G. and WIGZELL, G. (1972)

Surface Markers as Human B and T. Lymphocytes. 1. A large Population of Lymphocytes forming Non-immune Rosettes with SRBC. J. Exp. Med. 136 : 207.

JULIUS, M.H. and HERZENBERG, L.A. (1974)

Isolation of antigen-binding cells from unprimed mice.

Demonstration of antibody forming cell precursor activity and correlation between precursor and secreted antibody.

Avidities. J. Exp. Med. 140 : 904

KEITH, H. I. and CURREY, H.L.F. (1973)

Rosette formation by peripheral blood lymphocytes in rheumatoid arthritis. Ann. Rheum. Dis. 32 : 202.

KING, R.A., MESSNER, R.P. and WILLIAMS, R.C. (1968)

Possible adjuvant effect of rheumatoid factors - an autologous immunostat. Arth. Rheum. 11 : 828.

KINOSHITA, Y., KIMURA, S., TAKESHITA, T., KIMURA, G.,

YUKIOKA, M. and MIROSAWA, S. (1970)

Isolation of heterogenous lymphocytes according to their cellular densities by multilayer centrifugation.

Exptl. Cell. Res. 59 : 299.

KNAPP, W., SCHUIT, H.R.E., BOLHUIS, R.L.H. and HIJAMS, W. (1974)

Surface immunoglobulins in chronic lymphatic leukaemia, macroglobulinaemia and myelomatosis. Clin. Exp. Immunol. 16 : 541.

KOBAYASHI I. and ZIFF, M. (1973)

Electron microscopic studies of lymphoid cells in the rheumatoid synovial membrane. Arth. Rheum. , 16 : 471.

KONTAINEN, S. and FELDMANN, M. (1975)

Conditions for inducing T helper cells in vitro. Scand. J. Immunol. 4 : 121.

KONTAINEN, S. and FELDMANN, M. (1976)

Suppressor cell induction in vitro. 1. Kinetics of induction of antigen specific suppressor cells. Eur. J. Immunol. 6 : 296.

KOSTER, F. and MCGREGOR, D.D. (1970)

Rat thoracic duct lymphocytes : Types that participate in inflammation. Science. 167 : 1137.

KOSTER , F.T., MCGREGOR, D.D. and MACKNESS, S.G.B. (1971)

The mediator of cellular immunity. II. Migration of immunologically committed lymphocytes into inflammatory exudates. J. Exp. Med. 133 : 400.

KRAKUER, R.S., WALDMAN, T.A. and STROBER, W. (1976)

Loss of suppressor T cells in adult NZB/NZW mice. J. Exp. Med. 144 : 66.

KRAMMER, P.H. (1978)

Alloantigen receptors on activated T cells in mice . 1. Binding of alloantigens and anti-idiotypic antibodies to the same receptors.

J. Exp. Med. 147 : 25.

KULKA, J.P. (1966)

Vascular derangement in rheumatoid arthritis. In: Modern Trends in Rheumatology. p. 49. Ed. Hill, A.G.S., Butterworth, London.

KVEIM, A. (1941)

En. ny og Spesifikk Kutan - reaksjon ved Boecks Sarcoid.

Nord. Med. 2 : 169.

LAWTON, A.R., SELF, K.S., ROYAL, S.A. and COOPER, M.D. (1972)

Ontogeny of B-lymphocytes in the human foetus. Clin. Immunol.

Immunopathol. 1 :84.

LEIF, R.C. and VINOGRAD, J. (1964)

The distribution of buoyant density of human erythrocytes in bovine albumin solutions. Proc. N.A.S. 51 : 520.

LEVIS, W.R. and ROBBINS, J.H. (1972)

Methods for obtaining purified lymphocytes, glass-adherent

mononuclear cells and a population containing both cell types from human peripheral blood. Blood. 40 : 77.

LEWIS, H., MITCHELL, J. and NOSSAL, G.J.V. (1969)

Subpopulations of rat and mouse thoracic duct small lymphocytes in the Salmonella flagellar antigen system. Immunology, 17 : 955.

LIN, P.S., COOPER, A.G. and WORTIS, H.H. (1973)

Scanning electron microscopy of human T-cell and B-cell rosettes.

N. Engl. J. Med. 289 : 548.

LIN, P.S., WALLACH, D.F.M. and TSAI, S. (1973)

Temperature-induced variations in the surface topology of cultured lymphocytes are revealed by scanning electron microscopy. Proc.

Nat. Acad. Sci. 70 : 2492.

LINDNER, J. (1971)

Glycosaminoglycan metabolism in rheumatoid arthritis in hypergic knee joint arthritis in the rabbit. In: Rheumatoid Arthritis, ed.

W.Miller,Harwerth,H.G. and Fehr,K.Academic Press,London and New York.

LING, N.R. (1968)

Lymphocyte stimulation. Amsterdam: North Holland Publishing Company.

LITT, M. (1964)

Studies in experimental eosinophilia. VI. Uptake of immune complexes by eosinophils. J. Cell. Biol. 23 : 335.

LOBO, P.I., WESTERVELT, F.B. and HORWITZ, D.A. (1975)

Identification of two populations of immunoglobulin bearing lymphocytes in man. J. Immunol. 114 : 116.

LOCKSHIN, M.D., EISENHAUER, A.C., KOHN, R. et al (1975)

Cell mediated immunity in rheumatic diseases. II. Mitogen responses in RA, SLE and other illnesses : correlation with T and B lymphocyte populations. Arth. Rheum. 18 : 245.

LOEWI, G., DORLING, J. and HOWARD, A. (1974)

Mononuclear cells from inflammatory joint effusions : Electron-microscopic appearances and immunoglobulin synthesis. J. Rheumatol. 1 , 34.

LOEWI, G., LANCE, E.M. and REYNOLDS, J. (1975)

Study of lymph cells from inflamed synovial membranes. Ann.Rheum.Dis. 34 : 524.

LOOS, J.A. and ROOS, D. (1974)

Ficoll - Isopaque gradients for the determination of density distributions of human blood lymphocytes and other reticulo-endothelial cells. Exp. Cell. Res. 86 : 333.

LOWRY, O.H., ROSEBROUGH, N.J., FARR, A.L. and RANDALL, R.J. (1951)
Protein measurement with the Folin phenol reagent. J. Biol. Chem.,
193 : 265.

LUCAS, D.O., SHOHEET, S.B. and MERLER, E. (1971)
Changes in phospholipid metabolism which occur as a consequence of
mitogenic stimulation of lymphocytes. J. Immunol. 106 : 768.

MAINI, R.N.. (1977)

Immunology of the Rheumatic Diseases. Current Topics in
Immunology, 7, Arnold, London.

MALE, D., ROITT, I.M. and HAY, F.C.. (1978)

Analysis of Complexes in Rheumatoid Arthritis. (abstract).
4th Eur. Immunol. Meeting. (Budapest).

MARCHESI, V.T. and GOWANS, J.L.. (1964)

The Migration of Lymphocytes Through the Endothelium of Venules
in Lymph Nodes: an electron microscope study. Proc. Roy. Soc.
B. 159, 283.

MARCUS, R.L. and TOWNES, A.S.. (1971)

The Occurrence of Cryoproteins in Synovial Fluid; the
Association of a Complement - Fixing Activity in Rheumatoid
Synovial Fluid with Cold - Precipitable Protein.

MARKHAM, R.B., STASHAK, P.W., PRESCOTT, B., AMSBAUGH, D.F. and
BAKER, P.J.. (1978).

Selective Sensitivity to Hydrocortisone of Regulatory Junctions
that Determine the Magnitude of the Antibody Response to Type
III Pneumococcal Polysaccharide. J. Immunol. 121: 829.

MATTHES, M.L., AX, W. and FISCHER, H. (1971)

Morphometriccinematographic Analysis of the Co-operation Between
Lymphocytes and Dendritic Macrophages. In "Cell Interactions
and Receptor Antibodies in Immune Responses". Eds. Makela, O.,
Cross, A. and Kosunen, T.U.. Academic Press, London.

MACGREGOR, D.D. and LOGIE, P.S.. (1974)

The Mediator of Cellular Immunity. VII Localization of
Sensitized Lymphocytes in Inflammatory Exudates. J. Exp. Med.
139: 1415.

McCLUSKEY, R.T., BENACERRAF, B. and McCLUSKEY, J.W.. (1963)

Studies of the Specificity of the Cellular Infiltrate in
Delayed Hypersensitivity Reactions. J. Immunol. 90: 466

McCOOL, D., MILLER, R.G., PAINTER, R.H. and BRUCE, W.R.. (1970)

Erythropoietin Sensitivity of Rat Bone Marrow Cells Separated
by Velocity Sedimentation. Cell. Tiss. Kinet. 3: 55

McWILLIAMS, M., PHILLIPS-QUAGLIATA, J.M. and LAMM, M.E.. (1977)

Mesenteric Lymph Node B Lymphoblasts which home to the Small
Intestine are Precommitted to IgA Synthesis. J. Exp. Med.
145: 866.

MELLBYE, O.J., MESSNER, R.P., DeBORD, J.R. and WILLIAMS, R.C..
(1972).

Immunoglobulin and Receptors for C3 on Lymphocytes from
Patients with Rheumatoid Arthritis. Arth. Rheum. 15: 371

MENARD, H.A., DION, J. and RICHARD, C.. (1977)

Antinuclear Antibody: Predictive of Lymphocyte Response
in Rheumatoid Arthritis. J. Rheumatol. 4: 21-26.

METCALF, W.K., LUDWIG, S., GOULD, R.N. and METCALF, N.F.. (1976)

The Lymphocyte Cytoplasmic Refractive Index in Rheumatoid
Arthritis. Wadley Med. Bull. 6: 109.

MICHELI, A. and BRON, J.. (1974)

Studies on Blood T and B Lymphocytes in Rheumatoid Arthritis.

Ann. Rheum. Dis. 33: 435.

MILLER, R.G. and PHILLIPS, R.A.. (1970)

Sedimentation Analysis of the Cells in Mice Required to Initiate
an In Vivo Immune Response to Sheep Erythrocytes. Proc. Soc.

Exp. Biol. (N.Y.) 135: 63.

MIYAMA, M., KURIBAYASHI, K., YODOI, J., TAKABAYASHI, A. and

MASUDA, T.. (1978)

Immunological Properties of Fc Receptor on Lymphocytes. I.

Function Differences Between Fc Receptor - Positive and

Negative Lymphocytes in Humoral Immune Responses. Cell. Immunol.

35: 253.

MINDERHOUD, J.M. and SMITH, J.K.. (1972)

Immunological Activity of Blood Lymphocyte Fractions. A Study
by the Macrophage Electrophoretic Mobility Method. Clin. Exp.

Immunol. 10: 571.

MOCHIZUKI, D., HEDRICK, S., WATSON, J. and KINGSBURY, D.T.. (1977)

The Interaction of Herpes Simplex Virus with Murine Lymphocytes.

I. Mitogenic Properties of Herpes Simplex Virus. J. Exp. Med.

146: 1500.

MOLLER, G.. (1969)

Induction of DNA Synthesis in Normal Human Lymphocyte Cultures by

Antigen - Antibody Complexes. Clin. Exp. Immunol. 4: 65.

MOORE, A.R. and HALL, J.G.. (1972)

Evidence for a Primary Association Between Immunoblasts and Small Gut. *Nature* 239: 161.

MOORE, A.R. and HALL, J.G.. (1973)

Nonspecificity of Thoracic Duct Immunoblasts into Intradermal Foci of Antigens. *Cell. Immun.* 8: 112.

MOORE, T.L., ROBBINS, D.L., ROSE, J.E. and VAUGHAN, J.H.. (1978)

Drugs Affecting the Release of Rheumatoid Factor in a Plaque-Forming Cell Assay. *Arth. Rheum.* 21: 114.

MORETTA, L., FERRARRINI, M., MINGARI, M.C., MORETTA, A. and WEBB, S.R.. (1976)

Subpopulations of Human T Cells Identified by Receptors for Immunoglobulins and Mitogen Responsiveness. *J. Immunol.* 117: 2171

MORETTA, L., MINGARI, M.C. and ROMANZI, C.A.. (1978)

Loss of Fc Receptors from Human T lymphocytes Exposed to IgG Immune Complexes. *Nature* 272: 618

MORETTA, L., WEBB, S.R., GROSSI, C.E., LYDYARD, P.M. and COOPER, M.D.. (1977)

Functional Analysis of Two Human T Cell Subpopulations: Help and Suppression of B-Cell Responses by T Cells Bearing Receptors for IgM. *J. Exp. Med.* 146: 184.

MORGAN, J.E.. (1978)

The Search for a Synovial Antigen in Rheumatoid Arthritis. Ph.D. Thesis. University of Bath.

MULLER, W. (1971)

Immunological Reactivity in Rheumatoid Arthritis. In Rheumatoid Arthritis: Pathogenetic Mechanisms and Consequences in Therapeutics. Eds. Muller, W., Harwerth, H.G. and Fehr, K.. Academic Press. New York.

MUNTHE, E. and NATVIG, J.B.. (1972)

Immunoglobulin Classes, Subclasses and Complexes of IgG
Rheumatoid Factor in Rheumatoid Plasma Cells. Clin. Exp. Immunol. 12: 55.

MUIRDEN, K.D. (1966)

Ferritin in Synovial Cells in Patients with Rheumatoid Arthritis. Ann. Rheum. Dis. 25: 377.

MURPHY, M.J., HAY, J.B., MORRIS, B and BESSIS, M.C.. (1972)

Ultrastructural Analysis of Antibody Synthesis in Cells from Lymph and Lymph nodes. Am. J. Path. 66: 25.

NAGY, Z., ELLIOTT, B.E., NABHOLZ, M., KRAMMER, P.H. and PERNIS, B.. (1976).

Specific Binding of Alloantigens to T Cells Activated in the Mixed Lymphocyte Reaction. J. Exp. Med. 143: 648

NASPITZ, C.K. and RICHTER, M.. (1968)

The Action of Phytohaemagglutinin in Vivo and in Vitro, a Review. Prog. Allergy 12: 1.

NATVIG, J.B. and MUNTHER, E.. (1975)

Self Associating IgG Rheumatoid Factor Represents a Major
Response of Plasma Cells in Rheumatoid Inflammatory Tissue.

Ann. N.Y. Acad. Sci. 256: 88

NORDLING, S., ANDERSSON, L.C. and HAYRY, P.. (1972)

Thymus-dependent and Thymus-independent Lymphocyte Separation:
Relation to Exposed Sialic Acid on Cell Surface.

Science. 178: 1001

NORTH, R.J.. (1973)

Cellular Mediators of Anti-Listeria Immunity as an Enlarged
Population of Short Lived Replicating T Cells. J. Exp. Med.

138: 342.

OSOBA, D.. (1970)

Some Physical and Radiobiological Properties of Immunologically
Reactive Mouse Spleen Cells. J. Exp. Med. 132: 368.

PAGE-THOMAS, D.P.. (1968)

The Role of Lysosomes in Rheumatoid Disease. In: Rheumatology,
an Annual Review. Vol. 1. Ed. Rotstein, J.. Karger Basel.

PAGE-THOMAS, D.P. (1969)

Lysosomal Enzymes in Experimental and Rheumatoid Arthritis. In:
Lysosomes in Biology and Pathology. Vol. 2. 877: Ed. Dingle, J.T.
and Fell, H.B.. North Holland. Amsterdam and London.

PANAYI, G.S.. (1976)

Antibody Mediated Leucocyte Cytotoxicity to Change Human Liver Cells in Rheumatoid Arthritis and Other Diseases. Ann. Rheum. Dis. 35: 27.

PANAYI, G.S. and CORRIGAL, V.. (1977)

Functional Assay of Cytotoxic Lymphocytes Involved in Antibody Mediated Cytotoxicity in Normal and Rheumatoid Subjects. Ann. Rheum. Dis. 36: 257.

PAPAMICHAIL, M., BROWN, J.C. and HOLBOROW, E.J.. (1971)

Immunoglobulins on the Surface of Human Lymphocytes. Lancet (ii): 850

PARROTT, D.M.V. and De SOUSA, M.A.B.. (1971)

Thymus-Dependent and Thymus-Independent Populations: Origin, Migratory Patterns and Life-Span. Clin. Exp. Immunol. 8: 663.

PARROTT, D.M.V. and FERGUSON, A.. (1974)

Selective Migration of Lymphocytes Within the Mouse Small Intestine. Immunology, 26: 571.

PARWARESCH, M.R., DESAGA, J.F. and NEBELIN, K. (1973)

Die Kinetik der basophilen Granulozyten im peripheren Blut beim Kaninchen. Blut. 27: 33

PATY, J.G., SIENKNECHT, C.W., TOWNES, A.S., HANNISIAN, A.S. and
MASI, A.T.. (1974)

Impaired Cell-Mediated Immunity in SLE and RA. Arth. Rheum. 17:
324.

PAULUS, H.E., MACHLEDER, H.I., LEVINE, S., YU, D.T.Y. and
MACDONALD, N.S.. (1977)

Lymphocyte Involvement in Rheumatoid Arthritis. Studies During
Thoracic Duct Drainage. Arth. Rheum. 20(6): 1249.

PEKÁREK, J., VEPŘEKOVA, A and DOUTLIK, S.. (1972)

Activation of Peripheral Lymphocytes in the Course of Experimental
Allergic Encephalo-myelitis in the Guinea Pig. 2. Immun.-Forsch.
143: 139.

PEKIN, T.J. Jr. and ZVAIFLER, N.J.. (1964).

Haemolytic Complement in Synovial Fluid. J. Clin. Invest.
43: 1372.

PERLMANN, P. and HOLM, G. (1969)

Cytotoxic Effects of Lymphoid Cells in Vitro. Adv. Immunol. 11: 117

PEARSON, D.A., SHARP, J.T. and LIDSKY, M.D. (1976)

The Cytotoxicity of Leukocytes and Lymphocytes from Patients with
Rheumatoid Arthritis from Synovial Cells. J. Clin. Invest. 58:
690.

PHILLIPS, R.A. and MILLER, R.G.. (1970)

Physical separation of Haemopoietic Stem Cells from Cells
Causing Graft-versus- Host Disease. 1. Sedimentation
properties of Cells Causing Graft-versus-Hose Disease. J.
Immunol. 105: 1168

PLOTZ, P.H. and TALAL, N.. (1967)

Fractionation of Splenic Antibody-Forming Cells Fractionated
on Glass Bead Columns. Clin. Exp. Immunol. 5: 209

POLLIACK, A., FU, S.M., DOUGLAS, S.D., BENTWICK, Z., LAMPEN, N.
and De HARVEN, E.. (1974)

Scanning Electron Microscopy of Human Lymphocyte - Sheep
Erythrocyte Rosettes. J. Exp. Med. 140: 146

POLLIACK, A., LAMPEN, N., CLARKSON, B.D. and De HARVEN, E.,
BENTWICK, Z., SIEGAL, F.P. and KUNKEL, H.G.. (1973)

Identification of Human B and T lymphocytes by Scanning Electron
Microscopy. J. Exp. Med. 138: 607.

PRITCHARD, H. and MICKLEM, H.S.. (1972)

Immune Responses in Congenitally Thymus Less Mice. 1. Absence
of Response to Oxazolone. Clin. Exp. Immunol. 10: 151.

RAFF, M.C.. (1970).

Two Distinct Populations of Peripheral Lymphocytes in Mice
Distinguishable by Immunofluorescence. Immunol. 19: 637.

RAIDT, D.J., MISCHELL, R.I. and DUTTON, R.W.. (1968).

Cellular Events in the Immune Response. J. Exp. Med. 128: 681

RAWSON, A.J.. (1975)

Abnormalities of Peripheral Blood Lymphocytes in Rheumatoid
Arthritis. Rheum. Annual Rev. 6: 190.

RAWSON, A.J., ABELSON, N.M. and McCARTY, D.J.. (1961)

The Isohaemagglutinin Partition in Rheumatoid Arthritis.
Arth. Rheum. 4: 463.

RAWSON, A.J. and HUANG, T.C.. (1974)

Lymphocytes in Rheumatoid Arthritis. I. Responses to
Allogeneic Cells and to Phytoimitogens. Clin. Exp. Immunol.
16: 41.

RAWSON, A.J. and HUANG, T.C.. (1974)

Lymphocytes in Rheumatoid Arthritis. II. Quantitative Binding
of Phytoimitogens. Clin. Exp. Immunol. 16: 47.

RENOUX, G., RENOUX, M., TELLER, M.N., McMAHON, S. and

GUILLAUMIN, J.M.. (1976).

Potentiation of T Cell Mediated Immunity by Levamisole.
Clin. Exp. Immunol. 25: 288.

REYNOLDS, M.D. and ABDON, N.I. (1973)

Comparative study of the in vitro proliferative responses of blood and synovial fluid leukocytes of rheumatoid arthritis patients.

J. Clin. Invest. 52 : 1627.

RHODES, K., SCOTT, A., MARKHAM, R.L. and MONK-JONES, M. (1969)

Immunological sex differences: a study of patients with RA, their relatives and controls. Ann. Rheum. Dis. 28 : 104.

RIEKE, W.O. and SCHWARZ, M.R. (1967)

The proliferative and immunological potential of thoracic duct lymphocytes. "The lymphocyte in immunology and haemopoiesis".

Ed. J. Yoffey. p. 224. Edward Arnold, London.

ROBERTS, P.J. and HAYRY, P. (1977)

Effector mechanisms in allograft rejection II. Density, electrophoresis and size fractionation of allograft-infiltrating cells demonstrating several classes of killer cells. Cell Immunol.

30 : 236

ROCKLIN, R.E. (1973)

Production of migration inhibitory factor by non-dividing lymphocytes. J. Immunol. 110 : 673.

RODNAN, G.P. (1973)

ed: Primer on the Rheumatic Diseases. J.A.M.A. 224. Suppl.5

ROIT, I.M. (1976)

Essential Immunology. Blackwell. Oxford and London.

ROITT, I.M., CORBETT, M., FESTENSTEIN, H., PAPASTERIADIS, C.,
JARAQUEMADA, D., HAY, F.C. and NINEHAM, L.J. (1978)

The prognostic significance of DR⁴ in rheumatoid arthritis.

Presentation to Heberden Society Annual General Meeting.

ROOS, D. and LOOS, J.A. (1973)

Charges in the carbohydrate metabolism of mitogenically stimulated
human peripheral lymphocytes. II. Relative importance of glyco-
lysis and oxidative phosphorylation on phytohaemagglutinin
stimulation. *Expl. Cell Res.* 77, 127

ROSS, G.D., RABELLINO, E.M., POLLEY, M.J. and GREY, H.M. (1973)

Combined studies of complement receptor and surface immunoglobulin-
bearing cells and sheep erythrocyte rosette-forming cells in normal
and leukaemic human lymphocytes. *J. Clin. Invest.* 52 : 377.

ROSS, G.D., WINCHESTER, R.J., RABELLINO, E.M. and HOFFMAN, J. (1978)

Surface markers of complement receptor lymphocytes. *J. Clin. Invest.*
62: 1086.

RUNGE, L.A. (1976)

Search for antigen in rheumatoid synovial macrophages. *Ann. Rheum.*
Dis. 35 : 133.

RUSSELL, R.J., WILKINSON, P.C., SLESS, F. and PARROTT, D.M.V. (1975)

Chemotaxis of lymphoblasts. *Nature* 256 : 646.

SAGONE, A.L. (Jr.) and MURPHY, S.G. (1975)

The chronic lymphatic leukemia lymphocyte : correlation of functional,
metabolic and surface immunoglobulin characteristics. *Cell Immunol.* 18(i):1

SANDILANDS, G.P., GRAY, K., COONEY, A., BROWNING, J.D. and
ANDERSON, J.R. (1975)

Formation of auto-rosettes by peripheral blood lymphocytes.

Clin. exp. Immunol. 22 : 493

SANY, J., CLOT, J., FREITAS, M., CHARMASSON, E. and SERRE, H. (1977)

Antibody dependent cell cytotoxicity (ADCC) of rheumatoid arthritis
blood lymphocytes. Rev. Rhum (2) 1-6. Trans. from Rev. Rhum.

43 : 333.

SANY, J., CLOT, J., MASSIP, H., CHARMASSON, E. and SERRE, H. (1975)

Sous-populations lymphocytaires impliquées dans le phenomene des
rosettes rheumatoides. Rev. Rhum 42 : 309.

SATO, C. and KOJIMA, K. (1976)

Phytohaemagglutinin-induced change in the distribution of acidic
lugons in surface membrane of lymphoid cells and blocking of the
radiation effect. Exp. Cell Res. 98 : 90.

SCATCHARD, C., BATCHELDER, A.C. and BROWN, A. (1946)

Preparation and properties of serum proteins. VI. Osmotic
equilibria in solutions of serum albumin and sodium chloride

J. Amer. Chem. Soc. 68 : 2320.

SCHOOLEY, J.C. (1961)

Autoradiographic observations of plasma cell formation.

J. Immunol. 86 : 331.

SCHRADER, J.W. and FELDMANN, M. (1973)

261

The mechanism of antigenic competition. I. The macrophage as a site of a reversible block of T-B lymphocyte collaboration.

Eur. J. Immunol. 3 : 711.

SCHREK, R. (1963)

Cell transformations and mitoses produced in vitro. Am. Rev.

Respir. Dis. 87 : 734.

SCHUMACHER, H.R. and KITRIDOU, R.C. (1972)

Synovitis of recent onset. Arth. Rheum. 15 : 465.

SCOTT-LINTHICUM, D. and SELL, S. (1974)

A rabbit blood lymphocyte in both smooth and hairy. N. Engl. J. Med.

290 : 1327.

SEBOK, J., TALERMAN, A and WOUTERS, M.W. (1977)

Lymphocyte activation in rheumatoid synovial effusion.

Arth. Rheum 20 (8) 1481.

SELA, M., MOZES, E., SHEARER, G.M. and KARNIELY, Y. (1970)

Cellular aspects of the immune relationship between the net change of immunogens and of antibodies elicited. Proc. Nat. Acad.

Sci. 67 : 1288.

SELIGMAN, M., PREUD'HOMME, J.L. and BROUET, J.C. (1974)

B and T cell markers in human proliferative blood diseases and primary immunodeficiencies with special regard to membrane bound immunoglobulin. Transplant Rev. 16

SHARPIN, R.K.C. and WILSON, J.D. (1977a)

262

Increased EA rosette formation by lymphocytes from patients with rheumatoid arthritis. Clin. Exp. Immunol. 29 : 205.

SHARPIN, R.K.C. and WILSON, J.D. (1977b)

Studies on the nature of EA binding by lymphocytes from rheumatoid arthritis patients. Clin. Exp. Immunol. 29 : 213.

SHELDON, P.J. and HOLBOROW, E.J. (1975)

Demonstration of T and B lymphocytes in synovial membrane sections of patients with rheumatoid arthritis by the technique of cyto-adherence. Scand. J. Rheumatol. Suppl. 8.

SHORTMAN, K. (1966)

Separation of lymphocytes on glass bead columns. Aust.J. Exp. Biol. Med. Sci. 44 : 271.

SHORTMAN, K. (1968)

The separation of different cell classes from lymphoid organs.
II. The purification and analysis of lymphocyte populations by equilibrium density gradient centrifugation. Aust. J. Exp. Biol. Med. Sci. 46 : 375.

SHORTMAN, K. (1971)

The density distribution of thymus, thoracic duct and spleen lymphocytes. J. Cell. Physiol. 77 : 319.

SHORTMAN, K., HASKILL, J.S., SZENBERG, A. and LEGGE, D.G. (1967)

Density distribution analysis of lymphocyte populations.
Nature, 216 : 1227.

SHORTMAN, K. and JACKSON, H. (1974)

263

The differentiation of T-lymphocytes. 1. Proliferation kinetics and interrelationships of sub populations of mouse thymus cells.

Cell. Immunol. 12 : 230.

SHORTMAN, K. and SZENBERG, A. (1969)

The size and density distribution of fowl blood lymphocytes initiating a graft-versus-host reaction. Aust. J. Exp. Biol. Med.

Sci. 47 : 1

SHORTMAN, K., WILLIAMS, N., JACKSON, H., RUSSELL, P., BYRT, P. and DIENER, E. (1971)

The separation of different cell classes from lymphoid organs. IV.

The separation of lymphocytes from phagocytes on glass bead columns, and its effect on subpopulations of lymphocytes and antibody forming cells. J. Cell. Biol. 48 : 566.

SILVERMAN, H.A., JOHNSON, J.S., VAUGHAN, J.H. and McGLAMORY, J.C. (1976)

Altered lymphocyte reactivity in rheumatoid arthritis. Arth. Rheum. 19 : 509.

SKRAMKOVA, L. and FORTOVA, D. (1972)

Investigation of lymphocyte activation in peripheral blood in infectious diseases. Z. Immun. Forsch. 144 : 129.

SLIWINSKI, A.J., ZVAIFLER, N.J. (1970)

In vivo synthesis of IgG by the rheumatoid synovium. J. Lab. Clin. Med. 76 : 304.

SMETANA, K. and POTMESIL, M. (1970)

A further contribution on the incidence of ring shaped nucleoli and micronucleoli in mature human lymphocytes. *Folia Haematol.* 93 : 16.

SMILEY, J.D., SACHS, C. and ZIFF, M. (1968)

In vitro synthesis of immunoglobulin by rheumatoid synovial membrane. *J. Clin. Invest.* 47 : 624.

SMITH, J.W., STEINER, A.L., NEWBERRY, W.M. and PARKER, C.W. (1971)

Cyclic AMP in human lymphocytes. Alterations following PHA stimulation. *J. Clin. Invest.* 50 : 432.

SPECTOR, W.G. (1969)

The granulomatous inflammatory exudate. *Int. Rev. Exp. Pathol.* 8 : 1.

SPECTOR, W.G., WALTERS, M.N.I. and WILLOUGHBY, D.A. (1967)

A quantitative study of leucocytic emigration in chronic inflammatory granulomata. *J. Path. Bact.* 93 : 101.

SPLINTER, T.A.W. and REISS, M. (1974)

Separation of lymphoid line cells according to volume and density. *Exp. Cell. Res.* 89 : 343.

SPRENT, J. and MILLER, J.F.A.P. (1972a).

Interaction of thymus lymphocytes with histoincompatible cells.
I. Quantitation of the proliferative response of thymus cells.
Cell. Immunol. 3 : 361.

SPRENT, J. and MILLER, J.F.A.P. (1972b)

Interaction of thymus lymphocytes with histoincompatible cells.

II. Recirculating lymphocytes derived from antigen-activated thymus cells. *Cellular Immunology* 3 : 385.

SPRENT, J. and MILLER, J.F.A.P. (1972c)

Interaction of thymus lymphocytes with histoincompatible cells.

III Immunological characteristics of recirculating lymphocytes derived from activated thymus cells. *Cell. Immunol.* 3 : 213.

STASTNY, P. (1974)

• MLC typing cells from patients with RA. *Tissue Antigens.* 4 : 571.

STEINMAN, R.M., MACHTINGER, B.G., FRIED, J. and COHN, Z.A. (1978)

Mouse spleen lymphoblasts generated in vitro. Recovery in high yield and purity after flotation in dense bovine plasma albumin solutions. *J. Exp. Med.* 147 : 279.

STOBO, J.D., ROSENTHAL, A.S. and PAUL, W.E. (1972)

Functional heterogeneity of murine lymphoid cells. I. Responsiveness to and surface binding to Concanavalin A and phyto-haemagglutinin. *J. Immunol.* 108 : 1.

STOUT, R.D. and HERZENBERG, L.A. (1975a).

The Fc receptor on thymus-derived lymphocytes. I. Detection of a subpopulation of murine T lymphocytes bearing the Fc receptor.

J. Exp. Med. 142 : 611.

STOUT, R.D. and HERZENBERG, L.A. (1975b)

The Fc receptor on thymus-derived lymphocytes. II. Mitogen responsiveness of T lymphocytes bearing the Fc receptor. J. exp. Med. 142 : 1041.

STOUT, R.D., WAKSAL, S.D. and HERZENBERG, L.A. (1976)

The Fc receptor on thymus-derived lymphocytes. III. Mixed lymphocyte reactivity and cell-mediated lymphocytic activity of Fc^- and Fc^+ T lymphocytes. J. exp. Med. 144 : 54.

STRELKAUSKAS, A.J., WILSON, B.S., CALLERY, R.T., CHESS, L. and SCHLOSSMAN, S.F. (1977)

T-cell regulation of human peripheral blood B-cell responsiveness. J. Exp. Med. 146 : 1765.

STROBER, S. (1972)

Initiation of antibody responses by different classes of lymphocytes. V. Fundamental changes in the physiological characteristics of virgin thymus-independent ("B") lymphocytes and "B" memory cells. J. Exp. Med. 136 : 851.

STROBER, S. and DILLEY, J. (1973)

Biological characteristics of T and B memory lymphocytes in the rat. J. Exp. Med. 137 : 1275.

STUART, J. and SIMPSON, J.S. (1970)

Dehydrogenase enzyme cytochemistry of unfixed leucocytes. J. Clin. Path. 23 : 517.

SVEDMYR, E., WIZZELL, H. and JONDAL, M. (1974)

Sensitization of human lymphocytes against autologous or allogeneic lymphoblastoid cell lines: characteristics of the reactive cells.

Scand. J. Immunol. 3 : 499.

TAKAHASI, T., OLD, L.J., McINTRE, K.R. and BOYSE, E.A. (1971)

Immunoglobulin and other surface antigens of cells of the immune response. J. exp. Med. 134 : 815.

TANNENBAUM, J. and SCHUR, P.M. (1974)

The role of lymphocytes in rheumatic diseases. J.Rheumatol.1: 392.

TATEISHI, H., COOKE, T.D. and ZIFF, M. (1971)

Electron microscope studies of pannus-free cartilage in rheumatoid and experimental arthritis. Arth. Rheum. 14 : 417.

TENNANT, J.R. (1964)

Evaluation of the trypan blue technique for determination of cell viability. Transplantation,2 : 685.

TERASAKI, P.I., VREDEVOE, D.L. and MICKEY, M.R. (1967)

Serotyping for homotransplantation. Transplantation, 5 : 1057.

THOMAS, C.B. and CATER, D.B. (1972)

Technique of separating the lymph node cells of C57 BL mice allergized with EP8 ascites tumour into populations with a slow and fast mobility. Br. J. Exp.Path. 53 : 111.

THOMAS, D.B. and PHILLIPS, B. (1973)

268

Evidence for membrane antigen(s) specified for human B lymphoblasts.

Clin. exp. Immunol. 14 : 91.

TRAYCOFF, R.B., PASCUAL, E. and SCHUMACHER, H.R. (1976)

Mononuclear cells in human synovial fluid. Identification of lymphoblasts in rheumatoid arthritis. Arth. Rheum. 19 : 743.

TSE, H. and DUTTON, R.W. (1976)

Separation of helper and suppressor T lymphocytes on a Ficoll velocity sedimentation gradient. J. Exp. Med. 143 : 1199.

TURK, W. (1907)

Septische Erkrankungen bei Verkümmerng des Granulozytensystems. Wein. Klin. Wchnschr. 20 : 157.

UTSINGER, P.D. (1975)

Synovial fluid lymphocytes in rheumatoid arthritis. Arth. Rheum. 18 : 595.

UTSINGER, P.D. and BLUESTEIN, H.G. (1974)

Sequential studies of T and B cells in RA and SLE. J. Rheumatol, 1 (Suppl): 75.

VAN BOXEL, J. A. and PAGET, S. A. (1975).

Predominantly T cell infiltrate in rheumatoid membranes.

New Eng. J. Med. 293: 517

VAN BOXEL, J. A. and ROSENSTREICH, D. C. (1974).

Bindings of aggregated δ -globulin to activated T lymphocytes in the guinea pig. J. Exp. Med. 139: 1002

VAN DE PUTTE, L. B. A., MEIJER, C. J. L. M., LAFEVER, G. J. M., et al. (1976).

Lymphocytes in rheumatoid and non-rheumatoid synovial fluids. Non-specificity of high T cell and low B cell percentages.

Ann. Rheum. Dis. 35: 451

VAN DILLA, M. A., FULWYLER, M. J., and BOONE, I. U. (1967).

Volume distribution and separation of normal human leukocytes.

Proc. Soc. Exp. Biol. Med. 125: 367

VAN EWYK, W., BRONS, N. H. C., and ROZING, J. (1975).

Scanning electron microscopy of homing and recirculating lymphocyte populations. Cell. Immunol. 19: 245

VAN FURTH, R. (1970).

Origin and kinetics of monocytes and macrophages. Semin. Haematol.

7: 125

VAN FURTH, R., SCHUIT, H. R. C., and HIJMANS, W. (1965).

The immunological development of the human foetus. J. Exp. Med.

122: 1173

VAUGHAN, J. H., BARNETT, E. V., and SOBEL, M. V. (1968).

Intracytoplasmic inclusions of immunoglobulins in rheumatoid arthritis and other diseases. Arth. Rheum. 11: 125

VAUGHAN, J. H., JACOX, R. F., and NOELL, P. (1968).

Relation of intracytoplasmic inclusions in joint fluid leukocytes to anti- δ -globulins. Arth. Rheum. 11: 135

VERNON-ROBERTS, B. (1969).

Lymphocyte to macrophage transformation in the peritoneal cavity preceeding the mobilization of peritoneal macrophages to inflamed areas.

Nature 222: 1286

- VERNON-ROBERTS, B., CURREY, H. L. F., and PERRIN, J. (1974).
T and B cells in the blood and synovial fluid of rheumatoid patients. *Ann. Rheum. Dis.* 33: 430
- VITETTA, E. S., FORMAN, J., and KETTMAN, J. R. (1976).
Cell surface immunoglobulin. XVIII. Functional differences of B lymphocytes bearing different surface immunoglobulin isotypes. *J. Exp. Med.* 143: 1055
- VON BOEHMER, H., SHORTMAN, K. and NOSSAL, G. J. V. (1974).
The separation of different cell classes from lymphoid organs. X. Preparative electrophoretic separation of lymphocyte sub-populations from mouse spleen and thoracic duct lymph. *J. Cell. Physiol.* 83: 231
- VOSSEN, J. M. and HIJMANS, W. (1975).
Membrane associated determinants on bone marrow and blood lymphocytes in the paediatric age group and on foetal tissues. *Ann. N. Y. Acad. Sc.* 254: 262
- WAITHE, W. I., HATHAWAY, P. and HIRSCHHORN, K. (1971).
Protein synthesis in stimulated lymphocytes. I. Conditions and results of an assay system. *Clin. Exp. Immunol.* 9: 903
- WAITHE, W. I., and HIRSCHHORN, K. (1973).
The lymphocyte response to activators. In *Handbook of experimental immunology*. Vol. 2. Ed. Weir, D. M. Blackwell, London.
- WALDER, I. A. and LUNSETH, J. B. (1963).
A technique for separation of the cells of the gastric mucosa. *Proc. Soc. Exp. Biol. (N. Y.)* 112: 494
- WALLER, M. H., ELLMAN, M. and TOONE, E. C. (1966).
Brucella immunization in patients with seropositive and seronegative rheumatoid arthritis. *Acta. Rheum. Scand.* 12: 250
- WANG, T., MARQUARDT, G., FOKER, J. (1976).
Aerobic glycolysis during lymphocyte proliferation. *Nature* 261: 702
- WANGEL, A. and KLOCKARS, M. (1977).
Lymphocyte sub-populations in rheumatoid synovial tissue. *Ann. Rheum. Dis.* 36: 176

WATSON, J. (1975).

The influence of intracellular levels of cyclic nucleotides on cell proliferation and the induction of antibody synthesis. *J. Exp. Med.* 141: 97

WAXMAN, J., LOCKSHIN, M.D., SCHNAPP, J. J. and DENESON, I. N. (1973). Cellular immunity in rheumatic diseases. I. Rheumatoid arthritis. *Arth. Rheum.* 16: 499

WEISMANN, G. (1971).

Lysosomes and the mediation of tissue injury in arthritis. In: *Rheumatoid Arthritis*. Ed. W. Muller, H. G., Harwerth and K. Fehr. Academic Press. London and New York.

WETZEL, B., CANNON, G. B., ALEXANDER, E. L., ERICKSON, B. W., and WESTBROOK, E. W. (1974).

In: *Scanning Electron Microscopy, 1974*. Eds, Jahari, O., and Corvin, I. pp. 582-587. III Research Inst. Chicago, Illinois.

WHALEY, K., GLENN, A. C. A., MACSWEEN, R. N. M., DEODHAR, S., DICK, W. C., NUKI, G., WILLIAMSON, AND BUCHANAN, W. W. (1971). Immunological responses in Sjogren's Syndrome and rheumatoid arthritis. *Clin. Exp. Immunol* 9(b): 721

WHITEHOUSE, D. J., WHITEHOUSE, M. W., and PEARSON, C. M. (1969). Passive transfer of adjuvant-induced arthritis and allergic encephalomyelitis in rats using thoracic duct lymphocytes. *Nature* 224: 1322

WHITESIDE, T. and RABIN, B. S. (1976).

Surface immunoglobulin on activated human peripheral blood thymus-derived cells. *J. Clin. Invest.* 57: 762

WHO/IARC Sponsored Workshop on Human B and T cells. (1974).

Identification, enumeration and isolation of B and T lymphocytes from human peripheral blood. *Scand. J. Immunol.* 3: 521

WIGZELL, H., and MÄKELÄ, O. (1970).

Separation of normal and immune lymphoid cells by antigen coated columns. Antigen-binding characteristics of membrane antibodies as analysed by hapten protein conjugates. *J. Exp. Med.* 132: 110

WIIG, J. N. (1973).

Electrophoresis of lymphoid cells. Characterization of T- and B-lymphocytes and two populations of thymocytes in the mouse.

Acta. Pathol. Microbiol. Scand. (A) 236 (Suppl): 101

WILKINSON, P. C., PARROTT, D. M. V., RUSSELL, R. J. and SLESS, F. (1977).

Antigen-induced locomotor responses in lymphocytes. J. Exp. Med.

145: 1158

WILKINSON, P. C., ROBERTS, J. A., RUSSELL, R. J., and McLOUGHLIN, M.

(1976).

Chemotaxis of mitogen-activated human lymphocytes and the effects of membrane-active enzymes. Clin. Exp. Immunol. 25: 280

WILLIAMS, N., KRAFT, N. and SHORTMAN, K. (1972).

The separation of different cell classes from lymphoid organs.

VI. The effect of osmolarity of gradient media on the density distribution of cells. Immunology 22: 885

WILLIAMS, R. C. Jnr. (1975).

Blood and tissue distribution of lymphocytes in rheumatoid arthritis.

Rheum. Annual Rev. 6: 219

WILLIAMS, R. C. Jnr., DeBOARD, J. R., MELLBYE, O. J., MESSNER, R. P.

and LINDSTRÖM, F. D. (1973).

Studies of T and B lymphocytes in patients with connective tissue diseases. J. Clin. Invest. 52: 283

WILSON, D. B. and NOWELL, P. C. (1971).

Quantitative studies on the mixed lymphocyte interaction in rats.

V. Tempo and specificity of the proliferative response and the number of reactive cells from immunized donors. J. Exp. Med.

133: 442

WINCHESTER, R. J., SEIGAL, F. P., HENTWICH, Z. H. and KUNKEL, H. G.

(1973).

Alteration in the proportion of B and T lymphocytes in rheumatoid arthritis joint fluids with low complement and increased complexes.

Arth. Rheum. 16: 138.

WISLOFF, F., FROLAND, S. S., and NATVIG, J. B. (1975).

Deficient lymphoid cell mediated, PHA induced cytotoxicity in rheumatoid arthritis patients. Scand. J. Immunol. 4: 303

WOOD, T. A. and FRENKEL, E. P. (1967).

The atypical lymphocyte. *Ann. J. Med.* 42: 923

WOOLEY, P. H. and PANAYI, G. S. (1978).

Studies of lymphocytes in rheumatoid arthritis. I.

Ann. Rheum. Dis. 37: 343

WOOLEY, P. H., PANAYI, G. S., and BATCHELOR, R. J. (1978).

HLA-DR antigens in rheumatoid arthritis. Presentation to Heberden Society annual general meeting.

WYBRAN, J., CARR, M. C., and FUDENBERG, H. H. (1972).

The human rosette-forming cell as a marker of a population of thymus-derived cells. *J. Clin. Invest.* 51: 2537

WYBRAN, J. and FUDENBERG, H. H. (1973).

Thymus derived rosette-forming cells in various human disease states: cancer, lymphoma, bacterial and viral infections, and other diseases. *J. Clin. Invest.* 52: 1026

YAM, L. T., LI, C. Y., and CROSBY, W. H. (1971).

Cytochemical identification of monocytes and granulocytes.

Amer. J. Clin. Path. 55: 283

YODOI, J., MIYAMA, M., and MASUDA, T. (1978).

Immunological properties of Fc receptors on lymphocytes. 2.

Differentiation from FCR- to FCR+ cells and their functional

differences in the in vitro antibody response. *Cell. Immunol.*

35: 266

YOSHIDA, T. O., and ANDERSSON, B. (1972).

Evidence for a receptor recognizing antigen complexed immunoglobulin on the surface of activated mouse thymus lymphocytes. *Scand.*

J. Immunol. 1: 401

YOSHIDA, Y., and OSMOND, D. G. (1971).

Identification and proliferation of small lymphocyte precursors

in cultures of lymphocyte-rich fractions of guinea-pig bone marrow.

Blood 37: 73

- YU, D. T. Y., CLEMENTS, P. J., PETER., J. B., et al (1974).
Study of lymphocyte characteristics in rheumatic patients
and the effect of azathioprine. *Arth. Rheum.* 17:37
- YU, D. T. Y., PETER, J. B., PAULUS, H. E. et al (1973).
Lymphocyte population: separation by discontinuous density
gradient centrifugation. *J. Immunol.* 110: 1615
- ZEIDERS, R. S., JOHNSON, J. S., EDGINGTON, T. S., and
VAUGHAN, J. H. (1973).
T lymphocytes in rheumatic diseases. *Arth. Rheum.* 16: 478
- ZETTERGREN, J. G., LUBEROFF, D. E. and PRETLOW, T. G., (1973).
Separating of lymphocytes from disaggregated mouse malignant
neoplasms by sedimentation in gradients of Ficoll in tissue
culture medium. *J. Immunol* 111: 836
- ZIFF, M. (1974).
Relation of cellular infiltration of rheumatoid synovial membrane
to its immune response. *Arth. and Rheum.* 17(3): 313
- ZVAIFLER, N. J. (1972).
In: *Inflammation (Mechanisms and Control)*. p 228. Eds. Lepar
I. H., and Ward, P. A. Academic Press, London and New York.
- ZVAIFLER, N. J. (1973).
The immunopathology of joint inflammation in R.A. *Advances
Immunology* 16: 265
- ZVAIFLER, N. J. (1974).
Rheumatoid synovitis. An extravascular immune complex disease.
Arth. and Rheum. 17 (3): 297